#### MARKERS FOR DETECTION OF GASTRIC CANCER

### Related Application

This application claims priority under 35 U.S.C. 119 to United States Provisional Patent Application Serial No: 60/487,906, filed July 17, 2003, titled "Markers for Detection of Gastric Cancer," listing Parry John Guilford as inventor. The above application is herein incorporated fully by reference.

#### Field of the Invention

This invention relates to detection of cancer. Specifically, this invention relates to the use of genetic and/or protein markers for detection of cancer, and more particularly to the use of genetic and/or protein markers for detection of gastric cancer.

#### **BACKGROUND**

Survival of cancer patients is greatly enhanced when the cancer is detected and treated early. In the case of gastric cancer, patients diagnosed with early stage disease have 5-year survival rates of 90%, compared to approximately 10% for patients diagnosed with advanced disease. However, the vast majority of gastric cancer patients currently present with advanced disease. Therefore, developments that lead to early diagnosis of gastric cancer can lead to an improved prognosis for the patients.

Identification of specific cancer-associated markers in biological samples, including body fluids, for example, blood, urine, peritoneal washes and stool extracts can provide a valuable approach for the early diagnosis of cancer, leading to early treatment and improved prognosis. Specific cancer markers also can provide a means for monitoring disease progression, enabling the efficacy of surgical, radiotherapeutic and chemotherapeutic treatments to be tracked. However, for a number of major cancers, the available markers suffer from insufficient sensitivity and specificity. For example, the most frequently used markers for gastric cancer, ca19-9, ca72-4 and chorioembryonic antigen (CEA) detect only about 15-50% of gastric tumors of any stage, declining to approximately 2-11% for early stage disease. Thus, there is a very high frequency of false negative tests that can lead patients and health care practitioners to believe that no disease exists, whereas in fact, the patient may have

1

severe cancer that needs immediate attention. Moreover, these markers can give false positive signals in up to 1/3 of individuals affected by benign gastric disease.

### SUMMARY OF THE INVENTION

Thus, there is an acute need for better methods for detecting the presence of cancer. Aspects of this invention provide methods, compositions and devices that can provide for detection of early stage cancer, and decreasing the frequency of false positives and false negative test results.

In certain embodiments, molecular analysis can be used to identify genes that are over-expressed in gastric tumor tissue compared to non-malignant gastric tissue. Such analyses include microarray and quantitative polymerase chain reaction (qPCR) methods. Cancer genes and proteins encoded by those genes are herein termed gastric tumor markers (GTM). It is to be understood that the term GTM does not require that the marker be specific only for gastric tumors. Rather, expression of GTM can be increased in other types of tumors, including malignant or non-malignant tumors, including gastric, bladder, colorectal, pancreatic, ovarian, skin (e.g., melanomas), liver, esophageal, endometrial and brain cancers, among others. It should be understood, however that the term GTM does not include prior the art markers, ca19-9, ca72-4 and CEA. Some GTM are sufficiently over-expressed to be diagnostic of gastric cancer with a high degree of reliability, and in other cases, over-expression of two or more GTM can provide reliable diagnosis of gastric cancer.

In certain embodiments, microarray methods can be used to detect patterns of over-expression of one or more genes associated with cancer.

In other embodiments, quantitative polymerase chain reaction (qPCR) can be used to identify the presence of markers over expressed in tumor or other biological samples.

Some of the embodiments of GTM detection disclosed herein are over expressed in a highly selective fashion in tumor cells and little, if at all, in non-tumor cells, permitting sensitive and accurate detection of cancer with measurement of only one over expressed GTM. In other embodiments, over-expression of two, three or more GTM can be detected in a sample and can provide greater certainty of diagnosis.

Selected genes that encode proteins can be secreted by or cleaved from the cell. These proteins, either alone or in combination with each other, have utility as serum or body fluid markers for the diagnosis of gastric cancer or as markers for

monitoring the progression of established disease. Detection of protein markers can be carried out using methods known in the art, and include the use of monoclonal antibodies, polyclonal antisera and the like.

### BRIEF DESCRIPTION OF THE FIGURES

This invention is described with reference to specific embodiments thereof and with reference to the figures, in which:

Figure 1 depicts a table of markers and oligonucleotide sequences of markers for gastric cancer of this invention.

Figure 2 depicts a table of results obtained of studies carried out using microarray methods.

Figure 3 depicts a table of results obtained of studies carried out using quantitative PCR.

Figures 4a – 4d depict relationships between log2 fold results obtained using array and qPCR methods, in which the data is centered on the median normal for four gastric cancer markers. Grey squares correspond to non-malignant ("normal") samples and black triangles to tumor samples. Figure 4a: ASPN. Figure 4b: SPP1. Figure 4c: SPARC. Figure 4d: MMP12.

Figures 5a-5w depict histograms showing the relative frequency vs. log2 fold change data obtained from quantitative PCR studies of various tumor markers. Figure 5a: ASPN; Figure 5b: CST1,2 & 4; Figure 5c: CSPG2; Figure 5d: IGFBP7; Figure 5e: INHBA; Figure 5f: LOXL2; Figure 5g: LUM; Figure 5h: SFRP4; Figure 5i: SPARC; Figure 5j: SPP1; Figure 5k: THBS2; Figure 5l: TIMP1; Figure 5m: adlican; Figure 5n: PRS11; Figure 5o: ASAH1; Figure 5p: SFRP2; Figure 5q: GGH; Figure 5r: MMP12; Figure 5s: KLK10; Figure 5t: LEPRE1; Figure 5u: TG; Figure 5v: EFEMP2 and Figure 5w: TGFBI.

Figure 6 is a histogram showing the number of markers with a higher expression than the 95<sup>th</sup> percentile of the median normal expression. Results are based on qPCR data and are shown separately for each tumor sample.

Figures 7a- 7c depicts graphs that show relative log2 expression of the markers in individual tumor samples and non-malignant samples compared to the expression of the gene for the tumor marker, CEA. CEA is the serum marker currently most used to monitor progression of gastric cancer.

Figure 8 shows a table that complements Figure 3. Figure 8 summarizes expression levels determined by qPCR for the candidate tumor markers, but using the paired data (i.e., tumor ("T") and non-malignant ("N") samples from the same individual) to provide a T:N ratio. Figure 8 also includes additional markers not included in Figure 3, namely MMP2, CGR11, TGFB1, PCSK5, SERPINB5, SERPINH1. For comparison, the expression level of the established serum marker gene, CEACAM5 (CEA), is also shown. 27 of the 29 markers have a median T:N difference greater than or equal to CEA. Further, compared to CEA, 29/29 of the markers have a higher percentage of paired samples in which the expression in the tumor sample exceeds the expression in the normal sample. Three markers, CST1,2,44, ASPN and SFRP4 showed 100% discrimination between the paired tumor and normal samples. The gene sequences of these markers, and the location of the primers and probes used to detect them, are shown herein.

Figures 9a - 9d depict individual and median T:N fold change data for 29 gastric cancer markers in 40 patients with paired samples.

Figures 10a – 10ad depict graphs of tumor stage and log2 fold change in expression of CEA and other GTM of this invention. Figure 10a: adlican; Figure 10b: ASPN; Figure 10c: CSPG2; Figure 10d: CST1,2,4; Figure 10e:EFEMP2; Figure 10f: GGF; Figure 10g: INHBA; Figure 10h: IGFBP7; Figure 10i: KLK10; Figure 10j: LEPRE1; Figure 10k: LUM; Figure 10l: LOXL2; Figure 10m: MMP12; Figure 10n; TIMP1; Figure 10o: ASAH1; Figure 10p: SPP1; Figure 10q: SFRP2; Figure 10r: SFRP4; Figure 10s: SPARC; Figure 10t: PRSS11; Figure 10u: THBS2: Figure 10v: TG; Figure 10w: TGFBI; Figure 10x: CGR11; Figure 10y: SERPINH1; Figure 10z: MMP2; Figure 10aa: PCSK5; Figure 10ab: SERPINB5; Figure 10ac: TGFB1 and Figure 10ad: CEA (CEACAM5).

Figures 11a – 11ad depict graphs of tumor type (diffuse (D) or intestinal (I)) and log2 fold change in expression 29 GTM of this invention and CEA. Figure 11a: adlican; Figure 11b: ASPN; Figure 11c: CSPG2; Figure 11d: CST1,2,4; Figure 11e: EFEMP2; Figure 11f: GGH; Figure 11g: INHBA; Figure 11h: IGFBP7; Figure 11i: KLK10; Figure 11j: LEPRE1: Figure 11k: LUM; Figure 11l: LOXL2; Figure 11m: MMP12; Figure 11n: TIMP1; Figure 11o: ASAH1; Figure 11p: SPP1; Figure 11q: SFRP2; Figure 11r: SFRP4: Figure 11s; SPARC; Figure 11t: PRSS11: Figure 11u: THBS2; Figure 11v: TG; Figure 11w: TGFBI; Figure 11x: CGR11: Figure 11y:

SERPINH1; Figure 11z: MIMP2; Figure 11aa: PCSK5; Figure 11ab:SERPINB5; Figure 11ac: TGFB1 and Figure 11ad: CEA (CEACAM5).

Figure 12 depicts a three-dimensional graph showing 3 markers, SERPINH1, CST1,2,4 and INHBA, in a series of gastric tumor samples and non-malignant gastric samples.

Figure 13 depicts a table that shows the effect of multiple markers on the ability to accurately discriminate between tumor tissue and non-malignant tissue. The table has been derived from normal distributions derived from qPCR data.

Figure 14 is a Western blot of 4 tumor markers derived from tumor and non-tumor tissue.

Figure 15 is a Western blot of the tumor marker SPARC in gastric tumor tissue and in serum.

Figure 16 is an immunoblot depicting cystatin SN in the supernatant of a gastric cell line, AGS.

#### **DETAILED DESCRIPTION**

#### **Definitions**

Before describing embodiments of the invention in detail, it will be useful to provide some definitions of terms as used herein.

The term "GTM" or "gastric tumor marker" or "GTM family member" means a gene, gene fragment, RNA, RNA fragment, protein or protein fragment related or other identifying molecule associated with gastric cancer that does not include molecules that are known in the prior art to be associated with gastric cancer, ca19-9, ca72-4 and CEA. Examples of GTMs are included herein below.

The term "marker" means a molecule that is associated quantitatively or qualitatively with the presence of a biological phenomenon. Examples of "markers" are GTMs, however, "markers" also includes metabolites, byproducts, whether related directly or indirectly to a mechanism underlying a condition.

The term "qPCR" means quantitative polymerase chain reaction.

The term "expression" includes production of mRNA from a gene or portion of a gene, and includes the production of a protein encoded by an RNA or gene or portion of a gene, and includes appearance of a detection material associated with expression. For example, the binding of a binding ligand, such as an antibody, to a gene or other oligonucleotide, a protein or a protein fragment and the visualization of

the binding ligand is included within the scope of the term "expression." Thus, increased density of a spot on an immunoblot, such as a Western blot, is included within the term "expression" of the underlying biological molecule.

The term "CPN2" means human carboxypeptidase N, polypeptide 2, 83 kDa chain; and carboxypeptidase N.

The term "HAPLN4" means human hyaluronan glycoprotein link protein 4.

The term "MMP12" means human matrix metalloproteinase 12.

The term "INHBA" means human inhibin, beta A (also includes activin A, activin AB or alpha polypeptide).

The term "IGFBP7" means human insulin-like growth factor 7.

The term "GGH" means human gamma-glutamyl hydrolase (also known as conjugase, folylpolygammaglutamyl hydrolase).

The term "LEPRE1" means human leucine proline-enriched proteoglycan (also known as leprecan 1).

The term "CST4" means human cystatin S.

The term "SFRP4" means human secreted frizzled-related protein 4.

The term "ASPN" means human asporin (also known as LRR class 1).

The term "CGREF1" or "CGR11" means human cell growth regulator with EF hand domain 1.

The term "KLK" means either human kallikrein 10, variant 1 or human kallikrein 10, variant 2, or both, unless specified otherwise.

The term "TIMP1" means human tissue inhibitor of metalloproteinase 1 (also known as erythroid potentiating activity or collagenase inhibitor).

The term "SPARC" means human secreted protein, acidic, cysteine-rich (also known as osteonectin).

The term "TGFBI" means human transforming growth factor, beta-induced, 68kDa.

The term "EFEMP2" means human EGF-containing fibulin-like extracellular matrix protein 2.

The term "LUM" means human lumican.

The term "SNN" means human stannin.

The term "SPP1" means human secreted phosphoprotein 1 (also known as osteopontin, or bone sialoprotein I, or early T-lymphocyte activation 1).

The term "CSPG2" means human chondroitin sulfate proteoglycan 2 (also known as versican).

The term "ASAH1" means human N-acylsphingosine amidohydrolase, variant 1, or N-acylsphingosine amidohydrolase, variant 2, or both N-acylsphingosine amidohydrolase variants 1 and 2 (also known as acid ceramidase 1, variants 1 and 2).

The term "PRSS11" means human protease, serine, 11 (also known as IGF binding serine protease).

The term "SFRP2" means human secreted frizzled-related protein 2.

The term "PLA2G12B" means human phospholipase A2, group XIIB.

The term "SPON2" means human spondin 2, extracellular matrix protein.

The term "OLFM1" means human olfactomedin 1.

The term "TSRC1" means human thrombospondin repeat containing 1.

The term "THBS2" means human thrombospondin 2.

The term "adlican" means DKFZp564I1922.

The term "CST2" means human cystatin SA.

The term "CST1" means human cystatin SN.

The term "LOXL2" means human lysyl oxidase-like enzyme 2.

The term "TG" means human thyroglobulin.

The term "TGFB1" means human transforming growth factor, beta1.

The term "SERPINH1" means human serine or cysteine proteinase inhibitor clade H (also known as heat shock protein 47, member 1, or collagen binding protein 1).

The term "SERPINB5" means human serine or cysteine proteinase inhibitor, clade B (also known as ovalbumin, member 5).

The term "CEACAM5" or "CEA" means human carcinoembryonic antigenrelated cell adhesion molecule 5.

The term "MMP2" means human matrix metalloproteinase 2 (also known as gelatinase A, or 72 kDa gelatinase, or 72 kDa type IV collagenase).

The term "PCSK5" means human proprotein convertase subtilisin/kexin type 5.

It is to be understood that the above terms may refer to protein, DNA sequence and/or RNA sequence. It is also to be understood that the above terms also refer to non-human proteins, DNA and/or RNA having the same sequences as depicted herein.

## Description of Embodiments of the Invention

Markers for detection and evaluation of tumors including gastric cancer are provided that have a greater reliability in detecting gastric cancer than prior art markers. By the term "reliability" we include the absence of false positives and/or false negatives. Thus, with higher reliability of a marker, fewer false positives and/or false negatives are associated with diagnoses made using that marker Therefore, in certain embodiments, markers are provided that permit detection of gastric cancer with reliability greater than the reliability of prior art markers of about 50%. In other embodiments, markers are provided that have reliability greater than about 70%; in other embodiments, greater than about 73%, in still other embodiments, greater than about 80%, in yet further embodiments, greater than about 90%, in still others, greater than about 95%, in yet further embodiments greater than about 98%, and in certain embodiments, about 100% reliability.

Thus, we have surprisingly found numerous genes and proteins whose presence is associated with gastric tumors. Detection of gene products (e.g., oligonucleotides such as mRNA) and proteins and peptides translated from such oligonucleotides therefore can be used to diagnose tumors, such as gastric tumors. Array analysis of samples taken from patients with gastric tumors and from non-malignant tissues of the same subjects has led us to the surprising discovery that in many gastric tumors, specific patterns of over-expression of certain genes are associated with the disease.

Cancer markers can also be detected using antibodies raised against cancer markers.

By analyzing the presence and amounts of expression of a plurality of cancer markers can thus increase the sensitivity of diagnosis while decreasing the frequency of false positive and/or false negative results.

### General Approaches to Cancer Detection

The following approaches are non-limiting methods that can be used to detect cancer including gastric cancer using GTM family members.

 Microarray approaches using oligonucleotide probes selective for products of GTM genes.

• Real-time quantitative PCR (qPCR) on tumor samples and normal samples using marker specific primers and probes.

- Enzyme-linked immunological assays (ELISA).
- Immunohistochemistry using anti-marker antibodies on gastric tumors and lymph node metastases.
- Immunohistochemistry using anti-marker antibodies on other tumors including but not limited to colorectal, pancreatic, ovarian, melanoma, liver, esophageal, bladder, endometrial, and brain.
- Immunodetection of marker family members in sera from gastric cancer patients taken before and after surgery to remove the tumor.
- Immunodetection of marker family members in sera from healthy individuals and individuals with non-malignant diseases such as gastritis, ulceration, gastric metaplasia and dysplasia.
- Immunodetection of marker family members in patients with other cancers including but not limited to colorectal, pancreatic, ovarian, melanoma, liver, oesophageal, bladder, endometrial, and brain.
- Detection of markers in body fluids, including serum, lymph, peritoneal fluid, cerebrospinal fluid, synovial fluid and the like.
- Immunodetection of marker family members in gastric fluid, peritoneal washes, urine and stool from gastric cancer patients. Using array methods and/or qPCR.
- Analysis of array or qPCR data using computers. Primary data is collected and fold change analysis is performed by comparison of levels of gastric tumor gene expression with expression of the same genes in non-tumor tissue. A threshold for concluding that expression is increased is provided (e.g., 1.5 x increase, 2-fold increase, and in alternative embodiments, 3-fold increase, 4-fold increase or 5-fold increase). It can be appreciated that other thresholds for concluding that increased expression has occurred can be selected without departing from the scope of this invention. Further analysis of tumor gene expression includes matching those genes exhibiting increased expression with expression profiles of known gastric tumors to provide diagnosis of tumors.

In certain aspects, this invention provides methods for detecting cancer, comprising:

- (a) providing a biological sample; and
- (b) detecting the over expression of a GTM family member in said sample.

In other aspects, the invention includes a step of detecting over expression of GTM mRNA.

In other aspects, the invention includes a step of detecting over expression of a GTM protein.

In yet further aspects, the invention includes a step of detecting overexpression of a GTM peptide.

In still further aspects, the invention includes a device for detecting a GTM, comprising:

- a substrate having a GTM capture reagent thereon; and
- a detector associated with said substrate, said detector capable of detecting a GTM associated with said capture reagent, wherein the capture reagent includes an oligonucleotide or an antibody.

Additional aspects include kits for detecting cancer, comprising:

- a substrate:
- a GTM capture reagent, including one or more of a GTM-specific oligonucleotide and a GTM-specific antibody; and

instructions for use.

Yet further aspects of the invention include method for detecting a GTM using qPCR, comprising:

- a forward primer specific for said GTM;
- a reverse primer specific for said GTM;

PCR reagents;

a reaction vial; and

instructions for use.

Additional aspects of this invention comprise a kit for detecting the presence of a GTM protein or peptide, comprising:

a substrate having a capture agent for said GTM protein or peptide;

an antibody specific for said GTM protein or peptide;

a reagent capable of labeling bound antibody for said GTM protein or peptide;

and

instructions for use.

Additional aspects of this invention include a method for manufacturing a monoclonal antibody, comprising the steps of:

In yet further aspects, this invention includes a method for detecting gastric cancer, comprising the steps of:

providing a sample from a patient suspected of having gastric cancer; measuring the presence of a GTM protein using an ELISA method.

As described herein, detection of tumors can be accomplished by measuring expression of one or more tumor-specific markers. We have unexpectedly found that the association between increased expression of GTMs and the presence of diagnosed gastric cancer is extremely high. The least significant association detected had a p value of about  $1.6 \times 10^{-6}$ . Many of the associations were significant at p values of less than  $10^{-20}$ . With such a high significance, it may not be necessary to detect increased expression in more than one GTM. However, the redundancy in the GTMs of this invention can permit detection of gastric cancers with an increased reliability.

The methods provided herein also include assays of high sensitivity. qPCR is extremely sensitive, and can be used to detect gene products in very low copy number (e.g., 1-100) in a sample. With such sensitivity, very early detection of events that are associated with gastric cancer is made possible.

### Methods

The following general methods were used to evaluate the suitability of various approaches to molecular identification of markers associated with gastric tumors.

## **Tumor Collection**

Gastric tumor samples and non-malignant gastric tissues were collected from surgical specimens resected at Seoul National University Hospital, Korea and Dunedin Hospital, New Zealand. Diagnosis of gastric cancer was made on the basis of symptoms, physical findings and histological examination of tissues.

#### RNA Extraction

In some embodiments, expression of genes associated with gastric tumors was analyzed by determining the changes in RNA from samples taken from tumors. Frozen surgical specimens were embedded in OCT medium. 60µm sections were sliced from the tissue blocks using a microtome, homogenized in a TriReagent: water

(3:1) mix, then chloroform extracted. Total RNA was then purified from the aqueous phase using the RNeasy<sup>TM</sup> procedure (Qiagen). RNA was also extracted from 16 cancer cell lines and pooled to serve as a reference RNA.

## Microarray Slide Preparation

Epoxy coated glass slides were obtained from MWG Biotech AG, Ebersberg, Germany) and were printed with ~30,000 50mer oligonucleotides using a Gene Machines microarraying robot, according to the manufacturer's protocol. Reference numbers (MWG oligo #) for relevant oligonucleotides, and the NCBI mRNA and protein reference sequences are shown in Figure 2. Full DNA sequences of the GTM of this invention are shown herein below.

# RNA labeling and Hybridization

cDNA was transcribed from 10µg total RNA using Superscript II reverse transcriptase (Invitrogen) in reactions containing 5-(3-aminoallyl)- 2' deoxyuridine – 5'-triphosphate. The reaction was then de-ionized in a Microcon column before being incubated with Cy3 or Cy5 in bicarbonate buffer for 1 hour at room temperature. Unincorporated dyes were removed using a Qiaquick column (Qiagen) and the sample concentrated to 15ul in a SpeedVac. Cy3 and Cy5 labeled cDNAs were then mixed with Ambion ULTRAhyb buffer, denatured at 100°C for 2 minutes and hybridized to the microarray slides in hybridization chambers at 42°C for 16 hours. The slides were then washed and scanned twice in an Axon 4000A scanner at two power settings to yield primary fluorescence data on gene expression.

#### **Normalization Procedure**

To compare expression of cancer genes from tumors and non-cancerous tissues, median fluorescence intensities detected by Genepix<sup>TM</sup> software were corrected by subtraction of the local background fluorescence intensities. Spots with a background corrected intensity of less than zero were excluded. To facilitate normalization, intensity ratios and overall spot intensities were log-transformed. Log-transformed intensity ratios were corrected for dye and spatial bias using local regression implemented in the LOCFIT<sup>TM</sup> package. Log-transformed intensity ratios were regressed simultaneously with respect to overall spot intensity and location. The

residuals of the local regression provided the corrected log-fold changes. For quality control, ratios of each normalized microarray were plotted with respect to spot intensity and localization. The plots were subsequently visually inspected for possible remaining artifacts. Additionally, an analysis of variance (ANOVA) model was applied for the detection of pin-tip bias. All results and parameters of the normalization were inserted into a Postgres-database for statistical analysis.

# Statistical Analysis

Statistically significant changes in gene expression in tumor samples vs. normal tissues were identified by measured fold changes between arrays. To accomplish this, log2 (ratios) were scaled to have the same overall standard deviation per array. This standardization procedure reduced the average within-tissue class variability. The log2 (ratios) were further shifted to have a median value of zero for each oligonucleotide to facilitate visual inspection of results. A rank-test based on fold changes was then used to improve the noise robustness. This test consisted of two steps: (i) calculation of the rank of fold change (Rfc) within arrays and ii) subtraction of the median (Rfc) for normal tissue from the median(Rfc) for tumor tissue. The difference of both median ranks defines the score of the fold change rank presented in Figure 2. Two additional statistical tests were also performed on this standardized data: 1) Two sample student's t-test, with and without the Bonferroni adjustment and 2) the Wilcoxon test.

## Statistical Analysis of Marker Combinations

To determine the value of using combinations of two or three of the markers to discriminate between tumor and non-malignant samples, the qPCR data from 40 paired samples (tumor and non-malignant samples from the same patient) were subjected to the following analysis. Normal distributions for the non-malignant and tumor samples were generated using the sample means and standard deviations. The probability that values taken from the tumor expression data would exceed a defined threshold (e.g., greater than 50%, 70%, 73%, 80%, 90%, 95%, 98%, 99% or 100%) in the non-malignant distribution was then determined (i.e., sensitivity). For combinations of markers, the probability that at least one marker exceeded the threshold was determined.

(I)
Avollab

		Applied						Γ
	sombol	Blosystems "assay on demand" assay #	forward pelmer	Se di		<b>3</b> 83	1	ğaş
esportn (irr class 1)	ASPN		AAATACAAAAGGACACATTCAAAGGA	1	TOCTICISCAATICISATATISSA	23	TIGGAAATGAGTGCAAACCCTCTTGATAATAATG	\$
chondroltin sulfate proteoglycan 2 (versican)	CSPG2		GCCAGTGGAATGATGTTCCC	1	:TCTTGGCATTTTCTACAACAGGG	Z	AGGAACAGTTGCTTGGGGCCAGC	\$
cystatins SN, SA & S	CST1, 2, 4		AGTOCCAGOCCAACTTGGA	3	GGGAACTTCGTAGATCTGGAAAGA	25	AGCCAGAACTGCAGAAGAACAGTTGTGC	6
gamma-glutamyl hydrolase	HOO		GTGGCAATGCCGCTGAA	•	TEACAGCAACAACTCAGTAGGAAAA	×	TTCACTGGAGGTCAATTGCACAGCAGAAT	2
Insulin-like growth factor binding protein 7	IGFBP7		CAGGTCAGCAAGGCCACC		TCACAGCTCAAGTACACCTGGG	a	AGCAAGGTCTTCCATAGTGACGCC	8
kallikrein 10	KLK10		ACAACATGATATGTGCTGGACTGG	•	GAGAGGATGCCTTGGAGGGT	R	CITIGOCAGAIGACTCTGGAGGCCC	8
leudne profine-enriched proteoglycan 1(leprecan 1)	LEPRE1		CTTGAGTACAACGCTGACCTCTTC	_	CCSTGACACAGTICTGCTTACAG	20	CCATCACAGATOATTACATOCAGGTOCTCA	3
	WOT		GATTETTGTCCATAGTGCATETGC	•	CCAATCAATTCCAGGAAGAGA	ş	TAGGATTCAACATTTGCCAAAATGAGTCTAA	5
lysyl oxidasa-like 2	1000		AGGCCAGCTTCTIGCTTTGGA		CONTRACTOR			2
matrix metalloproteinase 12	MMP12		GCCTCTCTGCTGATGACATACGT	10	AGTGACAGCATCAAAACTCAAATTG	T	TCAGTCCCTGTATGCACACCCAAACACCA	z z
metalloproteinese inhibitor 1	TIMP		CCAGACCACCTTATACCAGOG	11	GGACCTGTGGAAGTATCCGC	22	CANGATGACCANGATGTATAAAGGGTTTCCAAGC	8
n-acylsphingosine amidohydrolase	ASAHI		CCCAGACGCTGCAA	12	ACAGGACATCATACATGGTTTCAAA	T-	TGTCTGAACCGCACCAACAATA	8
secreted fritzled-related protein 2	SFRP2		COCTAGCAGGACCACCT		TITIGCAGGCTTCACATACCTT		CTGCCAGCCACCGAGGAAGCTC	6
secreted protein, addic, cysteine rich	SPARC		TCTTCCCTGIACACTGGCAGTTC		GAAAAGCGGGTGGTCCA	~	TGGACCAGDACCCATTGACGG	58
serine protease 11 (IGF binding)	PRSS11		TCGGCAGGCCCTTAGTAA	15	AAGGAGATICCAGCIGICACTITIC		AGTOTTANTICCAATCACTTCACCGTCCAGG	8
thrombospondin 2	THBS2		TGGAAGGACTACAGGGCTATAG		TAGGITTGGTCATAGATAGGTCCTGAGT	**	AGGCCCAAGACCGGCTACATCAGAGTC	8
	,TG		GAGGETTCCTCGCAGTTCAA		TGTAAACCGCTCCACTTCACAT		TCTGGCAGATTCCGATGCCCCACAA	5
human cell growth regulator with EF hand domain 1	:CGR11		CTGCCCACCCTTCCA	18	TICIGIOCITICCIAGIOOCITIAGG	-7	CCAGGCAGGAGCAGCTCGG	Ø
human serine of cysteine proteinase inhibitor dade 8	SERPTINBS		TOCAGGCATTTTCCAGGATAA	61	AAGCGAATTTGCTAGTTGCA	=	TCACTOTACCONOCIA	Ę
transforming growth factor \$1	TGFB1		GGTCCATGTCATCACCAATGTT	R	TCTGCAAGTTCATCCCCTTT	Т	CAGCITICAGCCACACTCAGG	3
human proprotein convertase subtilisin/kexin type S	. PCSK5		AAAATCTTTGCCGGAAATGC ;	. 21	AGTCCTGGCCGTTGAAATACC		ACAGAATGTAGGGATGGGTTAAGCCTGCA	3
matrix metalloproteinase 2	MMP2		TTGATGGCATCGCTCAGATC	[ [	TGTCACGTGGCGTCACAGT		TICMGGACCGGTICATTIGGGG	8
human serine of cysteing omtalnase inhibitor clade H	SERPTHIA	Hs00241844 m1					•	
adilcan		Hs00377849_m1			·			
eof-containing fibuilin-like extracellular matrix protein 2 FFEW92	FEEMPS	H:00213545 m1						
secreted frizzled-related protein 4	SFRD4	Hs00180066_m1				T		T
Inhibin beta A chain	INHBA	Hs00170103_m1				Γ		T
osteopontin	SPP1	Hs00167093_m1						İ
transforming growth factor 8-induced	189	Hs00165908 mi						П
	1					T		1
			rigure 4			1		

Table 1

## Quantitative Real-Time PCR

In other embodiments, real-time, or quantitative PCR (qPCR) can be used for absolute or relative quantitation of PCR template copy number. Taqman<sup>TM</sup> probe and primer sets were designed using Primer Express V 2.0TM (Applied Biosystems). Where possible, all potential splice variants were included in the resulting amplicon, with amplicon preference given to regions covered by the MWG-Biotech-derived microarray oligonucleotide. Alternatively, if the target gene was represented by an Assay-on-Demand<sup>TM</sup> expression assay (Applied Biosystems) covering the desired amplicons, these were used. The name of the gene, symbol, the Applied Biosystems "assay on demand" number, forward primer, reverse primer and probe sequence used for qPCR are shown in Table 1 and in Figure 1. In the in-house designed assays, primer concentration was titrated using a SYBR green labeling protocol and cDNA made from the reference RNA. Amplification was carried out on an ABI Prism™ 7000 sequence detection system under standard cycling conditions. When single amplification products were observed in the dissociation curves, standard curves were generated over a 625-fold concentration range using optimal primer concentrations and 5'FAM - 3'TAMRA phosphate Taqman<sup>TM</sup> probe (Proligo) at a final concentration of 250nM. Assays giving standard curves with regression coefficients over 0.98 were used in subsequent assays. It can be appreciated that in other embodiments, regression coefficients need not be as high. Rather, any standard curve can be used so long as the regression coefficients are sufficiently high to permit statistically significant determination of differences in expression. Such regression coefficients may be above about 0.7, above about 0.8, above about 0.9 or above about 0.95 in alternative embodiments.

Assays were performed over two 96 well plates with each RNA sample represented by a single cDNA. Each plate contained a reference cDNA standard curve, over a 625-fold concentration range, in duplicate. Analysis consisted of calculating the ΔCT (target gene CT – mean reference cDNA CT). ΔCT is directly proportional to the negative log2 fold change. Log2 fold changes relative to the median non-malignant log2 fold change were then calculated (log2 fold change – median normal log2 fold change). These fold changes were then clustered into frequency classes and graphed.

# Microarray Analysis of Cancer Marker Genes

RNA from 58 gastric tumors and 58 non-malignant ("normal") gastric tissue samples were labeled with Cy5 and hybridized in duplicate or triplicate with Cy3 labeled reference RNA. After normalization, the change in expression in each of 29,718 genes was then estimated by three measures: (i) fold change: the ratio of the gene's median expression (un-standardized) in the tumor samples divided by the median level in the non-malignant samples. (ii) fold change rank and (iii) the statistical probability that the observed fold changes were significant.

# Selection of Serum Markers for Gastric Malignancy

In certain embodiments, the cancer marker can be found in biological fluids, including serum. Serum markers were selected from the array data based on (i) the presence of a signal sequence characteristic of secreted proteins or cleaved from the outside of the membrane, (ii) the median level of over-expression (fold change) in tumors compared to non-malignant controls, (iii) the median change in expression rank between tumors and non-malignant controls, and (iv) the degree of overlap between the ranges of expression in the tumor and the non-malignant controls.

All 29 GTMs are known to have a signal peptide sequence at the 5'end of their coding sequences. The signal sequence targets the GTM proteins for transport to an extracellular compartment through the plasma membrane (Gunner von Heijne, Journal of Molecular Biology 173:243-251 (1984). In addition, none of the GTMs have transmembrane sequence motifs that would result in the full-length protein being retained within the plasma membrane. Consequently, all of the GTM markers of this invention are likely to be secreted into the extracellular compartment, and therefore can be in contact with the vasculature, either being taken up by capillaries, or by being transported into the lymphatic system and then into the vasculature. As a result, each of these tumor-derived markers will be present in the blood.

Next, genes were excluded if >50% of the tumor samples showed expression levels within the 95<sup>th</sup> percentile of the non-malignant range. The variation in the degree of over-expression in the tumor samples reflects not only tumor heterogeneity but also variations in the extent of contamination of the tumor samples with "normal" tissue including muscle, stromal cells and non-malignant epithelial glands. This "normal" contamination ranged from 5 to 70% with a median of approximately 25%. Other genes were excluded because of high relative expression in hematopoietic cells,

or elevated expression in metaplastic gastric tissue. It can be appreciated that depending on the degree of contamination by normal cells or cells that normally express the marker, different threshold ranges can be selected that can provide sufficient separation between a cancer source and a normal source.

GTM that we have found to be useful include genes (DNA), complementary DNA (cDNA), RNA, proteins, and protein fragments of the following markers: carboxypeptidase N, polypeptide 2, 83 kDa chain (also known as carboxypeptidase N (CPN2), matrix metalloproteinase 12 (MMP12), inhibin ("INHBA"), insulin-like growth factor 7 ("IGFBP7"), gamma-glutamyl hydrolase ("GGH"), leucine prolineenriched proteoglycan ("LEPRE1"), cystatin S ("CST4"), secreted frizzled-related protein 4 ("SFRP4"), asporin ("ASPN"), cell growth regulator with EF hand domain 1 ("CGREF1"), kallikrein (KLK10), tissue inhibitor of metalloproteinase 1 ("TIMP1"), secreted acidic cysteine-rich protein ("SPARC"), transforming growth factor, βinduced ("TGFBI"), EGF-containing fibulin-like extracellular matrix protein 2 ("EFEMP2"), lumican ("LUM"), stannin ("SNN"), secreted phosphoprotein 1 ("SPP1"), chondroitin sulfate proteoglycan 2 ("CSPG2"), N-acylsphingosine amidohydrolase ("ASAH1"), serine protease 11 ("PRSS11"), secreted frizzled-related protein 2 ("SFRP2"), phospholipase A2, group XIIB ("PLA2G12B"), spondin 2, extracellular matrix protein ("SPON2"), olfactomedin 1 ("OLFM1"), thrombospondin repeat containing 1 ("TSRC1"), thrombospondin 2 ("THBS2"), adlican, cystatin SA ("CST2"), cystatin SN (CST1), lysyl oxidase-like enzyme 2 ("LOXL2"), thyroglobulin ("TG"), transforming growth factor beta1 ("TGFB1"), serine or cysteine proteinase inhibitor clade H ("SERPINH1"), serine or cysteine proteinase inhibitor clade B ("SERPINB5"), matrix metalloproteinase 2 ("MMP2"), proprotein convertase subtilisin/kexin type 5 ("PCSK5"), and hyalronan proteoglycan link protein 4 ("HAPLN4").

DNA sequences of GTM of this invention along with identifying information are shown herein below.

## Matrix Metalloproteinase 12

>gi|4505206|ref|NM\_002426.1| Homo sapiens matrix metalloproteinase 12 (macrophage elastase) (MMP12), mRNA | qPCR forward\_primer match [758..780] | qPCR reverse\_primer match [888..864] | qPCR probe match [786..815]

TAGAAGTTTACAATGAAGTTTCTTCTAATACTGCTCCTGCAGGCCA CTGCTTCTGGAGCTCTTCCCCTGAACAGCTCTACAAGCCTGGAAAAAAAT AATGTGCTATTTGGTGAGAGATACTTAGAAAAATTTTATGGCCTTGAGATA AACAAACTTCCAGTGACAAAAATGAAATATAGTGGAAACTTAATGAAGG AAAAAATCCAAGAAATGCAGCACTTCTTGGGTCTGAAAGTGACCGGGCAA CTGGACACATCTACCCTGGAGATGATGCACGCACCTCGATGTGGAGTCCC CGATCTCCATCATTTCAGGGAAATGCCAGGGGGGCCCGTATGGAGGAAAC ATTATATCACCTACAGAATCAATAATTACACACCTGACATGAACCGTGAG GATGTTGACTACGCAATCCGGAAAGCTTTCCAAGTATGGAGTAATGTTAC CCCCTTGAAATTCAGCAAGATTAACACAGGCATGGCTGACATTTTGGTGG TTTTTGCCCGTGGAGCTCATGGAGACTTCCATGCTTTTGATGGCAAAGGTG GAATCCTAGCCCATGCTTTTGGACCTGGATCTGGCATTGGAGGGGATGCA CATTTCGATGAGGACGAATTCTGGACTACACATTCAGGAGGCACAAACTT GTTCCTCACTGCTGTTCACGAGATTGGCCATTCCTTAGGTCTTGGCCATTCT AGTGATCCAAAGGCTGTAATGTTCCCCACCTACAAATATGTCGACATCAA CACATTTCGCCTCTCTGCTGATGACATACGTGGCATTCAGTCCCTGTATGG AGACCCAAAAGAGAACCAACGCTTGCCAAATCCTGACAATTCAGAACCAG CTCTCTGTGACCCCAATTTGAGTTTTGATGCTGTCACTACCGTGGGAAATA AGATCTTTTCTTCAAAGACAGGTTCTTCTGGCTGAAGGTTTCTGAGAGAC CAAAGACCAGTGTTAATTTAATTTCTTCCTTATGGCCAACCTTGCCATCTG GCATTGAAGCTGCTTATGAAATTGAAGCCAGAAATCAAGTTTTTCTTTTTA AAGATGACAAATACTGGTTAATTAGCAATTTAAGACCAGAGCCAAATTAT CCCAAGAGCATACATTCTTTTGGTTTTCCTAACTTTGTGAAAAAAATTGAT GCAGCTGTTTTTAACCCACGTTTTTATAGGACCTACTTCTTTGTAGATAAC CAGTATTGGAGGTATGAAAAGGAGACAGATGATGGACCCTGGTTATCC CAAACTGATTACCAAGAACTTCCAAGGAATCGGGCCTAAAATTGATGCAG TCTTCTATTCTAAAAACAAATACTACTATTTCTTCCAAGGATCTAACCAAT TTGAATATGACTTCCTACTCCAACGTATCACCAAAACACTGAAAAGCAAT AGCTGGTTTGGTTAGAAATGGTGTAATTAATGGTTTTTGTTAGTTCAC TTCAGCTTAATAAGTATTTATTGCATATTTGCTATGTCCTCAGTGTACCACT ACTTAGAGATATGTATCATAAAAATAAAATCTGTAAACCATAGGTAATGA TTATATAAAATACATAATATTTTCAATTTTGAAAACTCTAATTGTCCATTC TTGCTTGACTCTACTATTAAGTTTGAAAATAGTTACCTTCAAAGCAAGATA ATTCTATTTGAAGCATGCTCTGTAAGTTGCTTCCTAACATCCTTGGACTGA GAAATTATACTTACTTCTGGCATAACTAAAATTAAGTATATATTTTTGGC TCAAATAAAATTG **SEQ ID NO:67** 

#### Inhibin Beta A

>gi|4504698|ref|NM\_002192.1| Homo sapiens inhibin, beta A (activin A, activin AB alpha polypeptide) (INHBA), mRNA | qPCR assay\_on\_demand\_context match [457..481]

GAAAGCTTCATGTGGGCAAAGTCGGGGAGAACGGGTATGTGGAGATAGA GGATGACATTGGAAGGAGGCAGAAATGAATGAACTTATGGAGCAGACC TCGGAGATCATCACGTTTGCCGAGTCAGGAACAGCCAGGAAGACGCTGCA CTTCGAGATTTCCAAGGAAGGCAGTGACCTGTCAGTGGTGGAGCGTGCAG AAGTCTGGCTCTTCCTAAAAGTCCCCAAGGCCAACAGGACCAGGACCAAA GTCACCATCCGCCTCTTCCAGCAGCAGCAGCACCCGCAGGGCAGCTTGGA CACAGGGGAAGAGGCCGAGGAAGTGGGCTTAAAGGGGGAGAGGAGTGA ACTGTTGCTCTCTGAAAAAGTAGTAGACGCTCGGAAGAGCACCTGGCATG TCTTCCCTGTCTCCAGCAGCATCCAGCGGTTGCTGGACCAGGGCAAGAGC TCCCTGGACGTTCGGATTGCCTGTGAGCAGTGCCAGGAGAGTGGCGCCAG CTTGGTTCTCCTGGGCAAGAAGAAGAAGAAGAAGAGGGGGGGAAGGG AAAAAGAAGGCGGAGGTGAAGGTGGGGCAGGAGCAGATGAGGAAAAG GAGCAGTCGCACAGACCTTTCCTCATGCTGCAGGCCCGGCAGTCTGAAGA CCACCCTCATCGCCGGCGTCGGCGGGGCTTGGAGTGTGATGGCAAGGTCA ACATCTGCTGTAAGAAACAGTTCTTTGTCAGTTTCAAGGACATCGGCTGGA ATGACTGGATCATTGCTCCCTCTGGCTATCATGCCAACTACTGCGAGGGTG AGTGCCCGAGCCATATAGCAGGCACGTCCGGGTCCTCACTGTCCTTCCACT CAACAGTCATCAACCACTACCGCATGCGGGGCCATAGCCCCTTTGCCAAC CTCAAATCGTGCTGTGTGCCCACCAAGCTGAGACCCATGTCCATGTTGTAC TATGATGATGGTCAAAACATCATCAAAAAGGACATTCAGAACATGATCGT GGAGGAGTGTGGTGCTCATAGAGTTGCCCAGCCCAGGGGGAAAGGGAG CAAGAGTTGTCCAGAGAAGACAGTGGCAAAATGAAGAAATTTTTAAGGTT AAAAACAAAAAAAACAAAAGTAAATTAAAAACA:AACCTGATGAAACAG CTCAGAGATGAAGCAGTGAAGAGACAGATTGGGAGGGAAAGGGAGAATG GTGTACCCTTTATTTCTTCTGAAATCACACTGATGACATCAGTTGTTTAAA CGGGGTATTGTCCTTTCCCCCCTTGAGGTTCCCTTGTGAGCTTGAATCAAC CAATCTGATCTGCAGTAGTGTGGACTAGAACAACCCAAATAGCATCTAGA AAGCCATGAGTTTGAAAGGGCCCATCACAGGCACTTTCCTAGCCTAAT SEQ ID NO:68

# Insulin-Like Growth Factor Binding Protein 7

>gi|4504618|ref|NM\_001553.1| Homo sapiens insulin-like growth factor binding protein 7 (IGFBP7), mRNA | qPCR forward\_primer match [470..487] | qPCR reverse\_primer match [567..546] | qPCR probe match [492..517]

## Gamma-Glutamyl Hydrolase

>gi|4503986|ref|NM\_003878.1| Homo sapiens gamma-glutamyl hydrolase (conjugase, folylpolygammaglutamyl hydrolase) (GGH), mRNA | qPCR forward\_primer match [531..547] | qPCR reverse\_primer match [611..587] | qPCR probe match [549..577]

TGCCGCAGCCCCGCCGCCGCAGAGCTTTTGAAAGGCGGCGGG AGGCGCGAGCCCATGCCAGTCCGGGCTGCTGTGCGTGCTGGGC CTGCTACTCTGCGGGGCGGCGAGCCTCGAGCTGTCTAGACCCCACGGCGA CACCGCCAAGAAGCCCATCATCGGAATATTAATGCAAAAATGCCGTAATA AAGTCATGAAAAACTATGGAAGATACTATATTGCTGCGTCCTATGTAAAG TACTTGGAGTCTGCAGGTGCGAGAGTTGTACCAGTAAGGCTGGATCTTAC AGAGAAAGACTATGAAATACTTTTCAAATCTATTAATGGAATCCTTTTCCC TGGAGGAAGTGTTGACCTCAGACGCTCAGATTATGCTAAAGTGGCCAAAA TATTTATAACTTGTCCATACAGAGTTTTGATGATGGAGACTATTTTCCTGT GTGGGGCACATGCCTTGGATTTGAAGAGCTTTCACTGCTGATTAGTGGAG AGTGCTTATTAACTGCCACAGATACTGTTGACGTGGCAATGCCGCTGAACT TCACTGGAGGTCAATTGCACAGCAGAATGTTCCAGAATTTTCCTACTGAGT TGTTGCTGTCATTAGCAGTAGAACCTCTGACTGCCAATTTCCATAAGTGGA GCCTCTCCGTGAAGAATTTTACAATGAATGAAAAGTTAAAGAAGTTTTTC AATGTCTTAACTACAAATACAGATGGCAAGATTGAGTTTATTTCAACAAT GGAAGGATATAAGTATCCAGTATATGGTGTCCAGTGGCATCCAGAGAAAG CACCTTATGAGTGGAAGAATTTGGATGGCATTTCCCATGCACCTAATGCTG TGAAAACCGCATTTTATTTAGCAGAGTTTTTTGTTAATGAAGCTCGGAAAA ACAACCATCATTTTAAATCTGAATCTGAAGAGGAGAAAGCATTGATTTAT CAGTTCAGTCCAATTTATACTGGAAATATTTCTTCATTTCAGCAATGTTAC ATATTTGATTGAAAGTCTTCAATTTGTTAACAGAGCAAATTTGAATAATTC CATGATTAAACTGTTAGAATAACTTGCTACTCATGGCAAGATTAGGAAGT ACTATTATATAACATTAGATAATTAAATAGTGAGACATAAATAGAGTGC GAAATACAAAAAAAAAAAAAA **SEQ ID NO: 70** 

## Leucine Proline-Enriched Proteoglycan 1

>gi|21361917|ref|NM\_022356.2| Homo sapiens leucine proline-enriched proteoglycan (leprecan) 1 (LEPRE1), mRNA | qPCR forward\_primer match [813..836] | qPCR reverse\_primer match [894..872] | qPCR probe match [841..870]

GGTGGCGGGTGGCTGGCGGTTCCGTTAGGTCTGAGGGAGCGATGG CGGTACGCGCGTTGAAGCTGCTGACCACACTGCTGGCTGTCGTGGCCGCT GCCTCCCAAGCCGAGGTCGAGTCCGAGGCAGGATGGGGCATGGTGACGCC CCGGGGTGGTCCTGAGCATGGAACGGGCGCTGCGCTCCCGGGCAGCCCTC CGCGCCCTTCGCCTGCGCTGCCGCACCCAGTGTGCCGCCGACTTCCCGTGG GAGCTGGACCCGACTGGTCCCCAGCCCGGCCCAGGCCTCGGGCGCCGC CGCCTGCGCGACCTGAGCTTCTTCGGGGGCCTTCTGCGTCGCGCTGCCTG CCTGCGCCGCTGCCTCGGGCCGCCGCCCACTCGCTCAGCGAAGAGA TGGAGCTGGAGTTCCGCAAGCGGAGCCCCTACAACTACCTGCAGGTCGCC TACTTCAAGATCAACAAGTTGGAGAAAGCTGTTGCTGCAGCACACACCTT CTTCGTGGGCAATCCTGAGCACATGGAAATGCAGCAGAACCTAGACTATT ACCAAACCATGTCTGGAGTGAAGGAGGCCGACTTCAAGGATCTTGAGACT CAACCCCATATGCAAGAATTTCGACTGGGAGTGCGACTCTACTCAGAGGA ACAGCCACAGGAAGCTGTGCCCCACCTAGAGGCGGCGCTGCAAGAATACT TTGTGGCCTATGAGGAGTGCCGTGCCCTCTGCGAAGGGCCCTATGACTAC GATGGCTACAACTACCTTGAGTACAACGCTGACCTCTTCCAGGCCATCAC AGATCATTACATCCAGGTCCTCAACTGTAAGCAGAACTGTGTCACGGAGC TTGCTTCCCACCCAAGTCGAGAGAAGCCCTTTGAAGACTTCCTCCCATCGC ATTATAATTATCTGCAGTTTGCCTACTATAACATTGGGAATTATACACAGG CTGTTGAATGTGCCAAGACCTATCTTCTCTTCTTCCCCAATGACGAGGTGA TGAACCAAAATTTGGCCTATTATGCAGCTATGCTTGGAGAAGAACACACC AGATCCATCGGCCCCCGTGAGAGTGCCAAGGAGTACCGACAGCGAAGCCT ACTGGAAAAAGAACTGCTTTTCTTCGCTTATGATGTTTTTTGGAATTCCCTTT GTGGATCCGGATTCATGGACTCCAGGAGAAGTGATTCCCAAGAGATTGCA AGAGAAACAGAAGTCAGAACGGGAAACAGCCGTACGCATCTCCCAGGAG ATTGGGAACCTTATGAAGGAAATCGAGACCCTTGTGGAAGAGAAGACCA AGGAGTCACTGGATGTGAGCAGACTGACCCGGGAAGGTGGCCCCCTGCTG TATGAAGGCATCAGTCTCACCATGAACTCCAAACTCCTGAATGGTTCCCA GCGGGTGGTGATGGACGGCGTAATCTCTGACCACGAGTGTCAGGAGCTGC AGAGACTGACCAATGTGGCAGCAACCTCAGGAGATGGCTACCGGGGTCA GACCTCCCACATACTCCCAATGAAAAGTTCTATGGTGTCACTGTCTTCAA AGCCCTCAAGCTGGGGCAAGAAGCCAAAGTTCCTCTGCAGAGTGCCCACC TGTACTACAACGTGACGGAGAAGGTGCGGCGCATCATGGAGTCCTACTTC CGCCTGGATACGCCCCTCTACTTTTCCTACTCTCATCTGGTGTGCCGCACT GCCATCGAAGAGGTCCAGGCAGAGAGGAAGGATGATAGTCATCCAGTCC CCCCAGCCTACACCTTCCGCGACTACAGCGCCATCCTTTACCTAAATGGG GACTTCGATGGCGGAAACTTTTATTTCACTGAACTGGATGCCAAGACCGT GACGCAGAGGTGCAGCCTCAGTGTGGAAGAGCCGTGGGATTCTCTTCAG GCACTGAAAACCCACATGGAGTGAAGGCTGTCACCAGGGGGCAGCGCTGT AGCAGCTCGAGCGGTGAGAGCAGCTGGTGCTGTGGTGACCCGTTCCCAG

AGCGCCCTTGGTTTGCCTTTCTCTTCCCCAAATCCCATTGCCAGTGGCTGA GACACGAAAGGAGCACTTGGGACACCAGCTCCAACGCCCTGTCATTATGG TCACATTGCCTTGTCCTCCCTGGGCCTGCTGTGAACGGGATCCAGGTGGGG AAAGAGGTCAAGACAGGGAGCGATGCTGAGTTCTTGGTTCCCTCCTTGGG CCCCACTTCAGCTGTCCTTTTCCAGAGAGTAGGACCTGCTGGGAAGGAGA TGAGCCTGGGGCCATTAAGGAACCTTCCTTGTCCCCTGGGAAGTAGCAGC TGAGAGATAGCGAGTGTCTGGAGCGGAGGCCTCTCTGAATGGGCAGGGGT TTGTCCTTGCAGGACAGGGTGCAGGCAGATGACCTGGTGAAGATGCTCTT CAGCCCAGAAGAGATGGTCCTCTCCCAGGAGCAGCCCCTGGATGCCCAGC AGGCCCCCCGAACCTGCACAAGAGTCTCTCTCAGGCAGTGAATCGAAG CCCAAGGATGAGCTATGACAGCGTCCAGGTCAGACGGATGGGTGACTAGA CCCATGGAGAGGAACTCTTCTGCACTCTGAGCTGGCCAGCCCCTCGGGGC TGCAGAGCAGTGAGCCTACATCTGCCACTCAGCCGAGGGGACCCTGCTCA CAGCCTTCTACATGGTGCTACTGCTCTTGGAGTGGACATGACCAGACACC GCACCCCTGGATCTGGCTGAGGGCTCAGGACACAGGCCCAGCCACCCCC AGGGGCCTCCACAGGCCGCTGCATAACAGCGATACAGTACTTAAGTGTCT 

**SEQ ID NO: 71** 

#### Cystatin S

>gi|19882254|ref|NM\_001899.2| Homo sapiens cystatin S (CST4), mRNA | qPCR forward\_primer match [343..361] | qPCR reverse\_primer match [434..411] | qPCR probe match [382..410]

GGCTCTCACCCTCCTCCTGCAGCTCCAGCTTTGTGCTCTGCCTCT GAGGAGACCATGGCCCGGCCTCTGTGTACCCTGCTACTCCTGATGGCTACC CTGGCTGGGCTCTGGCCTCGAGCTCCAAGGAGAATAGGATAATCCC AGGTGGCATCTATGATGCAGACCTCAATGATGAGTGGGTACAGCGTGCCC TTCACTTCGCCATCAGCGAGTACAACAAGGCCACCGAAGATGAGTACTAC AGACGCCCGCTGCAGGTGCTGCGAGCCAGGGAGCAGACCTTTGGGGGGGT GAATTACTTCTTCGACGTAGAGGTGGGCCGCACCATATGTACCAAGTCCC AGCCCAACTTGGACACCTGTGCCTTCCATGAACAGCCAGAACTGCAGAAG AAACAGTTGTGCTCTTTCGAGATCTACGAAGTTCCCTGGGAGGACAGAAT GTCCCTGGTGAATTCCAGGTGTCAAGAAGCCTAGGGGTCTGTGCCAGGCC AGTCACACCGACCACCCACTCCCACCCACTGTAGTGCTCCCACCCCTG GACTGGTGGCCCCACCCTGCGGGAGGCCTCCCCATGTGCCTGTGCCAAG AGACAGACAGAGAAGGCTGCAGGAGTCCTTTGTTGCTCAGCAGGGCGCTC TGCCCTCCTTCCTTCTTGCTTCTAATAGACCTGGTACATGGTACACAC SEQ ID NO: 72 ACCCCCACCTCCTGCAATTAAACAGTAGCATCGCC

#### Secreted Frizzle-Related Protein 4

>gi|8400733|ref|NM\_003014.2| Homo sapiens secreted frizzled-related protein 4 (SFRP4), mRNA | qPCR assay\_on\_demand\_context match [1079..1103]

GGCGGGTTCGCGCCCCGAAGGCTGAGAGCTGGCGCTGCTCGTGCCC TGTGTGCCAGACGCGGAGCTCCGCGGCCGGACCCCGCGCCCCGCTTTG CTGCCGACTGGAGTTTGGGGGAAGAAACTCTCCTGCGCCCCAGAAGATTT CTTCCTCGGCGAAGGGACAGCGAAAGATGAGGGTGGCAGGAAGAGAGA CGCTTTCTGTCTGCCGGGGTCGCAGCGCGAGAGGGCAGTGCCATGTTCCTC

TCCATCCTAGTGGCGCTGTGCCTGTGGCTGCACCTGGCGCTGGGCGTGCGC GGCGCCCTGCGAGGCGGTGCGCATCCCTATGTGCCGGCACATGCCCTG GAACATCACGCGGATGCCCAACCACCTGCACCACAGCACGCAGGAGAAC GCCATCCTGGCCATCGAGCAGTACGAGGAGCTGGTGGACGTGAACTGCAG CGCCGTGCTGCGCTTCTTCTTCTGTGCCATGTACGCGCCCATTTGCACCCT GGAGTTCCTGCACGACCCTATCAAGCCGTGCAAGTCGGTGTGCCAACGCG CGCGCGACGACTGCGAGCCCCTCATGAAGATGTACAACCACAGCTGGCCC GAAAGCCTGCCTGCGACGAGCTGCCTGTCTATGACCGTGGCGTGTGCAT TTCGCCTGAAGCCATCGTCACGGACCTCCCGGAGGATGTTAAGTGGATAG ACATCACACCAGACATGATGGTACAGGAAAGGCCTCTTGATGTTGACTGT AAACGCCTAAGCCCCGATCGGTGCAAGTGTAAAAAGGTGAAGCCAACTTT GGCAACGTATCTCAGCAAAAACTACAGCTATGTTATTCATGCCAAAATAA AAGCTGTGCAGAGGAGTGGCTGCAATGAGGTCACAACGGTGGTGGATGTA AAAGAGATCTTCAAGTCCTCATCACCCATCCCTCGAACTCAAGTCCCGCTC ATTACAAATTCTTCTTGCCAGTGTCCACACATCCTGCCCCATCAAGATGTT CTCATCATGTGTTACGAGTGGCGTTCAAGGATGATGCTTCTTGAAAATTGC TTAGTTGAAAAATGGAGAGATCAGCTTAGTAAAAGATCCATACAGTGGGA AGAGAGGCTGCAGGAACAGCGGAGAACAGTTCAGGACAAGAAGAAAACA GCCGGGCGCACCAGTCGTAGTAATCCCCCCAAACCAAAGGGAAAGCCTCC TGCTCCCAAACCAGCCAGTCCCAAGAAGAACATTAAAACTAGGAGTGCCC AGAAGAGAACAAACCCGAAAAGAGTGTGAGCTAACTAGTTTCCAAAGCG GAGACTTCCGACTTCCTTACAGGATGAGGCTGGGCATTGCCTGGGACAGC CTATGTAAGGCCATGTGCCCCTTGCCCTAACAACTCACTGCAGTGCTCTTC ATAGACACATCTTGCAGCATTTTTCTTAAGGCTATGCTTCAGTTTTTCTTTG TAAGCCATCACAAGCCATAGTGGTAGGTTTGCCCTTTGGTACAGAAGGTG AGTTAAAGCTGGTGGAAAAGGCTTATTGCATTGCATTCAGAGTAACCTGT GTGCATACTCTAGAAGAGTAGGGAAAATAATGCTTGTTACAATTCGACCT TTTTTACAGTATGTTTTATTACCTTTTGATATCTGTTGTTGCAATGTTAGTG GGAATGAATGTTAAAAGATCTTTATGTGTTTATGGTCTGCAGAAGGATTTT TGTGATGAAAGGGGATTTTTTGAAAAATTAGAGAAGTAGCATATGGAAAA TTATAATGTGTTTTTTACCAATGACTTCAGTTTCTGTTTTTAGCTAGAAAC TTAAAAACAAAAATAATAATAAAGAAAAATAAATAAAAAGGAGAGGCAG ACAATGTCTGGATTCCTGTTTTTTGGTTACCTGATTTCCATGATCATGATGC TTCTTGTCAACACCCTCTTAAGCAGCACCAGAAACAGTGAGTTTGTCTGTA CCATTAGGAGTTAGGTACTAATTAGTTGGCTAATGCTCAAGTATTTTATAC CCACAAGAGAGGTATGTCACTCATCTTACTTCCCAGGACATCCACCCTGA GAATAATTTGACAAGCTTAAAAATGGCCTTCATGTGAGTGCCAAATTTTGT TTTTCTTCATTTAAATATTTTCTTTGCCTAAATACATGTGAGAGGAGTTAA ATATAAATGTACAGAGAGGAAAGTTGAGTTCCACCTCTGAAATGAGAATT ACTTGACAGTTGGGATACTTTAATCAGAAAAAAAGAACTTATTTGCAGCA AAACAATTTTATTGGCCTTTTGCTAACACAGTAAGCATGTATTTTATAAGG CATTCAATAAATGCACAACGCCCAAAGGAAATAAAATCCTATCTAATCCT ACTCTCCACTACACAGAGGTAATCACTATTAGTATTTTGGCATATTATTCT CCAGGTGTTTGCTTATGCACTTATAAAATGATTTGAACAAATAAAACTAG GAACCTGTATACATGTGTTTCATAACCTGCCTCCTTTGCTTGGCCCTTTATT GAGATAAGTTTTCCTGTCAAGAAAGCAGAAACCATCTCATTTCTAACAGC TGTGTTATATTCCATAGTATGCATTACTCAACAAACTGTTGTGCTATTGGA

TACTTAGGTGGTTTCTTCACTGACAATACTGAATAAACATCTCACCGGAAT
TC SEQ ID NO: 73

## Asporin

>gi|41350213|ref|NM\_017680.3| Homo sapiens asporin (LRR class 1) (ASPN), mRNA | qPCR forward\_primer match [798..823] | qPCR reverse\_primer match [934..912] | qPCR probe match [842..875]

AGTACTAACATGGACTAATCTGTGGGAGCAGTTTATTCCAGTATCA AAATGTAATACCTCCTCATCTTTTCTTCTTACACAGTGTCTGAGAACATTT ACATTATAGATAAGTAGTACATGGTGGATAACTTCTACTTTTAGGAGGACT ACTCTCTTCTGACAGTCCTAGACTGGTCTTCTACACTAAGACACCATGAAG GAGTATGTGCTCCTATTATTCCTGGCTTTGTGCTCTGCCAAACCCTTCTTTA GCCCTTCACACATCGCACTGAAGAATATGATGCTGAAGGATATGGAAGAC ACAGATGATGATGATGATGATGATGATGATGATGATGAGGA CAACTCTCTTTTCCAACAGAGAGCCAAGAAGCCATTTTTTCCATTTGA TCTGTTTCCAATGTGTCCATTTGGATGTCAGTGCTATTCACGAGTTGTACA TTGCTCAGATTTAGGTTTGACCTCAGTCCCAACCAACATTCCATTTGATAC TCGAATGCTTGATCTTCAAAACAATAAAATTAAGGAAATCAAAGAAAATG ATTTTAAAGGACTCACTTCACTTTATGGTCTGATCCTGAACAACAACAAGC TAACGAAGATTCACCCAAAAGCCTTTCTAACCACAAAGAAGTTGCGAAGG CTGTATCTGTCCCACAATCAACTAAGTGAAATACCACTTAATCTTCCCAAA TCATTAGCAGAACTCAGAATTCATGAAAATAAAGTTAAGAAAATACAAAA GGACACATTCAAAGGAATGAATGCTTTACACGTTTTGGAAATGAGTGCAA ACCCTCTTGATAATAATGGGATAGAGCCAGGGGCATTTGAAGGGGTGACG GTGTTCCATATCAGAATTGCAGAAGCAAAACTGACCTCAGTTCCTAAAGG CTTACCACCAACTTTATTGGAGCTTCACTTAGATTATAATAAAATTTCAAC AGTGGAACTTGAGGATTTTAAACGATACAAAGAACTACAAAGGCTGGGCC TAGGAAACAACAAATCACAGATATCGAAAATGGGAGTCTTGCTAACATA TTCAATTGCAAGAGTGGGAGTAAATGACTTCTGTCCAACAGTGCCAAAGA TGAAGAAATCTTTATACAGTGCAATAAGTTTATTCAACAACCCGGTGAAA TACTGGGAAATGCAACCTGCAACATTTCGTTGTGTTTTGAGCAGAATGAGT GTTCAGCTTGGGAACTTTGGAATGTAATAATTAGTAATTGGTAATGTCCAT TTAATATAAGATTCAAAAATCCCTACATTTGGAATACTTGAACTCTATTAA TAATGGTAGTATTATATATACAAGCAAATATCTATTCTCAAGTGGTAAGTC CACTGACTTATTTATGACAAGAAATTTCAACGGAATTTTGCCAAACTATT GATACATAAGGGTTGAGAGAAACAAGCATCTATTGCAGTTTCTTTTTGCGT ACAAATGATCTTACATAAATCTCATGCTTGACCATTCCTTTCTTCATAACA AAAAAGTAAGATATTCGGTATTTAACACTTTGTTATCAAGCATATTTTAAA AAGAACTGTACTGTAAATGGAATGCTTGACTTAGCAAAATTTGTGCTCTTT CATTTGCTGTTAGAAAAACAGAATTAACAAAGACAGTAATGTGAAGAGTG CATTACACTATTCTTATTCTTTAGTAACTTGGGTAGTACTGTAATATTTTTA ATCATCTTAAAGTATGATTTGATATAATCTTATTGAAATTACCTTATCATG TCTTAGAGCCCGTCTTTATGTTTAAAACTAATTTCTTAAAATAAAGCCTTC AGTAAATGTTCATTACCAACTTGATAAATGCTACTCATAAGAGCTGGTTTG GGGCTATAGCATATGCTTTTTTTTTTTAATTATTACCTGATTTAAAAAATCT

## Cell Growth Regulator with EF Hand Domain 1

>gi|33589823|ref|NM\_006569.2| Homo sapiens cell growth regulator with EF hand domain 1 (CGREF1), mRNA | qPCR forward\_primer match [378..394] | qPCR reverse primer match [455..431] | qPCR probe match [396..415]

CGCGCAGCCCTCCGGCCGCGGGCGCAGCGGGGGCGCTGGTGGAG CTGCGAAGGCCAGGTCCGGCGGCGGCGGCGGCGCTGGCACTGGCTCC GGACTCTGCCGGCCAGGGCGGCGCTCCAGCCGGGAGGGCGACGTGGA GCGCCACGTGGAGCGGCCCGGGGGAGGCTGGCGGGGGAGGCGAGGCG CGGGCGCGCAGCAGCAGCAGCGCCCACGAGCTGGACCCCCAGAGCC GCGCGCGCCGCAGCAGTTCCAGGAAGGATGTTACCTTTGACGATGACAG TGTTAATCCTGCTGCTGCTCCCCACGGGTCAGGCTGCCCCAAAGGATGGA GTCACAAGGCCAGACTCTGAAGTGCAGCATCAGCTCCTGCCCAACCCCTT CCAGCCAGGCCAGGAGCAGCTCGGACTTCTGCAGAGCTACCTAAAGGGAC TAGGAAGGACAGAAGTGCAACTGGAGCATCTGAGCCGGGAGCAGGTTCT CCTCTACCTCTTTGCCCTCCATGACTATGACCAGAGTGGACAGCTGGATGG CCTGGAGCTGCCATGTTGACAGCTGCTCTGGCCCCTGGAGCTGCCAA CTCTCCTACCACCCACCCGGTGATATTGATAGTGGACAAAGTGCTCGAGA CGCAGGACCTGAATGGGGATGGGCTCATGACCCCTGCTGAGCTCATCAAC TTCCCGGGAGTAGCCCTCAGGCACGTGGAGCCCCGGAGAGCCCCTTGCTCC ATCTCCTCAGGAGCCACAAGCTGTTGGAAGGCAGTCCCTATTAGCTAAAA GCCCATTAAGACAAGAAACACAGGAAGCCCCTGGTCCCAGAGAAGAAGC AAAGGCCAGGTAGAGGCCAGAAGGGAGTCTTTGGATCCTGTCCAGGAG CCTGGGGGCCAGGCAGAGGCTGATGGAGATGTTCCAGGGCCCAGAGGGG **AAGCTGAGGGCCAGGCAGAGGCTAAAGGAGATGCCCCTGGGCCCAGAGG** GGAAGCTGGGGGCCAGGCAGAGGCTGAAGGAGATGCCCCCGGGCCCAGA GGGGAAGCTGGGGGCCAGGCAGAGGCCAGGGAGAATGGAGAGGAGGCC AAGGAACTTCCAGGGGAAACACTGGAGTCTAAGAACACCCAAAATGACTT TGAGGTGCACATTGTTCAAGTGGAGAATGATGAGATCTAGATCTTGAAGA TACAGGTACCCCACGAAGTCTCAGTGCCAGAACATAAGCCCTGAAGTGGG CAGGGGAAATGTACGCTGGGACAAGGACCATCTCTGTGCCCCCTGTCTGG TCCCAGTAGGTATCAGGTCTTTCTGTGCAGCTCAGGGAGACCCTAAGTTAA GGGCAGATTACCAATAAAGAACTGAATGAATTCATCCCCCCGGGCCACC TCTCTACCCGTCCAGCCTGCCCAGACCCTCTCAGAGGAACGGGGTTGGGG ACCGAAAGGACAGGATGCCGCCTGCCCAGTGTTTCTGGGCCTCACGGTG CTCCGGCAGCAGAGCGCATGGTGCTAGCCATGGCCGGCTGCAGAGGACCC AGTGAGGAAAGCTCAGTCTATCCCTGGGCCCCAAACCCTCACCGGTTCCC CCTCACCTGGTGTTCAGACACCCCATGCTCTCCTGCAGCTCAGGGCAGGTG ACCCCATCCCAGTAATATTAATCATCACTAGAACTTTTTGAGAGCCTTGT ACACATCAGGCATCATGCTGGGCATTTTATATATGATTTTATCCTCACAAT

#### Kallikrein 10, Transcript Variant 1

>gi|22208981|ref|NM\_002776.3| Homo sapiens kallikrein 10 (KLK10), transcript variant 1, mRNA | qPCR forward\_primer match [851..874] | qPCR reverse\_primer match [950..931] | qPCR probe match [890..914]

CATCCTGCCACCCCTAGCCTTGCTGGGGACGTGAACCCTCTCCCCG CGCCTGGGAAGCCTTCTTGGCACCGGGACCCGGAGAATCCCCACGGAAGC CAGTTCCAAAAGGGATGAAAAGGGGGCGTTTCGGGCACTGGGAGAAGCC TGTATTCCAGGGCCCCTCCCAGAGCAGGAATCTGGGACCCAGGAGTGCCA GCCTCACCCACGCAGATCCTGGCCATGAGAGCTCCGCACCTCCACCTCTCC GCCGCCTCTGGCGCCCCGGGCTCTGGCGAAGCTGCTGCCGCTGCTGATGGC GCAACTCTGGGCCGCAGAGGCGCGCTGCTCCCCCAAAACGACACGCGCT TGGACCCGAAGCCTATGGCTCCCGTGCGCGCGCGCTCGCAGCCCTGG CAGGTCTCGCTCTTCAACGGCCTCTCGTTCCACTGCGCGGGTGTCCTGGTG GACCAGAGTTGGGTGCTGACGCCGCGCACTGCGGAAACAAGCCACTGTG GGCTCGAGTAGGGGATGACCACCTGCTGCTTCTTCAGGGAGAGCAGCTCC GCCGGACCACTCGCTCTGTTGTCCATCCCAAGTACCACCAGGGCTCAGGC CCCATCCTGCCAAGGCGAACGGATGAGCACGATCTCATGTTGCTGAAGCT GGCCAGGCCCGTAGTGCTGGGGCCCCGCGTCCGGGCCCTGCAGCTTCCCT ACGCCCCCGGAGAGTGAAGTACAACAAGGCCTGACCTGCTCCAGCAT CACTATCCTGAGCCCTAAAGAGTGTGAGGTCTTCTACCCTGGCGTGGTCAC CAACAACATGATATGTGCTGGACTGGACCGGGGCCAGGACCCTTGCCAGA GTGACTCTGGAGGCCCCCTGGTCTGTGACGAGACCCTCCAAGGCATCCTCT CGTGGGTGTTTACCCCTGTGGCTCTGCCCAGCATCCAGCTGTCTACACCC AGATCTGCAAATACATGTCCTGGATCAATAAAGTCATACGCTCCAACTGA TCCAGATGCTACGCTCCAGCTGATCCAGATGTTATGCTCCTGCTGATCCAG ATGCCCAGAGGCTCCATCGTCCATCCTCTCCCCCAGTCGGCTGAACTC TCCCCTTGTCTGCACTGTTCAAACCTCTGCCGCCCTCCACACCTCTAAACA TCTCCCCTCTCACCTCATTCCCCACCTATCCCCATTCTCTGCCTGTACTGA AGCTGAAATGCAGGAAGTGGTGGCAAAGGTTTATTCCAGAGAAGCCAGG AAGCCGGTCATCACCCAGCCTCTGAGAGCAGTTACTGGGGTCACCCAACC TGACTTCCTCTGCCACTCCTGCTGTGTGACTTTGGGCAAGCCAAGTGCCC TCTCTGAACCTCAGTTTCCTCATCTGCAAAATGGGAACAATGACGTGCCTA TAAAGGTTACCTGTTGTCGTGA **SEQ ID NO: 76** 

#### Kallikrein 10 Transcript Variant 2

>gi|22208983|ref|NM\_145888.1| Homo sapiens kallikrein 10 (KLK10), transcript variant 2, mRNA | qPCR forward\_primer match [714..737] | qPCR reverse\_primer match [813..794] | qPCR probe match [753..777]

ACCAGCGCAGACCACAGGCAGGCAGAGGCACGTCTGGGTCCCC TCCTCCTTCCTATCGGCGACTCCCAGGATCCTGGCCATGAGAGCTCCGCA CCTCACCTCTCGCCGCCTCTGGCGCCCGGGCTCTGGCGAAGCTGCTGCC GCTGCTGATGGCGCAACTCTGGGCCGCAGAGGCGGCGCTGCTCCCCCAAA TCGCAGCCTGGCAGGTCTCGCTCTTCAACGGCCTCTCGTTCCACTGCGCG GGTGTCCTGGTGGACCAGAGTTGGGTGCTGACGGCCGCGCACTGCGGAAA CAAGCCACTGTGGGCTCGAGTAGGGGATGACCACCTGCTGCTTCTTCAGG GAGAGCAGCTCCGCCGGACCACTCGCTCTGTTGTCCATCCCAAGTACCAC CAGGGCTCAGGCCCCATCCTGCCAAGGCGAACGGATGAGCACGATCTCAT GTTGCTGAAGCTGGCCAGGCCCGTAGTGCTGGGGCCCCGCGTCCGGGCCC TGCAGCTTCCCTACCGCTGTGCTCAGCCCGGAGACCAGTGCCAGGTTGCTG GCTGGGGCACCACGGCCGCCGGAGAGTGAAGTACAACAAGGGCCTGAC CTGCTCCAGCATCACTATCCTGAGCCCTAAAGAGTGTGAGGTCTTCTACCC ACCCTTGCCAGAGTGACTCTGGAGGCCCCCTGGTCTGTGACGAGACCCTC CAAGGCATCCTCTCGTGGGGTGTTTACCCCTGTGGCTCTGCCCAGCATCCA GCTGTCTACACCCAGATCTGCAAATACATGTCCTGGATCAATAAAGTCAT ACGCTCCAACTGATCCAGATGCTACGCTCCAGCTGATCCAGATGTTATGCT CCTGCTGATCCAGATGCCCAGAGGCTCCATCGTCCATCCTCTCCCCCA GTCGGCTGAACTCTCCCCTTGTCTGCACTGTTCAAACCTCTGCCGCCCTCC ACACCTCTAAACATCTCCCCTCTCACCTCATTCCCCCACCTATCCCCATTCT CTGCCTGTACTGAAGCTGAAATGCAGGAAGTGGTGGCAAAGGTTTATTCC AGAGAAGCCAGGAAGCCGGTCATCACCCAGCCTCTGAGAGCAGTTACTGG GGTCACCCAACCTGACTTCCTCTGCCACTCCTGTGTGACTTTGGGCA AGCCAAGTGCCCTCTGAACCTCAGTTTCCTCATCTGCAAAATGGGAACA ATGACGTGCCTACCTCTTAGACATGTTGTGAGGAGACTATGATATAACAT GTGTATGTAAATCTTCATGGTGATTGTCATGTAAGGCTTAACACAGTGGGT GGTGAGTTCTGACTAAAGGTTACCTGTTGTCGTGA **SEO ID NO: 77** 

### Tissue Inhibitor of Metalloproteinase 1

>gi|4507508|ref|NM\_003254.1| Homo sapiens tissue inhibitor of metalloproteinase 1 (erythroid potentiating activity, collagenase inhibitor) (TIMP1), mRNA | qPCR forward\_primer match [221..241] | qPCR reverse\_primer match [359..340] | qPCR probe match [251..283]

AGGGGCCTTAGCGTGCCGCATCGCCGAGATCCAGCGCCCAGAGAG
ACACCAGAGAACCCACCATGGCCCCCTTTGAGCCCCTGGCTTCTGGCATCC
TGTTGTTGCTGTGGCTGATAGCCCCCAGCAGGGCCTGCACCTGTGTCCCAC
CCCACCCACAGACGGCCTTCTGCAATTCCGACCTCGTCATCAGGGCCAAG
TTCGTGGGGACACCAGAAGTCAACCAGACCACCTTATACCAGCGTTATGA
GATCAAGATGACCAAGATGTATAAAAGGGTTCCAAGCCTTAGGGGATGCCG
CTGACATCCGGTTCGTCTACACCCCCGCCATGGAGAGTGTCTGCGGATACT
TCCACAGGTCCCACAACCGCAGCGAGGAGTTTCTCATTGCTGGAAAACTG
CAGGATGGACTCTTGCACATCACTACCTGCAGTTTCGTGGCTCCCTGGAAC
AGCCTGAGCTTAGCTCAGCGCCGGGGCTTCACCAAGACCTACACTGTTGG
CTGTGAGGAATGCACAGTGTTTCCCTGTTTATCCATCCCCTGCAAACTGCA
GAGTGGCACTCATTGCTTGTGGACGACCAGCTCCTCCAAGGCTCTGAAA

## Secreted Protein, Acidic, Cysteine-Rich

>gi|48675809|ref|NM\_003118.2| Homo sapiens secreted protein, acidic, cysteine-rich (osteonectin) (SPARC), mRNA | qPCR forward\_primer match [788..810] | qPCR reverse\_primer match [915..898] | qPCR probe match [818..839]

GTTGCCTGTCTCAAACCCCTCCACATTCCCGCGGTCCTTCAGACTG CCCGGAGAGCGCGCTCTGCCTGCCTGCCTGCCACTGAGGGTTCC CAGCACCATGAGGCCTGGATCTTCTTTTCTCCTTTGCCTGGCCGGGAGGGC CTTGGCAGCCCTCAGCAAGAAGCCCTGCCTGATGAGACAGAGGTGGTGG AAGAAACTGTGGCAGAGGTGACTGAGGTATCTGTGGGAGCTAATCCTGTC CAGGTGGAAGTAGGAGAATTTGATGATGGTGCAGAGGAAACCGAAGAGG AGGTGGTGGCGAAAATCCCTGCCAGAACCACCACTGCAAACACGGCAA GGTGTGCGAGCTGGATGAGAACAACACCCCCATGTGCGTGTGCCAGGACC CCACCAGCTGCCCAGCCCCATTGGCGAGTTTGAGAAGGTGTGCAGCAAT GACAACAAGACCTTCGACTCTTCCTGCCACTTCTTTGCCACAAAGTGCACC CTGGAGGCACCAAGAAGGCCACAAGCTCCACCTGGACTACATCGGGCC TTGCAAATACATCCCCCTTGCCTGGACTCTGAGCTGACCGAATTCCCCCT GCGCATGCGGACTGGCTCAAGAACGTCCTGGTCACCCTGTATGAGAGGG ATGAGGACAACAACCTTCTGACTGAGAAGCAGAAGCTGCGGGTGAAGAA GATCCATGAGAATGAGAAGCGCCTGGAGGCAGGAGACCACCCCGTGGAG CTGCTGGCCCGGGACTTCGAGAAGAACTATAACATGTACATCTTCCCTGTA CACTGGCAGTTCGGCCAGCTGGACCAGCACCCCATTGACGGGTACCTCTC CCACACCGAGCTGGCTCCACTGCGTGCTCCCCTCATCCCCATGGAGCATTG CACCACCGCTTTTTCGAGACCTGTGACCTGGACAATGACAAGTACATCG CCCTGGATGAGTGGGCCGGCTGCTTCGGCATCAAGCAGAAGGATATCGAC AAGGATCTTGTGATCTAAATCCACTCCTTCCACAGTACCGGATTCTCTCTT TAACCCTCCCTTCGTGTTTCCCCCAATGTTTAAAATGTTTGGATGGTTTGT TGTTCTGCCTGGAGACAAGGTGCTAACATAGATTTAAGTGAATACATTAA ACTTAACTATTAAGGCCTTTTCCACACGCATTAATAGTCCCATTTTTCTCTT GCCATTTGTAGCTTTGCCCATTGTCTTATTGGCACATGGGTGGACACGGAT CTGCTGGGCTCTGCCTTAAACACACATTGCAGCTTCAACTTTTCTCTTTAGT GTTCTGTTTGAAACTAATACTTACCGAGTCAGACTTTGTGTTCATTTCATTT CAGGGTCTTGGCTGCCTGTGGGCTTCCCCAGGTGGCCTGGAGGTGGGCAA AGGGAAGTAACAGACACACGATGTTGTCAAGGATGGTTTTGGGACTAGAG TGCCTGAGGCTGTAACTGAGAGAAAGATTCTGGGGCTGTGTTATGAAAAT ATAGACATTCTCACATAAGCCCAGTTCATCACCATTTCCTCCTTTACCTTTC AGTGCAGTTTCTTTTCACATTAGGCTGTTGGTTCAAACTTTTGGGAGCACG GACTGTCAGTTCTCTGGGAAGTGGTCAGCGCATCCTGCAGGGCTTCTCCTC CTCTGTCTTTTGGAGAACCAGGGCTCTTCTCAGGGGCTCTAGGGACTGCCA GGCTGTTTCAGCCAGGAAGGCCAAAATCAAGAGTGAGATGTAGAAAGTTG TAAAATAGAAAAAGTGGAGTTGGTGAATCGGTTGTTCTTTCCTCACATTTG GATGATTGTCATAAGGTTTTTAGCATGTTCCTCCTTTTCTTCACCCTCCCCT

TTTTCTTCTATTAATCAAGAGAAACTTCAAAGTTAATGGGATGGTCGGAT CTCACAGGCTGAGAACTCGTTCACCTCCAAGCATTTCATGAAAAAGCTGC TTCTTATTAATCATACAAACTCTCACCATGATGTGAAGAGTTTCACAAATC CTTCAAAATAAAAAGTAATGACTTAGAAACTGCCTTCCTGGGTGATTTGC ATGTGTCTTAGTCACCTTATTATCCTGACACAAAAACACATGAGC ATACATGTCTACACATGACTACACAAATGCAAACCTTTGCAAACACATTA TGCTTTTGCACACACACACCTGTACACACACCCGGCATGTTTATACACAG GGAGTGTATGGTTCCTGTAAGCACTAAGTTAGCTGTTTTCATTTAATGACC TGTGGTTTAACCCTTTTGATCACTACCACCATTATCAGCACCAGACTGAGC ATATTTATGATGTATTTACTCTGCACCAGGTCCCATGCCAAGCACTGGGGA CACAGTTATGGCAAAGTAGACAAAGCATTTGTTCATTTGGAGCTTAGAGT CCAGGAGGAATACATTAGATAATGACACAATCAAATATAAATTGCAAGAT GTCACAGGTGTGATGAAGGGAGAGTAGGAGAGACCATGAGTATGTGTAA CAGGAGGACACAGCATTATTCTAGTGCTGTACTGTTCCGTACGGCAGCCA CTACCCACATGTAACTTTTAAGATTTAAATTTAAATTAGTTAACATTCAA AACGCAGCTCCCCAATCACACTAGCAACATTTCAAGTGCTTGAGAGCCAT GCATGATTAGTGGTTACCCTATTGAATAGGTCAGAAGTAGAATCTTTTCAT CATCACAGAAAGTTCTATTGGACAGTGCTCTTCTAGATCATCATAAGACTA CAGAGCACTTTTCAAAGCTCATGCATGTTCATCATGTTAGTGTCGTATTTT GAGCTGGGGTTTTGAGACTCCCCTTAGAGATAGAGAAACAGACCCAAGAA ATGTGCTCAATTGCAATGGGCCACATACCTAGATCTCCAGATGTCATTTCC CCTCTCTTATTTTAAGTTATGTTAAGATTACTAAAACAATAAAAGCTCCTA **SEQ ID NO: 79** 

## Transforming Growth Factor, Beta-Induced

>gi|4507466|ref|NM\_000358.1| Homo sapiens transforming growth factor, beta-induced, 68kDa (TGFBI), mRNA | qPCR assay\_on\_demand\_context match [170..194]

GCTTGCCCGTCGGTCGCTAGCTCGCTCGGTGCGCGTCCTCC ATGGCGCTCTTCGTGCGGCTGCTGGCTCTCGCCCTGGCCCTGGGC CCCGCCGCGACCCTGGCGGGTCCCGCCAAGTCGCCCTACCAGCTGGTGCT GCAGCACAGCAGGCTCCGGGGCCGCCAGCACGCCCCAACGTGTGTGCTG TGCAGAAGGTTATTGGCACTAATAGGAAGTACTTCACCAACTGCAAGCAG TGGTACCAAAGGAAAATCTGTGGCAAATCAACAGTCATCAGCTACGAGTG CTGTCCTGGATATGAAAAGGTCCCTGGGGAGAAGGGCTGTCCAGCAGCCC TACCACTCTCAAACCTTTACGAGACCCTGGGAGTCGTTGGATCCACCACCA CTCAGCTGTACACGGACCGCACGGAGAAGCTGAGGCCTGAGATGGAGGG GCCGGCAGCTTCACCATCTTCGCCCCTAGCAACGAGGCCTGGGCCTCCTT GCCAGCTGAAGTGCTGGACTCCCTGGTCAGCAATGTCAACATTGAGCTGC CTGAAACACGGCATGACCCTCACCTCTATGTACCAGAATTCCAACATCCA GATCCACCACTATCCTAATGGGATTGTAACTGTGAACTGTGCCCGGCTCCT GAAAGCCGACCATGCAACCAACGGGTGGTGCACCTCATCGATAAGG TCATCTCCACCATCACCAACACATCAGCAGATCATTGAGATCGAGGAC ACCTTTGAGACCCTTCGGGCTGCTGTGGCTGCATCAGGGCTCAACACGAT GCTTGAAGGTAACGGCCAGTACACGCTTTTGGCCCCGACCAATGAGGCCT

TCGAGAAGATCCCTAGTGAGACTTTGAACCGTATCCTGGGCGACCCAGAA GCCTGAGAGACCTGCTGAACAACCACATCTTGAAGTCAGCTATGTGTGC TGAAGCCATCGTTGCGGGGCTGTCTGTAGAGACCCTGGAGGGCACGACAC TGGAGGTGGGCTGCAGCGGGACATGCTCACTATCAACGGGAAGGCGATC ATCTCCAATAAAGACATCCTAGCCACCAACGGGGTGATCCACTACATTGA TGAGCTACTCATCCCAGACTCAGCCAAGACACTATTTGAATTGGCTGCAG AGTCTGATGTCCACAGCCATTGACCTTTTCAGACAAGCCGGCCTCGGCA ATCATCTCTCTGGAAGTGAGCGGTTGACCCTCCTGGCTCCCCTGAATTCTG TATTCAAAGATGGAACCCCTCCAATTGATGCCCATACAAGGAATTTGCTTC GGAACCACATAATTAAAGACCAGCTGGCCTCTAAGTATCTGTACCATGGA CAGACCCTGGAAACTCTGGGCGGCAAAAAACTGAGAGTTTTTGTTTATCG TAATAGCCTCTGCATTGAGAACAGCTGCATCGCGGCCCACGACAAGAGGG GGAGGTACGGGACCCTGTTCACGATGGACCGGGTGCTGACCCCCCAATG GGGACTGTCATGGATGTCCTGAAGGGAGACAATCGCTTTAGCATGCTGGT AGCTGCCATCCAGTCTGCAGGACTGACGGAGACCCTCAACCGGGAAGGAG TCTACACAGTCTTTGCTCCCACAAATGAAGCCTTCCGAGCCCTGCCACCAA GAGAACGGAGCAGACTCTTGGGAGATGCCAAGGAACTTGCCAACATCCTG AAATACCACATTGGTGATGAAATCCTGGTTAGCGGAGGCATCGGGGCCCT GGTGCGCTAAAGTCTCTCCAAGGTGACAAGCTGGAAGTCAGCTTGAAAA ACAATGTGGTGAGTGTCAACAAGGAGCCTGTTGCCGAGCCTGACATCATG GCCACAAATGGCGTGGTCCATGTCATCACCAATGTTCTGCAGCCTCCAGCC AACAGACCTCAGGAAAGAGGGGATGAACTTGCAGACTCTGCGCTTGAGAT CTTCAAACAAGCATCAGCGTTTTCCAGGGCTTCCCAGAGGTCTGTGCGACT AGCCCCTGTCTATCAAAAGTTATTAGAGAGGATGAAGCATTAGCTTGAAG CACTACAGGAGGAATGCACCACGGCAGCTCTCCGCCAATTTCTCTCAGAT TTCCACAGAGACTGTTTGAATGTTTTCAAAACCAAGTATCACACTTTAATG TACATGGGCCGCACCATAATGAGATGTGAGCCTTGTGCATGTGGGGGAGG AGGGAGAGAGATGTACTTTTTAAATCATGTTCCCCCTAAACATGGCTGTTA ACCCACTGCATGCAGAAACTTGGATGTCACTGCCTGACATTCACTTCCAGA GAGGACCTATCCCAAATGTGGAATTGACTGCCTATGCCAAGTCCCTGGAA AAGGAGCTTCAGTATTGTGGGGCTCATAAAACATGAATCAAGCAATCCAG CCTCATGGGAAGTCCTGGCACAGTTTTTGTAAAGCCCTTGCACAGCTGGA GAAATGGCATCATTATAAGCTATGAGTTGAAATGTTCTGTCAAATGTGTCT CACATCTACACGTGGCTTGGAGGCTTTTATGGGGCCCTGTCCAGGTAGAA AAGAAATGGTATGTAGAGCTTAGATTTCCCTATTGTGACAGAGCCATGGT GTGTTTGTAATAATAAAACCAAAGAAACATA SEQ ID NO: 80

## EGF-Containing Fibulin-Like Extracellular Matrix Protein 2

>gi|8393298|ref|NM\_016938.1| Homo sapiens EGF-containing fibulin-like extracellular matrix protein 2 (EFEMP2), mRNA | qPCR assay\_on\_demand\_context match [1248..1272]

CATCAACGACCTACACGGCGAGGGACCCCGCCACCAGTGCCTCCCGCTC AACACCCCAACCCTGCCCACCAGGCTATGAGCCCGACGATCAGGACAGC TGTGTGGATGTGGACGAGTGTGCCCAGGCCCTGCACGACTGTCGCCCCAG CCAGGACTGCCATAACTTGCCTGGCTCCTATCAGTGCACCTGCCCTGATGG TTACCGCAAGATCGGGCCCGAGTGTGTGGACATAGACGAGTGCCGCTACC GCTACTGCCAGCACCGCTGCGTGAACCTGCCTGGCTCCTTCCGCTGCCAGT GCGAGCCGGGCTTCCAGCTGGGGCCTAACAACCGCTCCTGTGTTGATGTG AACGAGTGTGACATGGGGGCCCCATGCGAGCAGCGCTGCTTCAACTCCTA TGGGACCTTCCTGTGTCGCTGCCACCAGGGCTATGAGCTGCATCGGGATG GCTTCTCCTGCAGTGATATTGATGAGTGTAGCTACTCCAGCTACCTCTGTC AGTACCGCTGCGTCAACGAGCCAGGCCGTTTCTCCTGCCACTGCCCACAG GGTTACCAGCTGCTGGCCACACGCCTCTGCCAAGACATTGATGAGTGTGA GTCTGGTGCGCACCAGTGCTCCGAGGCCCAAACCTGTGTCAACTTCCATG GGGGCTACCGCTGCGTGGACACCAACCGCTGCGTGGAGCCCTACATCCAG GTCTCTGAGAACCGCTGTCTCTGCCCGGCCTCCAACCCTCTATGTCGAGAG CAGCCTTCATCCATTGTGCACCGCTACATGACCATCACCTCGGAGCGGAG AGTACCCGCTGACGTGTTCCAGATCCAGGCGACCTCCGTCTACCCCGGTGC CTACAATGCCTTTCAGATCCGTGCTGGAAACTCGCAGGGGGACTTTTACAT CGGGCCCCGGGAGTACGTGCTGGACCTGGAGATGGTCACCATGAATTCC CTCATGAGCTACCGGGCCAGCTCTGTACTGAGGCTCACCGTCTTTGTAGGG TAGCTGAGGAGCCTGTTGTGAGGGGCAGAATGAGAAAGGCCCAGGGGCC CCCATTGACAGGAGCTGGGAGCTCTGCACCACGAGCTTCAGTCACCCCGA GAGGAGAGGAGGTAACGAGGAGGGCGGACTCCAGGCCCCGGCCCAGAGA TTTGGACTTGGCTGCCTTGCAGGGGTCCTAAGAAACTCCACTCTGGACAG CGCCAGGAGGCCCTGGGTTCCATTCCTAACTCTGCCTCAAACTGTACATTT GGATAAGCCCTAGTAGTTCCCTGGGCCTGTTTTTCTATAAAACGAGGCAAC TGG **SEO ID NO: 81** 

#### Lumican

>gi|21359858|ref|NM\_002345.2| Homo sapiens lumican (LUM), mRNA | qPCR forward\_primer match [61..84] | qPCR reverse\_primer match [182..162] | qPCR probe match [117..152]

GTATCACTCAGAATCTGGCAGCCAGTTCCGTCCTGACAGAGTTCAC
AGCATATATTGGTGGATTCTTGTCCATAGTGCATCTGCTTTAAGAATTAAC
GAAAGCAGTGTCAAGACAGTAAGGATTCAAACCATTTGCCAAAAATGAGT
CTAAGTGCATTTACTCTCTCCTGGCATTGATTGGTGGTACCAGTGGCCAG
TACTATGATTATGATTTTCCCCTATCAATTTATGGGCAATCATCACCAAAC
TGTGCACCAGAATGTAACTGCCCTGAAAGCTACCCAAGTGCCATGTACTG
TGATGAGCTGAAATTGAAAAAGTGTACCAATGGTGCCTCCTGGAATCAAGT
ATCTTTACCTTAGGAATAACCAGATTGACCATATTGATGAAAAAGGCCTTTG
AGAATGTAACTGATCTGCAGTGGCTCATTCTAGATCACAACCTTCTAGAA
AACTCCAAGATAAAAGGGAGAGTTTTCTCTAAATTGAAACAACTGAAGAA
GCTGCATATAAACCACAACAACCTGACAGAGTCTGTGGGCCCACTTCCCA
AATCTCTGGAGGATCTGCAGCTTACTCATAACAAGATCACAAAGCTGGGC
TCTTTTGAAGGATTGGTAAACCTGACCTTCATCCATCTCCAGCACAATCGG
CTGAAAGAGGATGCTGTTTCAGCTGCTTTTAAATCACTCGAA

TACCTTGACTTGAGCTTCAATCAGATAGCCAGACTGCCTTCTGGTCTCCCT GTCTCTCTAACTCTCTACTTAGACAACAATAAGATCAGCAACATCCCT GATGAGTATTTCAAGCGTTTTAATGCATTGCAGTATCTGCGTTTATCTCAC AACGAACTGGCTGATAGTGGAATACCTGGAAATTCTTTCAATGTGTCATCC CTGGTTGAGCTGGATCTGTCCTATAACAAGCTTAAAAACATACCAACTGTC AATGAAAACCTTGAAAACTATTACCTGGAGGTCAATCAACTTGAGAAGTT TGACATAAGAGCTTCTGCAAGATCCTGGGGCCATTATCCTACTCCAAGA TCAAGCATTTGCGTTTGGATGGCAATCGCATCTCAGAAACCAGTCTTCCAC ATCTGTATCCTGGAACAATATTTTATGGTTATGTTTTTCTGTGTGTCAGTTT TCATAGTATCCATATTTTATTACTGTTTATTACTTCCATGAATTTTAAAATC TGAGGGAAATGTTTTGTAAACATTTATTTTTTTAAAGAAAAGATGAAAG GCAGGCCTATTCATCACAAGAACACACACATATACACGAATAGACATCA GATGTGCAAAACCTTTTACTGGTTGCATGGAAATCAGCCAAGTTTTATAAT CCTTAAATCTTAATGTTCCTCAAAGCTTGGATTAAATACATATGGATGTTA CTCTCTTGCACCAAATTATCTTGATACATTCAAATTTGTCTGGTTAAAAAA TAGGTGGTAGATATTGAGGCCAAGAATATTGCAAAATACATGAAGCTTCA TGCACTTAAAGAAGTATTTTTAGAATAAGAATTTGCATACTTACCTAGTGA ATTTGCATGTTATGTTTAATAAGCTACTAGCAAAATAAAACATAGCAAAT GAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA **SEO ID NO: 82** 

#### Stannin

>gi|29893560|ref|NM\_003498.3| Homo sapiens stannin (SNN), mRNA

AGCGGGGCCGGACCGGGCGGGCGGAGCCGGGGCCCGCGGGCTGCT GCGGGCGATCGGGCCGGGCCGCTGCCGCGCCATGGACTCCCGTGTCCAG CCTGAGTTCCAGCCTCACTGAGTGGCCACCCCCAAAGTGCTGCCAGCCGA GGAAGCCCCAGCACTGACCATGTCTATTATGGACCACAGCCCCACCACG GGCGTGGTCACAGTCATCGTCATCCTCATTGCCATCGCGGCCCTGGGGGCC TTGATCCTGGGCTGCTGCTGCTGCGGCTGCAGCGCATCAGCCAGTCA GAGGACGAGGAGCATCGTGGGGGATGGGGAGACCAAGGAACCCTTCC TGCTGGTGCAGTATTCGGCCAAGGGACCGTGCGTGGAGAGAAAGGCCAA GCTGATGACTCCCAACGGCCCGGAAGTCCACGGCTGAGCCAGGATGCAAG GCTCCTGGTCCTGTTTGCAGCCGGCCAAGAGGCGCTGGGAGGGGCAAAAC CATACGGATGCGCTGCTGTCTGAGAGGGAAGGGCTGACACTTGCTGGCATG GCCTCTGCGGGCTTCGTCATCGCATGCACTGATGCCCGGGGACCTGGCTGT CCTGGGCTTCCCCTCGGCCTCCAGGTGAGGCTGCCCATTGCAGGCACTGG GCAGGCCTGACCTTGCTGGGGCTCATGGCCCTGTAGCGCTTTTGTTACTTG AATGTCTAGCTGAGCCTGTTTTTGATGGAGCTACTACTGTAATGCGTGAAC TAACAAACCTGTGAACTGTAAATAGGCCCCTGGAAGCACGTGCTTAAGCC CTTTTGCTGATTTTTAAAAATATCATCTAGCGCACACGGGACTGGTATTCT GGCTGTACTAATGACAAGCTGAGTCAAGACCCTGGAGGGTCATAGGCTTG TAAAGGCCCACGCCACACTCGGCAGGGGTCTCTCATGTGTGTCCATCTGC GTGTATGTCAAGGAAGTGAGATGCCAATTTGGGGTCTTGAGGCTGACCAG TTGGGGTGCTTGGGTGATCTCTGCTTCATTAGTCATGGGTGGAAGAAAAA CCACACCCCCGCACCCCTCCGTTCTTTCTGCATAGACTCACTTGTTAAAT AGCAGTTCTGTTGAGAGTGGAGTTACTGCAGGGAAGCTACCGGACCTGCC TGGGAGCCAGTGAAGGCCGAGTCAGGGCACGCGTCCTGGAGGCTGCCAG

CGTCCTTGTAGCAGAGCAGTTTCTTGCCGCTTGGGTCTTCAGCACGCCAAG CCCCCACCACCTCCACCCGAGTGAAGGCTTCGCTGAAATTGCTTTGG TCCTCATAGAGCCTGTGGTGGCTACTTTTGGTCTGAAACCCACTTGGCCCA GGAAAGAGAAAAGGTTGTATGTTTTGTGTTTGGTGTTTCCTATTTTCTGCAC TGGAGGGGAGGGACTGTTGAGGTTCTGTCTTTTTCTTCTTTTCCTCTTCC CTCTTCACATCACTTGGCTTCCTTTCCTCTGATGACCGTCCGCCTATGGG GTTCTGACTTCACTTTCCTCAGCGGGTCTCCAGTCCCCTGACCCAGCTCTA AAGGCACTTAGGACCCAGGGAACATTCTCACGTGCACATTCCCCTAAGA GCCACCAGACTGCTTCCTGCCAGCCTGTGCTTGCGGCAGGGAGCCGGGGC AGGGCAGAGGTGAACTTGAAGTTCAGGACTTGACTCTCCCACAGGTGGTG AGCTGGTGGCTCTCTGGTGAGCTAGTGTCTCCACAGCCTGTCTCCAAGGCC TCCCCTATGTACATTTCAGTGAGCTCACTTTGATTTTTAATCCCACCACAA CTGTAATAGATGGAAGGTCAGCCCCGGCTTAACCACAGAGCACTGGCCCT TCATGGCTGAGCTCAGAGCTCTGGCCTCCTGCTCAGACTAAAGGCACCTCC CTGGAACCCGAGGCGGAGAAGTAGGGAGCTGTTCGTTTAAGCAGCATA CACCTAAATTGGGGGTTTAAACATTAAGTAGGAGCTTGGGGTGGAAGAGG GACAGCCGGCTGGGCCACCTGAGCAGAAGGTGGTAATGAAACACCTCAG CTGGGCTCTTGGGAGACCTTAGGAAGCAGGAGAGGCAACACCTCTGGCTA CTGATGGTGTGGCAAGTTCAGAAGAGGTGGTGGTGGGGTAGGCGTGATGT CAGCAGAAGCCCTGCAGGCTGGGTGGGCAGGACACGTGGTGGGGGCCAC TGAAACCAGGCCTAGGAGGGAGAACAAGTTCCAAAGGTGCCGACTGGAA GAAGGGGTAAAAGTTTGCTTTGGTGAGTGAGAAAAGGCTGGGGCGTGTG ATCCATCCCTCACGTTTCAGAACTTCCAGGCTTTCTACCTCGACTCTCAC CACAGCCAGCACATACACCTAGGCTGTTTTTCCTTCCTCCACACCTGAGGG ACGCAGCAACAGCTAGGATCTGCATTTTCAGGTTCCGAGCCTGACCCCTG GAACTGACCAGCGCTCGATTGTCAGCCTTGGCCTGGGGTTTTGACCTTGCC AGTGAAGTTTCGGTTTTGAAGTGATTAAATGTCACTTCCTCATCAGTTTCA CTTCTGGAGGTTTTCTTATCCTACTCCCTGGTGCCAGGGACGTACCTGGGA GTTTGAATCAGGCCCATTTGAGCGTGGCAGCCGTGTTGGGTGAAGGTCCG GGGCTCGGTGAGGCACTGGGGGGGGTTTTCGGGAGGAAAATGAAAATGCTT CTAGAATGAGTGAACCACATCATAGCTCTCACTGTTTTTTCAATAGCTACT TTTTTTAGCAGACACCAGAGCCACACTCAAATGGCTAAGTAGGTTATGAC CTCTCTGGATTATTTTTGAATGCCCAACTGTTGCATTCAAGTTTTCTGACTA ATAAGAAATTAAGCATTCATCCTTCGTATCACTGCAGAAGCAACAGTGGG GGCACAGGGAGGGAACTCTTGACACTGAGCCACTAAAATATGGACTAATT TTTTGGACAAATCTTCAAACGGACTGTGCTACTGTATTTGTCTCAAAGCTA CCAAGTTTGTGCAATAAGTGGAAGGGATGTCATCCTTCTTCAATAAATGCT AAAAAGAAAAAAAAAAA SEQ ID NO: 83

#### Secreted Phosphoprotein 1

>gi|38146097|ref|NM\_000582.2| Homo sapiens secreted phosphoprotein 1 (osteopontin, bone sialoprotein I, early T-lymphocyte activation 1) (SPP1), mRNA | qPCR assay on demand context match [253..277]

CTCCCTGTGTTGGTGGAGGATGTCTGCAGCAGCATTTAAATTCTGG GAGGGCTTGGTTGTCAGCAGCAGCAGGAGGAGGCAGAGCACAGCATCGT CGGGACCAGACTCGTCTCAGGCCAGTTGCAGCCTTCTCAGCCAAACGCCG ACCAAGGAAAACTCACTACCATGAGAATTGCAGTGATTTGCTTTTGCCTCC TAGGCATCACCTGTGCCATACCAGTTAAACAGGCTGATTCTGGAAGTTCTG AGGAAAAGCAGCTTTACAACAAATACCCAGATGCTGTGGCCACATGGCTA AACCCTGACCCATCTCAGAAGCAGAATCTCCTAGCCCCACAGACCCTTCC AAGTAAGTCCAACGAAAGCCATGACCACATGGATGATATGGATGATGAA GATGATGACCATGTGGACAGCCAGGACTCCATTGACTCGAACGACTC TGATGATGTAGATGACACTGATGATTCTCACCAGTCTGATGAGTCTCACCA TTCTGATGAATCTGATGAACTGGTCACTGATTTTCCCACGGACCTGCCAGC AACCGAAGTTTTCACTCCAGTTGTCCCCACAGTAGACACATATGATGGCC GAGGTGATAGTGTGGTTTATGGACTGAGGTCAAAATCTAAGAAGTTTCGC AGACCTGACATCCAGTACCCTGATGCTACAGACGAGGACATCACCTCACA CATGGAAAGCGAGGAGTTGAATGGTGCATACAAGGCCATCCCCGTTGCCC AGGACCTGAACGCGCCTTCTGATTGGGACAGCCGTGGGAAGGACAGTTAT GAAACGAGTCAGCTGGATGACCAGAGTGCTGAAACCCACAGCCACAAGC AGTCCAGATTATATAAGCGGAAAGCCAATGATGAGAGCAATGAGCATTCC GATGTGATTGATAGTCAGGAACTTTCCAAAGTCAGCCGTGAATTCCACAG CCATGAATTTCACAGCCATGAAGATATGCTGGTTGTAGACCCCAAAAGTA AGGAAGAAGATAAACACCTGAAATTTCGTATTTCTCATGAATTAGATAGT GCATCTTCTGAGGTCAATTAAAAGGAGAAAAAATACAATTTCTCACTTTG CATTTAGTCAAAAGAAAAATGCTTTATAGCAAAATGAAAGAGAACATGA AATGCTTCTTCTCAGTTTATTGGTTGAATGTGTATCTATTTGAGTCTGGAA ATAACTAATGTGTTTGATAATTAGTTTAGTTTGTGGCTTCATGGAAACTCC CTGTAAACTAAAAGCTTCAGGGTTATGTCTATGTTCATTCTATAGAAGAAA TGCAAACTATCACTGTATTTTAATATTTGTTATTCTCTCATGAATAGAAATT TATGTAGAAGCAAACAAAATACTTTTACCCACTTAAAAAGAGAATATAAC ATTTTATGTCACTATAATCTTTTGTTTTTTAAGTTAGTGTATATTTTGTTGT GATTATCTTTTTGTGGTGTGAATAAATCTTTTATCTTGAATGTAATAAGAA TTTGGTGTCAATTGCTTATTTGTTTTCCCACGGTTGTCCAGCAATTAAT AAAAAAAA **SEQ ID NO: 84** 

#### **Chondroitin Sulfate Proteoglycan 2**

>gi|21361115|ref|NM\_004385.2| Homo sapiens chondroitin sulfate proteoglycan 2 (versican) (CSPG2), mRNA | qPCR forward\_primer match [10087..10106] | qPCR reverse\_primer match [10185..10163] | qPCR probe match [10139..10161]

GCTGCCCGAGCCTTTCTGGGGAAGAACTCCAGGCGTGCGGACGCA ACAGCCGAGAACATTAGGTGTTGTGGACAGGAGCTGGGACCAAGATCTTC GGCCAGCCCGCATCCTCCGCATCTTCCAGCACCGTCCCGCACCCTCCGC ATCCTTCCCCGGGCCACCACGCTTCCTATGTGACCCGCCTGGGCAACGCCG AACCCAGTCGCGCAGCGCTGCAGTGAATTTTCCCCCCAAACTGCAATAAG CCGCCTTCCAAGGCCAAGATGTTCATAAATATAAAGAGCATCTTATGGAT GTGTTCAACCTTAATAGTAACCCATGCGCTACATAAAGTCAAAGTGGGAA AAAGCCCACCGGTGAGGGGCTCCCTCTCTGGAAAAGTCAGCCTACCTTGT

CATTITICAACGATGCCTACTTTGCCACCCAGTTACAACACCAGTGAATTT CTCCGCATCAAATGGTCTAAGATTGAAGTGGACAAAAATGGAAAAGATTT GAAAGAGACTACTGTCCTTGTGGCCCAAAATGGAAATATCAAGATTGGTC AGGACTACAAAGGGAGAGTGTCTGTGCCCACACATCCCGAGGCTGTGGGC GATGCCTCCCTCACTGTGGTCAAGCTGCTGGCAAGTGATGCGGGTCTTTAC CGCTGTGACGTCATGTACGGGATTGAAGACACACAAGACACGGTGTCACT GACTGTGGATGGGTTGTTTTCACTACAGGGCGCAACCAGCAGGTACA CACTGAATTTTGAGGCTGCTCAGAAGGCTTGTTTGGACGTTGGGGCAGTC ATAGCAACTCCAGAGCAGCTCTTTGCTGCCTATGAAGATGGATTTGAGCA GTGTGACGCAGGCTGGCTGATCAGACTGTCAGATATCCCATCCGGG CTCCCAGAGTAGGCTGTTATGGAGATAAGATGGGAAAAGGCAGGAGTCAG GACTTATGGATTCCGTTCTCCCCAGGAAACTTACGATGTGTATTGTTATGT GGATCATCTGGATGGTGATGTTTCCACCTCACTGTCCCCAGTAAATTCAC CTTCGAGGAGGCTGCAAAAGAGTGTGAAAACCAGGATGCCAGGCTGGCA ACAGTGGGGGAACTCCAGGCGGCATGGAGGAACGGCTTTGACCAGTGCG ATTACGGGTGGCTGTCGGATGCCAGCGTGCGCCACCCTGTGACTGTGGCC AGGGCCCAGTGTGGAGGTGGTCTACTTGGGGTGAGAACCCTGTATCGTTT TGAGAACCAGACAGGCTTCCCTCCCCTGATAGCAGATTTGATGCCTACTG CTTTAAACCTAAAGAGGCTACAACCATCGATTTGAGTATCCTCGCAGAAA CTGCATCACCCAGTTTATCCAAAGAACCACAAATGGTTTCTGATAGAACT ACACCAATCATCCCTTTAGTTGATGAATTACCTGTCATTCCAACAGAGTTC CCTCCCGTGGGAAATATTGTCAGTTTTGAACAGAAAGCCACAGTCCAACC TCAGGCTATCACAGATAGTTTAGCCACCAAATTACCCACACCTACTGGCA GTACCAAGAAGCCCTGGGATATGGATGACTACTCACCTTCTGCTTCAGGA CCTCTTGGAAAGCTAGACATATCAGAAATTAAGGAAGAAGTGCTCCAGAG TACAACTGGCGTCTCTCATTATGCTACGGATTCATGGGATGGTGTCGTGGA AGATAAACAAACACAAGAATCGGTTACACAGATTGAACAAATAGAAGTG TTCCCTGTAACTGAAACACCATTGGTAACTGCAAGAATGATCCTGGAATC CAAAACTGAAAAGAAAATGGTAAGCACTGTTTCTGAATTGGTAACCACAG GTCACTATGGATTCACCTTGGGAGAAGAGGATGATGAAGACAGAACACTT ACAGTTGGATCTGATGAGAGCACCTTGATCTTTGACCAAATTCCTGAAGTC ATTACGGTGTCAAAGACTTCAGAAGACACCATCCACACTCATTTAGAAGA CTTGGAGTCAGTCTCAGCATCCACAACTGTTTCCCCTTTAATTATGCCTGA TAATAATGGATCATCCATGGATGACTGGGAAGAGAGACAAACTAGTGGTA GGATAACGGAAGAGTTTCTTGGCAAATATCTGTCTACTACACCTTTTCCAT CACAGCATCGTACAGAAATAGAATTGTTTCCTTATTCTGGTGATAAAATAT TAGTAGAGGGAATTTCCACAGTTATTTATCCTTCTCTACAAACAGAAATGA CACATAGAAGAAGAACAGAAACACTAATACCAGAGATGAGAACAGA TACTTATACAGATGAAATACAAGAAGAGATCACTAAAAGTCCATTTATGG GAAAACAGAAGAAGTCTTCTCTGGGATGAAACTCTCTACATCTCTC TCAGAGCCAATTCATGTTACAGAGTCTTCTGTGGAAATGACCAAGTCTTTT GATTTCCCAACATTGATAACAAAGTTAAGTGCAGAGCCAACAGAAGTAAG AGATATGGAGGAAGACTTTACAGCAACTCCAGGTACTACAAAATATGATG AAAATATTACAACAGTGCTTTTGGCCCATGGTACTTTAAGTGTTGAAGCAG CCACTGTATCAAAATGGTCATGGGATGAAGATAATACAACATCCAAGCCT TTAGAGTCTACAGAACCTTCAGCCTCTTCAAAATTGCCCCCTGCCTTACTC ACAACTGTGGGGATGAATGGAAAGGATAAAGACATCCCAAGTTTCACTGA AGATGGAGCAGATGAATTTACTCTTATTCCAGATAGTACTCAAAAGCAGT TAGAGGAGGTTACTGATGAAGACATAGCAGCCCATGGAAAATTCACAATT

AGATTTCAGCCAACTACATCAACTGGTATTGCAGAAAAGTCAACTTTGAG AGATTCTACAACTGAAGAAAAAGTTCCACCTATCACAAGCACTGAAGGCC AAGTTTATGCAACCATGGAAGGAAGTGCTTTGGGTGAAGTAGAAGATGTG GACCTCTCTAAGCCAGTATCTACTGTTCCCCAATTTGCACACACTTCAGAG GTGGAAGGATTAGCATTTGTTAGTTATAGTAGCACCCAAGAGCCTACTAC TTATGTAGACTCTTCCCATACCATTCCTCTTTCTGTAATTCCCAAGACAGA CTGGGGAGTGTTAGTACCTTCTGTTCCATCAGAAGATGAAGTTCTAGGTGA ACCCTCTCAAGACATACTTGTCATTGATCAGACTCGCCTTGAAGCGACTAT TTCTCCAGAAACTATGAGAACAACAAAAATCACAGAGGGAACAACTCAG GAAGAATTCCCTTGGAAAGAACAGACTGCAGAGAAACCAGTTCCTGCTCT CAGTTCTACAGCTTGGACTCCCAAGGAGGCAGTAACACCACTGGATGAAC AAGAGGGCGATGGATCAGCATATACAGTCTCTGAAGATGAATTGTTGACA GGTTCTGAGAGGGTCCCAGTTTTAGAAACAACTCCAGTTGGAAAAATTGA TCACAGTGTGTCTTATCCACCAGGTGCTGTAACTGAGCACAAAGTGAAAA CAGATGAAGTGGTAACACTAACACCACGCATTGGGCCAAAAGTATCTTTA AGTCCAGGGCCTGAACAAAATATGAAACAGAAGGTAGTAGTACAACAG GATTTACATCATCTTTGAGTCCTTTTAGTACCCACATTACCCAGCTTATGG AAGAAACCACTACTGAGAAAACATCCCTAGAGGATATTGATTTAGGCTCA GGATTATTTGAAAAGCCCAAAGCCACAGAACTCATAGAATTTTCAACAAT CAAAGTCACAGTTCCAAGTGATATTACCACTGCCTTCAGTTCAGTAGACA GACTTCACACAACTTCAGCATTCAAGCCATCTTCCGCGATCACTAAGAAA CCACCTCTCATCGACAGGGAACCTGGTGAAGAACAACCAGTGACATGGT AATCATTGGAGAATCAACATCTCATGTTCCTCCCACTACCCTTGAAGATAT TGTAGCCAAGGAAACAGAAACCGATATTGATAGAGAGTATTTCACGACTT CAAGTCCTCCTGCTACACAGCCAACAAGACCACCCACTGTGGAAGACAAA AAAATTTCACCCTGACATTAATGTTTATATTATTGAGGTCAGAGAAAATAA GACAGGTCGAATGAGTGATTTGAGTGTAATTGGTCATCCAATAGATTCAG AATCTAAAGAAGATGAACCTTGTAGTGAAGAAACAGATCCAGTGCATGAT CTAATGGCTGAAATTTTACCTGAATTCCCTGACATAATTGAAATAGACCTA TACCACAGTGAAGAAATGAAGAAGAAGAAGAAGAGTGTGCAAATGCTA CTGATGTGACAACCACCCATCTGTGCAGTACATAAATGGGAAGCATCTC GTTACCACTGTGCCCAAGGACCCAGAAGCTGCAGAAGCTAGGCGTGGCCA GTTTGAAAGTGTTGCACCTTCTCAGAATTTCTCGGACAGCTCTGAAAGTGA TACTCATCCATTTGTAATAGCCAAAACGGAATTGTCTACTGCTGTGCAACC TAATGAATCTACAGAAACAACTGAGTCTCTTGAAGTTACATGGAAGCCTG AGACTTACCCTGAAACATCAGAACATTTTTCAGGTGGTGAGCCTGATGTTT TCCCACAGTCCCATTCCATGAGGAATTTGAAAGTGGAACAGCCAAAAAA GGGGCAGAATCAGTCACAGAGAGAGATACTGAAGTTGGTCATCAGGCAC ATGAACATACTGAACCTGTATCTCTGTTTCCTGAAGAGTCTTCAGGAGAGA TTGCCATTGACCAAGAATCTCAGAAAATAGCCTTTGCAAGGGCTACAGAA GTAACATTTGGTGAAGAGGTAGAAAAAAGTACTTCTGTCACATACACTCC CACTATAGTTCCAAGTTCTGCATCAGCATATGTTTCAGAGGAAGAAGCAG TTACCCTAATAGGAAATCCTTGGCCAGATGACCTGTTGTCTACCAAAGAA AGCTGGGTAGAAGCAACTCCTAGACAAGTTGTAGAGCTCTCAGGGAGTTC TTCGATTCCAATTACAGAAGGCTCTGGAGAAGCAGAAGAAGATGAAGATA CAATGTTCACCATGGTAACTGATTTATCACAGAGAAATACTACTGATACA CTCATTACTTTAGACACTAGCAGGATAATCACAGAAAGCTTTTTTGAGGTT CCTGCAACCACCATTTATCCAGTTTCTGAACAACCTTCTGCAAAAGTGGTG CCTACCAAGTTTGTAAGTGAAACAGACACTTCTGAGTGGATTTCCAGTACC

ACTGTTGAGGAAAAGAAAAGGAAGGAGGAGGAACTACAGGTACGG TACCCTTTGAATTAGAAAGTCCAAATGTAGCTACATCTAGTGATTCAGGTA CCAGGAAAAGTTTTATGTCCTTGACAACACCAACACACGTCTGAAAGGGAA ATGACAGATTCTACTCCTGTCTTTACAGAAACAAATACATTAGAAAATTTG GGGGCACAGACCACTGAGCACAGCAGTATCCATCAACCTGGGGTTCAGGA AGGGCTGACCACTCTCCCACGTAGTCCTGCCTCTGTCTTTATGGAGCAGGG CTCTGGAGAAGCTGCTGCCGACCCAGAAACCACCACTGTTTCTTCATTTTC ATTAAACGTAGAGTATGCAATTCAAGCCGAAAAGGAAGTAGCTGGCACTT TGTCTCCGCATGTGGAAACTACATTCTCCACTGAGCCAACAGGACTGGTTT TGAGTACAGTAATGGACAGAGTAGTTGCTGAAAATATAACCCAAACATCC AGGGAAATAGTGATTTCAGAGCGATTAGGAGAACCAAATTATGGGGCAG AAATAAGGGGCTTTTCCACAGGTTTTCCTTTGGAGGAAGATTTCAGTGGTG ACTTTAGAGAATACTCAACAGTGTCTCATCCCATAGCAAAAGAAGAAACG GTAATGATGGAAGGCTCTGGAGATGCAGCATTTAGGGACACCCAGACTTC ACCATCTACAGTACCTACTTCAGTTCACATCAGTCACATATCTGACTCAGA AGGACCCAGTAGCACCATGGTCAGCACTTCAGCCTTCCCCTGGGAAGAGT TTACATCCTCAGCTGAGGGCTCAGGTGAGCAACTGGTCACAGTCAGCAGC TCTGTTGTTCCAGTGCTTCCCAGTGCTGCAAAAGTTTTCTGGTACAGCT TCCTCCATTATCGACGAAGGATTGGGAGAAGTGGGTACTGTCAATGAAAT TGATAGAAGATCCACCATTTTACCAACAGCAGAAGTGGAAGGTACGAAAG CTCCAGTAGAGAAGGAGGAAGTAAAGGTCAGTGGCACAGTTTCAACAAA CTTTCCCCAAACTATAGAGCCAGCCAAATTATGGTCTAGGCAAGAAGTCA ACCCTGTAAGACAAGAAATTGAAAGTGAAACAACATCAGAGGAACAAAT TCAAGAAGAAAGTCATTTGAATCCCCTCAAAACTCTCCTGCAACAGAAC AAACAATCTTTGATTCACAGACATTTACTGAAACTGAACTCAAAACCACA GATTATTCTGTACTAACAACAAGAAAACTTACAGTGATGATAAAGAAAT GAAGGAGGAAGACACTTCTTTAGTTAACATGTCTACTCCAGATCCAGATG CAAATGGCTTGGAATCTTACACAACTCTCCCTGAAGCTACTGAAAAGTCA CATTTTTCTTAGCTACTGCATTAGTAACTGAATCTATACCAGCTGAACAT GTAGTCACAGATTCACCAATCAAAAAGGAAGAAAGTACAAAACATTTTCC GAAAGGCATGAGACCAACAATTCAAGAGTCAGATACTGAGCTCTTATTCT CTGGACTGGGATCAGGAGAAGAAGTTTTACCTACTCTACCAACAGAGTCA GTGAATTTTACTGAAGTGGAACAAATCAATAACACATTATATCCCCACAC TTCTCAAGTGGAAAGTACCTCAAGTGACAAAATTGAAGACTTTAACAGAA TGGAAAATGTGGCAAAAGAAGTTGGACCACTCGTATCTCAAACAGACATC TTTGAAGGTAGTGGGTCAGTAACCAGCACAACATTAATAGAAATTTTAAG TGACACTGGAGCAGAAGGACCCACGGTGGCACCTCTCCCTTTCTCCACGG ACATCGGACATCCTCAAAATCAGACTGTCAGGTGGGCAGAAGAAATCCAG ACTAGTAGACCACAAACCATAACTGAACAAGACTCTAACAAGAATTCTTC AACAGCAGAAATTAACGAAACAACAACCTCATCTACTGATTTTCTGGCTA GAGCTTATGGTTTTGAAATGGCCAAAGAATTTGTTACATCAGCACCAAAA CCATCTGACTTGTATTATGAACCTTCTGGAGAAGGATCTGGAGAAGTGGA TATTGTTGATTCATTTCACACTTCTGCAACTACTCAGGCAACCAGACAAGA AAGCAGCACCACATITGTTTCTGATGGGTCCCTGGAAAAACATCCTGAGG TGCCAAGCGCTAAAGCTGTTACTGCTGATGGATTCCCAACAGTTTCAGTGA TGCTGCCTCTTCATTCAGAGCAGAACAAAAGCTCCCCTGATCCAACTAGC ACACTGTCAAATACAGTGTCATATGAGAGGTCCACAGACGGTAGTTTCCA AGACCGTTTCAGGGAATTCGAGGATTCCACCTTAAAACCTAACAGAAAAA AACCCACTGAAAATATTATCATAGACCTGGACAAAGAGGACAAGGATTTA

ATATTGACAATTACAGAGAGTACCATCCTTGAAATTCTACCTGAGCTGAC ATCGGATAAAAATACTATCATAGATATTGATCATACTAAACCTGTGTATG AAGACATTCTTGGAATGCAAACAGATATAGATACAGAGGTACCATCAGAA CCACATGACAGTAATGAAGAAGTAATGATGACAGCACTCAAGTTCAAGA GATCTATGAGGCAGCTGTCAACCTTTCTTTAACTGAGGAAACATTTGAGG GCTCTGCTGATGTTCTGGCTAGCTACACTCAGGCAACACATGATGAATCA ATGACTTATGAAGATAGAAGCCAACTAGATCACATGGGCTTTCACTTCAC AACTGGGATCCCTGCTCCTAGCACAGAAACAGAATTAGACGTTTTACTTCC CACGGCAACATCCCTGCCAATTCCTCGTAAGTCTGCCACAGTTATTCCAGA GATTGAAGGAATAAAAGCTGAAGCAAAAGCCCTGGATGACATGTTTGAAT CAAGCACTTTGTCTGATGGTCAAGCTATTGCAGACCAAAGTGAAATAATA CCAACATTGGGCCAATTTGAAAGGACTCAGGAGGAGTATGAAGACAAAA AACATGCTGGTCCTTCTTTTCAGCCAGAATTCTCTTCAGGAGCTGAGGAGG CATTAGTAGACCATACTCCCTATCTAAGTATTGCTACTACCCACCTTATGG ATCAGAGTGTAACAGAGGTGCCTGATGTGATGGAAGGATCCAATCCCCCA TATTACACTGATACAACATTAGCAGTTTCAACATTTGCGAAGTTGTCTTCT CAGACACCATCATCTCCCCTCACTATCTCAGGCAGTGAAGCCTCTGGA CACACAGAGATCCCCAGCCCAGTGCTCTGCCAGGAATAGACGTCGGCTC ATCTGTAATGTCCCCACAGGATTCTTTTAAGGAAATTCATGTAAATATTGA AGCAACTTTCAAACCATCAAGTGAGGAATACCTTCACATAACTGAGCCTC CCTCTTTATCTCCTGACACAAAATTAGAACCTTCAGAAGATGATGGTAAAC CTGAGTTATTAGAAGAAATGGAAGCTTCTCCCACAGAACTTATTGCTGTG GAAGGAACTGAGATTCTCCAAGATTTCCAAAACAAACCGATGGTCAAGT TTCTGGAGAGCAATCAAGATGTTTCCCACCATTAAAACACCTGAGGCTG GAACTGTTATTACAACTGCCGATGAAATTGAATTAGAAGGTGCTACACAG TGGCCACACTCTACTTCTGCTTCTGCCACCTATGGGGTCGAGGCAGGTGTG GTGCCTTGGCTAAGTCCACAGACTTCTGAGAGGCCCACGCTTTCTTCTTCT CCAGAAATAAACCCTGAAACTCAAGCAGCTTTAATCAGAGGGCAGGATTC CACGATAGCAGCATCAGAACAGCAAGTGGCAGCAGAATTCTTGATTCCA ATGATCAGGCAACAGTAAACCCTGTGGAATTTAATACTGAGGTTGCAACA CCACCATTTTCCCTTCTGGAGACTTCTAATGAAACAGATTTCCTGATTGGC ATTAATGAAGAGTCAGTGGAAGGCACGGCAATCTATTTACCAGGACCTGA TCGCTGCAAAATGAACCCGTGCCTTAACGGAGGCACCTGTTATCCTACTG AAACTTCCTACGTATGCACCTGTGTGCCAGGATACAGCGGAGACCAGTGT GAACTTGATTTTGATGAATGTCACTCTAATCCCTGTCGTAATGGAGCCACT TGTGTTGATGGTTTTAACACATTCAGGTGCCTCTGCCTTCCAAGTTATGTT ATTCCAAGGCAGTGCTACAAATACTTTGCCCATCGACGCACATGGGATG CAGCTGAACGGGAATGCCGTCTGCAGGGTGCCCATCTCACAAGCATCCTG TCTCACGAAGAACAAATGTTTGTTAATCGTGTGGGCCATGATTATCAGTGG ATAGGCCTCAATGACAAGATGTTTGAGCATGACTTCCGTTGGACTGATGG TTTCTGCTGGAGAAGACTGTGTTGTAATCATTTGGCATGAGAATGGCCAGT GGAATGATGTTCCCTGCAATTACCATCTCACCTATACGTGCAAGAAAGGA ACAGTTGCTTGCGGCCAGCCCCCTGTTGTAGAAAATGCCAAGACCTTTGG AAAGATGAAACCTCGTTATGAAATCAACTCCCTGATTAGATACCACTGCA AAGATGGTTTCATCAACGTCACCTTCCAACTATCCGGTGCTTAGGAAATG GAAGATGGCTATACCTAAAATTACCTGCATGAACCCATCTGCATACCAA AGGACTTATTCTATGAAATACTTTAAAAATTCCTCATCAGCAAAGGACAA TTCAATAAATACATCCAAACATGATCATCGTTGGAGCCGGAGGTGGCAGG

**AGTCGAGGCGCTGATCCCTAAAATGGCGAACATGTGTTTTCATCATTTCAG** CCAAAGTCCTAACTTCCTGTGCCTTTCCTATCACCTCGAGAAGTAATTATC AGTTGGTTTGGATTTTTGGACCACCGTTCAGTCATTTTGGGTTGCCGTGCT CCCAAAACATTTTAAATGAAAGTATTGGCATTCAAAAAGACAGCAGACAA AATGAAAGAAAATGAGAGCAGAAAGTAAGCATTTCCAGCCTATCTAATTT CTTTAGTTTTCTATTTGCCTCCAGTGCAGTCCATTTCCTAATGTATACCAGC CTACTGTACTATTTAAAATGCTCAATTTCAGCACCGATGGCCATGTAAATA AGATGATTTAATGTTGATTTTAATCCTGTATATAAAAATAAAAAGTCACAAT GAGTTTGGGCATATTTAATGATGATTATGGAGCCTTAGAGGTCTTTAATCA TTGGTTCGCTGCTTTTATGTAGTTTAGGCTGGAAATGGTTTCACTTGCTCT TTGACTGTCAGCAAGACTGAAGATGGCTTTTCCTGGACAGCTAGAAAACA CAAAATCTTGTAGGTCATTGCACCTATCTCAGCCATAGGTGCAGTTTGCTT CTACATGATGCTAAAGGCTGCGAATGGGATCCTGATGGAACTAAGGACTC CAATGTCGAACTCTTCTTTGCTGCATTCCTTTTTCTTCACTTACAAGAAAGG **SEQ ID NO: 85** CCTGAATGGAGGACTTTTCTGTAACCAGG

# N-Acylsphingosine Amidohydrosase 1

>gi|30089929|ref|NM\_004315.2| Homo sapiens N-acylsphingosine amidohydrolase (acid ceramidase) 1 (ASAH1), transcript variant 2, mRNA | qPCR forward\_primer match [1212..1228] | qPCR reverse\_primer match [1290..1266] | qPCR probe match [1233..1260]

GGACTTTGAAATCCAACCGGTCACCTACCGCGCGACTGTGTCCA CGGATGGCACGAAAGCCAAGCGAGTCCCCCTGCCGAGCTACTCGCGTCCG CCTCCTCCCAAGCTGAGCTCTGCTCCGCCCACCTGAGTCCTTCGCCAGTTA GGAGGAAACACAGCCGCTTAATGAACTGCTGCATCGGGCTGGGAGAGAA AGCTCGCGGGTCCCACCGGGCCTCCTACCCAAGTCTCAGCGCGCTTTTCAC CGAGGCCTCAATTCTGGGATTTGGCAGCTTTGCTGTGAAAGCCCAATGGA CAGAGGACTGCAGAAAATCAACCTATCCTCCTTCAGGACCAACGTACAGA GGTGCAGTTCCATGGTACACCATAAATCTTGACTTACCACCCTACAAAAG ATGGCATGAATTGATGCTTGACAAGGCACCAATGCTAAAGGTTATAGTGA ATTCTCTGAAGAATATGATAAATACATTCGTGCCAAGTGGAAAAGTTATG CAGGTGGTGGATGAAAAATTGCCTGGCCTACTTGGCAACTTTCCTGGCCCT TTTGAAGAGGAAATGAAGGGTATTGCCGCTGTTACTGATATACCTTTAGG AGAGATTATTTCATTCAATATTTTTTATGAATTATTTACCATTTGTACTTCA ATAGTAGCAGAAGACAAAAAAGGTCATCTAATACATGGGAGAAACATGG ATTTTGGAGTATTTCTTGGGTGGAACATAAATAATGATACCTGGGTCATAA CTGAGCAACTAAAACCTTTAACAGTGAATTTGGATTTCCAAAGAAACAAC AAAACTGTCTTCAAGGCTTCAAGCTTTGCTGGCTATGTGGGCATGTTAACA GGATTCAAACCAGGACTGTTCAGTCTTACACTGAATGAACGTTTCAGTATA CATGTGGATAGGGTTCCTCACTAGAACAGTTCTGGAAAATAGCACAAGTT ATGAAGAAGCCAAGAATTTATTGACCAAGACCAAGATATTGGCCCCAGCC TACTTTATCCTGGGAGGCAACCAGTCTGGGGAAGGTTGTGTGATTACACG AGACAGAAAGGAATCATTGGATGTATATGAACTCGATGCTAAGCAGGGTA GATGGTATGTGGTACAAACAAATTATGACCGTTGGAAACATCCCTTCTTCC TTGATGATCGCAGAACGCCTGCAAAGATGTGTCTGAACCGCACCAGCCAA GAGAATATCTCATTTGAAACCATGTATGATGTCCTGTCAACAAAACCTGTC

CTCAACAAGCTGACCGTATACACAACCTTGATAGATGTTACCAAAGGTCA ATTCGAAACTTACCTGCGGGACTGCCCTGACCCTTGTATAGGTTGGTGAGC ACACGTCTGGCCTACAGAATGCGGCCTCTGAGACATGAAGACACCATCTC CATGTGACCGAACACTGCAGCTGTCTGACCTTCCAAAGACTAAGACTCGC GGCAGGTTCTCTTTGAGTCAAAAGCTTGTCTTCGTCCATCTGTTGACAAAT GACAGACCTTTTTTTTCCCCCATCAGTTGATTTTTCTTATTTACAGATAAC TTCTTTAGGGGAAGTAAAACAGTCATCTAGAATTCACTGAGTTTTGTTTCA CTTTGACATTTGGGGATCTGGTGGGCAGTCGAACCATGGTGAACTCCACCT CCGTGGAATAAATGGAGATTCAGCGTGGGTGTTGAATCCAGCACGTCTGT GTGAGTAACGGGACAGTAAACACTCCACATTCTTCAGTTTTTCACTTCTAC CTACATATTTGTATGTTTTCTGTATAACAGCCTTTTCCTTCTGGTTCTAAC TGCTGTTAAAATTAATATCATTATCTTTGCTGTTATTGACAGCGATATA ATTTTATTACATATGATTAGAGGGATGAGACAGACATTCACCTGTATATTT CTTTTAATGGGCACAAAATGGGCCCTTGCCTCTAAATAGCACTTTTTGGGG TTCAAGAAGTAATCAGTATGCAAAGCAATCTTTTATACAATAATTGAAGT GTTCCCTTTTTCATAATTACTGTACTTCCCAGTAACCCTAAGGAAGTTGCT AAAGACTTGTGGAAAATAGGAAGTGAACCCATATTTTAAATTCTCATAAG TAGCATTCATGTAATAAACAGGTTTTTAGTTTGTTCTTCAGATTGATAGGG AGTTTTAAAGAAATTTTAGTAGTTACTAAAATTATGTTACTGTATTTTTCA GAAATCAAACTGCTTATGAAAAGTACTAATAGAACTTGTTAACCTTTCTAA CCTTCACGATTAACTGTGAAATGTACGTCATTTGTGCAAGACCGTTTGTCC ACTTCATTTTGTATAATCACAGTTGTGTTCCTGACACTCAATAAACAGTCA TTGGAAAGAGTGCCAGTCAGCAGTCATGCA **SEO ID NO: 86** 

## N-Acylsphingosine Amidohydrolase 1 Transcript Variant 1

>gi|30089927|ref|NM\_177924.1| Homo sapiens N-acylsphingosine amidohydrolase (acid ceramidase) 1 (ASAH1), transcript variant 1, mRNA | qPCR forward\_primer match [1050..1066] | qPCR reverse\_primer match [1128..1104] | qPCR probe match [1071..1098]

GGCTCTTCTTTGCCTCTGCTGGAGTCCGGGGAGTGGCGTTGGCTGCT AGAGCGATGCCGGGCCGAGTTGCGTCGCCTTAGTCCTCCTGGCTGCCGC CGTCAGCTGTGCCGTCGCGCAGCACGCGCCGCGTGGACAGAGGACTGCA GAAAATCAACCTATCCTCCTTCAGGACCAACGTACAGAGGTGCAGTTCCA TGGTACACCATAAATCTTGACTTACCACCCTACAAAAGATGGCATGAATT GATGCTTGACAAGGCACCAATGCTAAAGGTTATAGTGAATTCTCTGAAGA ATATGATAAATACATTCGTGCCAAGTGGAAAAGTTATGCAGGTGGTGGAT GAAAAATTGCCTGGCCTACTTGGCAACTTTCCTGGCCCTTTTGAAGAGGAA ATGAAGGGTATTGCCGCTGTTACTGATATACCTTTAGGAGAGATTATTTCA TTCAATATTTTTATGAATTATTTACCATTTGTACTTCAATAGTAGCAGAA GACAAAAAGGTCATCTAATACATGGGAGAAACATGGATTTTGGAGTATT TCTTGGGTGGAACATAAATAATGATACCTGGGTCATAACTGAGCAACTAA AACCTTTAACAGTGAATTTGGATTTCCAAAGAAACAACAAAACTGTCTTC AAGGCTTCAAGCTTTGCTGGCTATGTGGGCATGTTAACAGGATTCAAACC AGGACTGTTCAGTCTTACACTGAATGAACGTTTCAGTATAAATGGTGGTTA TCTGGGTATTCTAGAATGGATTCTGGGAAAGAAGATGCCATGTGGATAG GGTTCCTCACTAGAACAGTTCTGGAAAATAGCACAAGTTATGAAGAAGCC

AAGAATTTATTGACCAAGACCAAGATATTGGCCCCAGCCTACTTTATCCTG GGAGGCAACCAGTCTGGGGAAGGTTGTGTGATTACACGAGACAGAAAGG **AATCATTGGATGTATATGAACTCGATGCTAAGCAGGGTAGATGGTATGTG** GTACAAACAAATTATGACCGTTGGAAACATCCCTTCTTCCTTGATGATCGC AGAACGCCTGCAAAGATGTGTCTGAACCGCACCAGCCAAGAGAATATCTC ATTTGAAACCATGTATGATGTCCTGTCAACAAACCTGTCCTCAACAAGCT GACCGTATACACAACCTTGATAGATGTTACCAAAGGTCAATTCGAAACTT ACCTGCGGGACTGCCCTGACCCTTGTATAGGTTGGTGAGCACACGTCTGG CCTACAGAATGCGGCCTCTGAGACATGAAGACACCATCTCCATGTGACCG AACACTGCAGCTGTCTGACCTTCCAAAGACTAAGACTCGCGGCAGGTTCT CTTTGAGTCAAAAGCTTGTCTTCGTCCATCTGTTGACAAATGACAGACCTT TTTTTTCCCCCATCAGTTGATTTTTCTTATTTACAGATAACTTCTTTAGGG GAAGTAAAACAGTCATCTAGAATTCACTGAGTTTTGTTTCACTTTGACATT TGGGGATCTGGTGGCAGTCGAACCATGGTGAACTCCACCTCCGTGGAAT AAATGGAGATTCAGCGTGGGTGTTGAATCCAGCACGTCTGTGTGAGTAAC TGTATGTTTTCTGTATAACAGCCTTTTCCTTCTGGTTCTAACTGCTGTTAA **AATTAATATCATTATCTTTGCTGTTATTGACAGCGATATAATTTTATTAC** ATATGATTAGAGGGATGAGACAGACATTCACCTGTATATTTCTTTTAATGG GCACAAAATGGGCCCTTGCCTCTAAATAGCACTTTTTGGGGTTCAAGAAG TAATCAGTATGCAAAGCAATCTTTTATACAATAATTGAAGTGTTCCCTTTT TCATAATTACTGTACTTCCCAGTAACCCTAAGGAAGTTGCTAACTTAAAAA ACTGCATCCCACGTTCTGTTAATTTAGTAAATAAACAAGTCAAAGACTTGT GGAAAATAGGAAGTGAACCCATATTTTAAATTCTCATAAGTAGCATTCAT GTAATAAACAGGTTTTTAGTTTGTTCTTCAGATTGATAGGGAGTTTTAAAG **AAATTTTAGTAGTTACTAAAATTATGTTACTGTATTTTTCAGAAATCAAAC** TGCTTATGAAAAGTACTAATAGAACTTGTTAACCTTTCTAACCTTCACGAT TAACTGTGAAATGTACGTCATTTGTGCAAGACCGTTTGTCCACTTCATTTT GTATAATCACAGTTGTGTTCCTGACACTCAATAAACAGTCATTGGAAAGA **SEO ID NO: 87 GTGCCAGTCAGCAGTCATGCA** 

#### Protease, Serine 11

>gi|21327712|ref|NM\_002775.2| Homo sapiens protease, serine, 11 (IGF binding) (PRSS11), mRNA | qPCR forward\_primer match [1030..1048] | qPCR reverse primer match [1106..1083] | qPCR probe match [1080..1050]

GGTGGCTAGTGGGTCTGGGTTTATTGTGTCGGAAGATGGACTGATCGTGA CAAATGCCCACGTGGTGACCAACAAGCACCGGGTCAAAGTTGAGCTGAAG AACGGTGCCACTTACGAAGCCAAAATCAAGGATGTGGATGAGAAAGCAG ACATCGCACTCAAAAATTGACCACCAGGGCAAGCTGCCTGTCCTGCTG CTTGGCCGCTCCTCAGAGCTGCGGCCGGGAGAGTTCGTGGTCGCCATCGG AAGCCCGTTTTCCCTTCAAAACACAGTCACCACCGGGATCGTGAGCACCA CCCAGCGAGGCGCAAAGAGCTGGGGCTCCGCAACTCAGACATGGACTA CATCCAGACCGACGCCATCATCAACTATGGAAACTCGGGAGGCCCGTTAG TAAACCTGGACGGTGAAGTGATTGGAATTAACACTTTGAAAGTGACAGCT GGAATCTCCTTTGCAATCCCATCTGATAAGATTAAAAAGTTCCTCACGGAG TCCCATGACCGACAGGCCAAAGGAAAAGCCATCACCAAGAAGAAGTATA TTGGTATCCGAATGATGTCACTCACGTCCAGCAAAGCCAAAGAGCTGAAG GACCGGCACCGGGACTTCCCAGACGTGATCTCAGGAGCGTATATAATTGA AGTAATTCCTGATACCCCAGCAGAAGCTGGTGGTCTCAAGGAAAACGACG TCATAATCAGCATCAATGGACAGTCCGTGGTCTCCGCCAATGATGTCAGC GACGTCATTAAAAGGGAAAGCACCCTGAACATGGTGGTCCGCAGGGGTA ATGAAGATATCATGATCACAGTGATTCCCGAAGAAATTGACCCATAGGCA GAGGCATGAGCTTCATGTTTCCCTCAAAGACTCTCCCGTGGATGAC GGATGAGGACTCTGGGCTGCTGGAATAGGACACTCAAGACTTTTGACTGC CATTTTGTTTCAGTGGAGACTCCCTGGCCAACAGAATCCTTCTTGATA GCCCTTCTGTATCCTATGTATGCAGTGTGCTTTTTCTTGCCAGCTTGGGCCA TTCTTGCTTAGACAGTCAGCATTTGTCTCCTCCTTTAACTGAGTCATCATCT TAGTCCAACTAATGCAGTCGATACAATGCGTAGATAGAAGAAGCCCCACG GGAGCCAGGATGGGACTGGTCGTGTTTTGTGCTTTTCTCCAAGTCAGCACCC AAAGGTCAATGCACAGAGACCCCGGGTGGGTGAGCGCTGGCTTCTCAAAC GGCCGAAGTTGCCTCTTTTAGGAATCTCTTTGGAATTGGGAGCACGATGAC TCTGAGTTTGAGCTATTAAAGTACTTCTTACACATTG **SEO ID NO: 88** 

#### Secreted Frizzled-Related Protein 2

>gi|42656988|ref|XM\_050625.4| Homo sapiens secreted frizzled-related protein 2 (SFRP2), mRNA | qPCR forward\_primer match [686..703] | qPCR reverse\_primer match [750..728] | qPCR probe match [705..726]

AAAATGATGATGACAACGACATAATGGAAACGCTTTGTAAAAATGATTT
TGCACTGAAAATAAAAGTGAAGGAGATAACCTACATCAACCGAGATACC
AAAATCATCCTGGAGACCAAGAGCAAGACCATTTACAAGCTGAACGGTGT
GTCCGAAAGGGACCTGAAGAAATCGGTGCTGTGGCTCAAAGACAGCTTGC
AGTGCACCTGTGAGGAGATGAACGACATCAACGCGCCCTATCTGGTCATG
GGACAGAAACAGGGTGGGGAGCTGGTGATCACCTCGGTGAAGCGGTGGC
AGAAGGGGCAGAGAGAGTTCAAGCGCATCTCCCGCAGCATCCGCAAGCT
GCAGTGCTAGTCCCGGCATCCTGATGGCTCCGACAGGCCTGCTCCAGAGC
ACGGCTGACCATTTCTGCTCCGGGATCTCAGCTCCCGTTCCCCAAGCACAC
TCCTAGCTGCTCCAGTCTCAGCCTGGGCAGCTTCCCCCTGCCTTTTGCACG
TTTGCATCCCCAGCATTTCCTGAGTTATAAGGCCACAGGAGTGGATAGCTG
TTTTCACCTAAAGGAAAAGCCCACCCGAATCTTGTAGAAATATTCAAACT
AATAAAATCATGAATATTTTTATGAAGTTTAAAAA
SEQ ID NO: 89

### Phospholipase A2, Group XIIB

>gi|45505134|ref|NM\_032562.2| Homo sapiens phospholipase A2, group XIIB (PLA2G12B), mRNA

TGTCCCTGGAATTCTGGGACACTGGCTGGGGTTTGAGGAGAGAAGC CAGTACCTACCTGGCTGCAGGATGAAGCTGGCCAGTGGCTTCTTGGTTTTG TGGCTCAGCCTTGGGGTGGCCTGGCTCAGAGCGACACGAGCCCTGACAC GGAGGAGTCCTATTCAGACTGGGGCCCTTCGGCACCTCCGGGGAAGCTTTG ATGGAGTCTGTCAGTACAGGTGCCGATATGGAAAGGCACCAATGCCCAGA CCTGGCTACAAGCCCCAAGAGCCCAATGGCTGCGGCTCCTATTTCCTGGGT CTCAAGGTACCAGAAAGTATGGACTTGGGCATTCCAGCAATGACAAAGTG CTGCAACCAGCTGGATGTCTGTTATGACACTTGCGGTGCCAACAAATATC GCTGTGATGCAAAATTCCGATGGTGTCTCCACTCGATCTGCTCTGACCTTA AGCGGAGTCTGGGCTTTGTCTCCAAAGTGGAAGCAGCCTGTGATTCCCTG GTTGACACTGTGTTCAACACCGTGTGGACCTTGGGCTGCCGCCCCTTTATG TATGAGGAAGAAGTGATTCCTTCCTGGTTTTGAGTGACACCACAGCTGTCA CAGTTTGGACACCACAAAGCAGGAGAAAAGGGAACATTTTTCTACAGCTGG AAAGTGAGTCCTATCCTTTGAGGAAATTTGAAAAAAGACATGGAGTGGTT ATTCCTGGACCTGATAGTTATATTCATGAGTGAAATTGTGGGGAGTCCAGC CATTTGGGAGGCAATGACTTTCTGCTGGCCCATGTTTCAGTTGCCAGTAAG CTTCTCACATTTAATAAAGTGTACTTTTTAGAACATT SEQ ID NO: 90

# Spondin 2, Extracellular Matrix Protein

>gi|6912681|ref|NM\_012445.1| Homo sapiens spondin 2, extracellular matrix protein (SPON2), mRNA

GCACGAGGAAGAGGTGATCCGACCCGGGAAGGTCGCTGGGCA GGGCGAGTTGGGAAAGCGGCAGCCCCCGCCGCCGCAGCCCTTCTCC TCCTTTCTCCCACGTCCTATCTGCCTCTCGCTGGAGGCCAGGCCGTGCAGC

ATCGAAGACAGGAGGAACTGGAGCCTCATTGGCCGGCCCGGGGCCCCGG CCTCGGGCTTAAATAGGAGCTCCGGGCTCTGGCTGGGACCCGACCGCTGC CGGCCGCGCTCCCGCTGCCCGGGTGATGGAAAACCCCAGCCCGGC CGCCGCCTGGGCAAGGCCCTCTGCGCTCTCCTGGCCACTCTCGGCGC CGCCGCCAGCCTCTTGGGGGAGAGTCCATCTGTTCCGCCAGAGCCCCGG CCAAATACAGCATCACCTTCACGGGCAAGTGGAGCCAGACGGCCTTCCCC **AAGCAGTACCCCTGTTCCGCCCCCTGCGCAGTGGTCTTCGCTGCTGGGG** GCCGCGCATAGCTCCGACTACAGCATGTGGAGGAAGAACCAGTACGTCAG TAACGGGCTGCGCGACTTTGCGGAGCGCGGCGAGGCCTGGGCGCTGATGA AGGAGATCGAGGCGGGGGGGGGGGCGCTGCAGAGCGTGCACGCGGTGTT TTCGGCGCCGCCGTCCCCAGCGGCACCGGGCAGACGTCGGCGGAGCTGG AGGTGCAGCGCAGGCACTCGCTGGTCTCGTTTGTGGTGCGCATCGTGCCC AGCCCGACTGGTTCGTGGGCGTGGACAGCCTGGACCTGTGCGACGGGGA CCGTTGGCGGGACAGGCGCGCTGGACCTGTACCCCTACGACGCCGGGA CGGACAGCGGCTTCACCTTCTCCCCCAACTTCGCCACCATCCCGCAGG ACACGGTGACCGAGATAACGTCCTCCTCTCCCAGCCACCCGGCCAACTCC TTCTACTACCCGCGGCTGAAGGCCCTGCCTCCCATCGCCAGGGTGACACTG GTGCGCTGCGACAGAGCCCCAGGGCCTTCATCCCTCCCGCCCCAGTCCT GCCCAGCAGGACAATGAGATTGTAGACAGCGCCTCAGTTCCAGAAACGC CGCTGGACTGCGAGGTCTCCCTGTGGTCGTCCTGGGGACTGTGCGGAGGC CACTGTGGGAGGCTCGGGACCAAGAGCAGGACTCGCTACGTCCGGGTCCA GCCGCCAACAACGGGAGCCCCTGCCCCGAGCTCGAAGAAGAGGCTGAG TGCGTCCTGATAACTGCGTCTAAGACCAGAGCCCCGCAGCCCCTGGGGC CCCGGAGCCATGGGGTGTCGGGGGCTCCTGTGCAGGCTCATGCTGCAGG CGGCCGAGGCACAGGGGTTTCGCGCTGCTCCTGACCGCGGTGAGGCCGC GCCGACCATCTCTGCACTGAAGGGCCCTCTGGTGGCCGGCACGGCATTG GGAAACAGCCTCCTCTTTCCCAACCTTGCTTCTTAGGGGCCCCCGTGTCC CGTCTGCTCTCGCCTCCTCCTGCAGGATAAAGTCATCCCCAAGGCTC CAGCTACTCTAAATTATGGTCTCCTTATAAGTTATTGCTGCTCCAGGAGAT TGTCCTTCATCGTCCAGGGGCCTGGCTCCCACGTGGTTGCAGATACCTCAG ACCTGGTGCTCTAGGCTGTGCTGAGCCCACTCTCCCGAGGGCGCATCCAA GCGGGGCCACTTGAGAAGTGAATAAATGGGGCGGTTTCGGAAGCGTCA GTGTTTCCATGTTATGGATCTCTCTGCGTTTGAATAAAGACTATCTCTGTTG **CTCAC SEQ ID NO: 91** 

## Olfactomedin 1, Transcript Variant 3

>gi|34335282|ref|NM\_058199.2| Homo sapiens olfactomedin 1 (OLFM1), transcript variant 3, mRNA

# **Thrombospondin Repeat Containing 1**

>gi|38016903|ref|NM\_019032.2| Homo sapiens thrombospondin repeat containing 1 (TSRC1), mRNA

GGGGCCCCAGTGGCCGCCGCGGAGCGAGGTTGCCTGGAGAGAGCG CCTGGGCGCAGAGGGTTAACGGCCACCGGGGGCTCGCAGAGCAGGAG GGTGCTCTCGGACGGTGTCCCCCACTGCACTCCTGAACTTGGAGGACA GGGTCGCCGCGAGGGACGCAGAGAGCACCCTCCACGCCCAGATGCCTGCG TAGTTTTTGTGACCAGTCCGCTCCTGCCTCCCCTGGGGCAGTAGAGGGGG AGCGATGGAGACTGGACTGGCAGGCCCTGGCTGTATCTGCTGCTTC TGTCCCTCCCTCAGCTCTGCTTGGATCAGGAGGTGTTGTCCGGACACTCTC TTCAGACACCTACAGAGGAGGGCCAGGGCCCCGAAGGTGTCTGGGGACCT TGGGTCCAGTGGGCCTCTTGCTCCCAGCCCTGCGGGGTGGGGGTGCAGCG CAGGAGCCGGACATGTCAGCTCCCTACAGTGCAGCTCCACCCGAGTCTGC CCTCCCTCCCGGCCCCAAGACATCCAGAAGCCCTCCTCCCCGGGGCC AGGGTCCCAGACCCCAGACTTCTCCAGAAACCCTCCCCTTGTACAGGACA CAGTCTCGGGGAAGGGGTGCCCACTTCGAGGTCCCGCTTCCCACCTAGG GAGAGAGGGCCAGGAGATTCGAGCGGCCAGGAGGTCCCGGCTTCGA GACCCATCAAGCCAGGAATGTTCGGTTATGGGAGAGTGCCCTTTGCATT TGTCCCTGATCTCTAGAGGGGAAGAGGCTATTCCGTCCCCTACTCCAA ACAGAACTGTCTGTCCACACCCCATCCCCCAAGCAGAACCTCTAAGCCC TGAAACTGCTCAGACAGAGGTGGCCCCCAGAACCAGGCCTGCCCCCTAC GGCATCACCCAGAGCCCAGGCCTCTGGCACAGAGCCCCCCTCACCCACG CACTCCTTAGGAGAAGGTGGCTTCTTCCGTGCATCCCCTCAGCCACGAAG GCCAAGTTCCCAGGGTTGGGCCAGTCCCCAGGTAGCAGGGAGACGCCCTG ATCCTTTCCTTCGGTCCCTCGGGGCCGAGGCCAGCAGGGCCAAGGGCCTT GGGGAACGGGGGACTCCTCACGGGCCCCGCCTGGAGCCTGACCCTCAG CACCCGGGCGCCTGCTGCTGAGCAACGGCCCCCATGCCAGCTC CCTCTGGAGCCTCTTTGCTCCCAGTAGCCCTATTCCAAGATGTTCTGGGGA GAGTGAACAGCTAAGAGCCTGCAGCCAAGCGCCCTGCCCCCTGAGCAGC CAGACCCCGGGCCCTGCAGTGCGCAGCCTTTAACTCCCAGGAATTCATG GGCCAGCTGTATCAGTGGGAGCCCTTCACTGAAGTCCAGGGCTCCCAGCG CTGTGAACTGCCGGCCCCGTGGCTTCCGCTTCTATGTCCGTCACAC TGAAAAGGTCCAGGATGGGACCCTGTGTCAGCCTGGAGCCCCTGACATCT

GTGTGGCTGGACGCTGTCTGAGCCCCGGCTGTGATGGGATCCTTGGCTCTG GCAGGCGTCCTGATGGCTGTGGAGTCTGTGGGGGTGATGATTCTACCTGTC GCCTTGTTTCGGGGAACCTCACTGACCGAGGGGGCCCCCTGGGCTATCAG AAGATCTTGTGGATTCCAGCGGGAGCCTTGCGGCTCCAGATTGCCCAGCT CCGGCCTAGCTCCAACTACCTGGCACTTCGTGGCCCTGGGGGCCGGTCCAT CATCAATGGGAACTGGGCTGTGGATCCCCCTGGGTCCTACAGGGCCGGCG GGACCGTCTTTCGATATAACCGTCCTCCCAGGGAGGAGGGCAAAGGGGAG AGTCTGTCGGCTGAAGGCCCCACCACCAGCCTGTGGATGTCTATATGATC TTTCAGGAGGAAAACCCAGGCGTTTTTTATCAGTATGTCATCTTCACCT CCTCCAATCCTTGAGAACCCCACCCCAGAGCCCCCTGTCCCCCAGCTTCAG CCGGAGATTCTGAGGGTGGAGCCCCACTTGCTCCGGCACCCCGCCAGC CCGGACCCCAGGCACCCTCCAGCGTCAGGTGCGGATCCCCCAGATGCCCG CCCCGCCCATCCCAGGACACCCCTGGGGTCTCCAGCTGCGTACTGGAAA CGAGTGGGACACTCTGCATGCTCAGCGTCCTGCGGGAAAGGTGTCTGGCG CCCCATTTCCTCTGCATCTCCCGTGAGTCGGGAGAGGAACTGGATGAAC GCAGCTGTGCCGGGTGCCAGGCCCCAGCCTCCCTGAACCCTGCCAC GGCACCCCATGCCCCCATACTGGGAGGCTGGCGAGTGGACATCCTGCAG CCGCTCCTGTGGCCCCGGCACCCAGCACCGCCAGCTGCAGTGCCGGCAGG AATTTGGGGGGGGTGCCCCCGGAGCGCTGTGGACATCTC CCCCGGCCCAACATCACCCAGTCTTGCCAGCTGCGCCTCTGTGGCCATTGG GAAGTTGGCTCTCCTTGGAGCCAGTGCTCCGTGCGGTGCGGCCGGGGCCA GAGAAGCCGGCAGGTTCGCTGTTGTTGGGAACAACGGTGATGAAGTGAGC GAGCAGGAGTGTGCGTCAGGCCCCCGCAGCCCCCCAGCAGAGAGGCCTG TGACATGGGGCCCTGTACTACTGCCTGGTTCCACAGCGACTGGAGCTCCA AGTGCTCAGCCGAGTGTGGGACGGGAATCCAGCGGCGCTCTGTGGTCTGC CTTGGGAGTGGGCAGCCCTCGGGCCAGGCCAGGGGAAGCAGGAGCAG GAACTGGGCAGAGCTGTCCAACAGGAAGCCGGCCCCCTGACATGCGCGCC TGCAGCCTGGGGCCCTGTGAGAGAACTTGGCGCTGGTACACAGGGCCCTG GGGTGAGTGCTCCTCCGAATGTGGCTCTGGCACACAGCGTAGAGACATCA TCTGTGTATCCAAACTGGGGACGGAGTTCAACGTGACTTCTCCGAGCAAC TGTTCTCACCTCCCAGGCCCCTGCCCTGCAGCCCTGTCAAGGGCAGGCC TGCCAGGACCGATGTTTTCCACGCCCTGGAGCCCATGTTCTCGCTCCTGC CCCTCAGCACCCGATGCCCTCCAACTGCGGCCCTCCAGGAAGCGCCCCT GTAACAGCCAACCCTGCAGCCAGCGCCCTGATGATCAATGCAAGGACAGC TCTCCACATTGCCCCCTGGTGGTACAGGCCCGGCTCTGCGTCTACCCCTAC TACACAGCCACCTGTTGCCGCTCTTGCGCACATGTCCTGGAGCGGTCTCCC CAGGATCCCTCCTGAAAGGGGTCCGGGGCACCTTCACGGTTTTCTGTGCCA CCATCGGTCACCCATTGATCGGCCCACTCTGAACCCCCTGGCTCTCCAGCC TGTCCCAGTCTCAGCAGGGATGTCCTCCAGGTGACAGAGGGTGGCAAGGT GACTGACACAAAGTGACTTTCAGGGCTGTGGTCAGGCCCATGTGGTGGTG TGATGGGTGTGCACATATGCCTCAGGTGTGCTTTTGGGACTGCATGGAT ATGTGTGTGCTCAAACGTGTATCACTTTTCAAAAAGAGGTTACACAGACT GAGAAGGACAAGACCTGTTTCCTTGAGACTTTCCTAGGTGGAAAGGAAAG CAAGTCTGCAGTTCCTTGCTAATCTGAGCTACTTAGAGTGTGGTCTCCCCA CCAACTCCAGTTTTGTGCCCTAAGCCTCATTTCTCATGTTCAGACCTCACA TCTTCTAAGCCGCCCTGTGTCTCTGACCCCTTCTCATTTGCCTAGTATCTCT GCCCTGCCTCAATTAGCTAGGGCTGGGGTCAGCCACTGCCAATCCTG CCTTACTCAGGAAGGCAGGAGGAAAGAGACTGCCTCTCCAGAGCAAGGC CCAGCTGGGCAGAGGGTGAAAAAGAGAAATGTGAGCATCCGCTCCCCA

# Thrombospondin 2

>gi|40317627|ref|NM\_003247.2| Homo sapiens thrombospondin 2 (THBS2), mRNA | qPCR forward\_primer match [3558..3580] | qPCR reverse\_primer match [3682..3655] | qPCR probe match [3597..3623]

GAGGAGGAGACGCATCCAGTACAGAGGGGCTGGACTTGGACCCC TGCAGCAGCCCTGCACAGGAGAAGCGGCATATAAAGCCGCGCTGCCCGG GAGCCGCTCGGCCACGTCCACCGGAGCATCCTGCACTGCAGGGCCGGTCT CTCGCTCCAGCAGAGCCTGCGCCTTTCTGACTCGGTCCGGAACACTGAAA CCAGTCATCACTGCATCTTTTTGGCAAACCAGGAGCTCAGCTGCAGGAGG CAGGATGGTCTGGAGGCTGGTCCTGCTGGGTGTGGCCCAGCA CGCAAGCTGGTCACCAGGACAAAGACACGACCTTCGACCTTTTCAGTATC AGCAACATCAACCGCAAGACCATTGGCGCCAAGCAGTTCCGCGGGCCCGA CCCCGGCGTGCCGGCTTACCGCTTCGTGCGCTTTGACTACATCCCACCGGT GAACGCAGATGACCTCAGCAAGATCACCAAGATCATGCGGCAGAAGGAG GGCTTCTTCCTCACGGCCCAGCTCAAGCAGGACGGCAAGTCCAGGGGCAC GCTGTTGGCTCTGGAGGGCCCCGGTCTCTCCCAGAGGCAGTTCGAGATCG TCTCCAACGGCCCGCGGACACGCTGGATCTCACCTACTGGATTGACGGC ACCCGGCATGTGGTCTCCCTGGAGGACGTCGGCCTGGCTGACTCGCAGTG GAAGAACGTCACCGTGCAGGTGGCTGGCGAGACCTACAGCTTGCACGTGG GCTGCGACCTCATAGACAGCTTCGCTCTGGACGAGCCCTTCTACGAGCAC CTGCAGGCGGAAAAGAGCCGGATGTACGTGGCCAAAGGCTCTGCCAGAG AGAGTCACTTCAGGGGTTTGCTTCAGAACGTCCACCTAGTGTTTGAAAACT CTGTGGAAGATATTCTAAGCAAGAAGGGTTGCCAGCAAGGCCAGGGAGCT GAGATCAACGCCATCAGTGAGAACACAGAGACGCTGCGCCTGGGTCCGCA TGTCACCACCGAGTACGTGGGCCCCAGCTCGGAGAGGAGGCCCGAGGTGT GCGAACGCTCGTGCGAGGAGCTGGGAAACATGGTCCAGGAGCTCTCGGG GCTCCACGTCCTCGTGAACCAGCTCAGCGAGAACCTCAAGAGAGTGTCGA ATGATAACCAGTTTCTCTGGGAGCTCATTGGTGGCCCTCCTAAGACAAGG AACATGTCAGCTTGCTGGCAGGATGGCCGGTTCTTTGCGGAAAATGAAAC GTGGGTGGACAGCTGCACCACGTGTACCTGCAAGAAATTTAAAACCA TITGCCACCAAATCACCTGCCCGCCTGCAACCTGCGCCAGTCCATCCTTTG TGGAAGGCGAATGCTGCCTTCCTGCCTCCACTCGGTGGACGGTGAGGAG GGCTGGTCTCCGTGGGCAGAGTGGACCCAGTGCTCCGTGACGTGTGGCTC TGGGACCCAGCAGAGAGGCCGGTCCTGTGACGTCACCAGCAACACCTGCT TGGGGCCCTCCATCCAGACACGGGCTTGCAGTCTGAGCAAGTGTGACACC CGCATCCGGCAGGACGGCGGCTGGAGCCACTGGTCACCTTGGTCTTCATG CTCTGTGACCTGTGGAGTTGGCAATATCACACGCATCCGTCTCTGCAACTC CCCAGTGCCCAGATGGGGGGCAAGAATTGCAAAGGGAGTGGCCGGGAG ACCAAAGCCTGCCAGGGCGCCCCATGCCCAATCGATGGCCGCTGGAGCCC CTGGTCCCGTGGTCGGCCTGCACTGTCACCTGTGCCGGTGGGATCCGGG AGCGCACCCGGGTCTGCAACAGCCCTGAGCCTCAGTACGGAGGGAAGGCC TGCGTGGGGGATGTGCAGGAGCGTCAGATGTGCAACAAGAGGAGCTGCC CCGTGGATGGCTGTTTATCCAACCCCTGCTTCCCGGGAGCCCAGTGCAGCA GCTTCCCCGATGGGTCCTGGTCATGCGGCTCCTGCCCTGTGGGCTTCTTGG

GCAATGGCACCCACTGTGAGGACCTGGACGAGTGTGCCCTGGTCCCCGAC ATCTGCTTCTCCACCAGCAAGGTGCCTCGCTGTGTCAACACTCAGCCTGGC TTCCACTGCCTGCCCGCCCCGATACAGAGGGAACCAGCCCGTCGG GGTCGGCCTGGAAGCAGCCAAGACGGAAAAGCAAGTGTGTGAGCCCGAA AACCCATGCAAGGACAAGACACACACTGCCACAAGCACGCGGAGTGCA TCTACCTGGGCCACTTCAGCGACCCCATGTACAAGTGCGAGTGCCAGACA GGCTACGCGGCCACGGGCTCATCTGCGGGGAGGACTCGGACCTGGACG GCTGGCCCAACCTCAATCTGGTCTGCGCCACCAACGCCACCTACCACTGC ATCAAGGATAACTGCCCCATCTGCCAAATTCTGGGCAGGAAGACTTTGA CAAGGACGGATTGCCTGTGATGACGATGACAATGACGGTG TGACCGATGAGAAGGACAACTGCCAGCTCCTCTTCAATCCCCGCCAGGCT GACTATGACAAGGATGAGGTTGGGGACCGCTGTGACAACTGCCCTTACGT GCACAACCCTGCCCAGATCGACACAGACAACAATGGAGAGGGTGACGCC TGCTCCGTGGACATTGATGGGGACGATGTCTTCAATGAACGAGACAATTG TCCCTACGTCTACAACACTGACCAGAGGGACACGGATGGTGACGGTGTGG GGGATCACTGTGACAACTGCCCCCTGGTGCACAACCCTGACCAGACCGAC GTGGACAATGACCTTGTTGGGGACCAGTGTGACAACAACGAGGACATAGA TGACGACGCCACCAGAACAACCAGGACAACTGCCCCTACATCTCCAACG CCAACCAGGCTGACCATGACAGAGACGCCAGGGCGACGCCTGTGACCCT GATGATGACAACGATGGCGTCCCCGATGACAGGGACAACTGCCGGCTTGT GTTCAACCCAGACCAGGAGGACTTGGACGGTGATGGACGGGGTGATATTT GTAAAGATGATTTTGACAATGACAACATCCCAGATATTGATGATGTGTGT CCTGAAAACAATGCCATCAGTGAGACAGACTTCAGGAACTTCCAGATGGT CCCCTTGGATCCCAAAGGGACCACCCAAATTGATCCCAACTGGGTCATTC GCCATCAAGGCAAGGAGCTGGTTCAGACAGCCAACTCGGACCCCGGCATC GCTGTAGGTTTTGACGAGTTTGGGTCTGTGGACTTCAGTGGCACATTCTAC GTAAACACTGACCGGGACGACGACTATGCCGGCTTCGTCTTTGGTTACCA GTCAAGCAGCCGCTTCTATGTGGTGATGTGGAAGCAGGTGACGCAGACCT ACTGGGAGGACCAGCCCACGCGGGCCTATGGCTACTCCGGCGTGTCCCTC AAGGTGGTGAACTCCACCACGGGGACGGGCGAGCACCTGAGGAACGCGC TGTGGCACACGGGGAACACGCCGGGGCAGGTGCGAACCTTATGGCACGA CCCCAGGAACATTGGCTGGAAGGACTACACGGCCTATAGGTGGCACCTGA CTCACAGGCCCAAGACTGGCTACATCAGAGTCTTAGTGCATGAAGGAAAA CAGGTCATGGCAGACTCAGGACCTATCTATGACCAAACCTACGCTGGCGG GCGCTGGGTCTATTTGTCTTCTCAAGAAATGGTCTATTTCTCAGACCT CAAGTACGAATGCAGAGATATTTAAACAAGATTTGCTGCATTTCCGGCAA TGCCCTGTGCATGCCATGGTCCCTAGACACCTCAGTTCATTGTGGTCCTTG TGGCTTCTCTCTAGCAGCACCTCCTGTCCCTTGACCTTAACTCTGATGGT TCTTCACCTCCTGCCAGCAACCCCAAACCCAAGTGCCTTCAGAGGATAAA TATCAATGGAACTCAGAGATGAACATCTAACCCACTAGAGGAAACCAGTT TGGTGATATATGAGACTTTATGTGGAGTGAAAATTGGGCATGCCATTACA TTGCTTTTCTTGTTTAAAAAGAATGACGTTTACATATAAAATGTAA TTACTTATTGTATTTATGTGTATATGGAGTTGAAGGGAATACTGTGCATAA GCCATTATGATAAATTAAGCATGAAAAATATTGCTGAACTACTTTTGGTGC TTAAAGTTGTCACTATTCTTGAATTAGAGTTGCTCTACAATGACACAAA TCCCATTAAATAAATTATAAACAAGGGTCAATTCAAATTTGAAGTAATGTT TTAGTAAGGAGAGATTAGAAGACAACAGGCATAGCAAATGACATAAGCT ACCGATTAACTAATCGGAACATGTAAAACAGTTACAAAAAATAAACGAACT CTCCTCTTGTCCTACAATGAAAGCCCTCATGTGCAGTAGAGATGCAGTTTC ATCAAAGAACAACATCCTTGCAAATGGGTGTGACGCGGTTCCAGATGTG

GATTTGGCAAAACCTCATTTAAGTAAAAGGTTAGCAGAGCAAAGTGCGGT CTTCCTTCCCAGCTTTGCTGCCTGAGAGGAACCAGAGCAGACGCACAGG CCGGAAAAGGCGCATCTAACGCGTATCTAGGCTTTGGTAACTGCGGACAA GTTGCTTTTACCTGATTTGATGATACATTTCATTAAGGTTCCAGTTATAAAT ATTTTGTTAATATTTAATTAAGTGACTATAGAATGCAACTCCATTTACCAGT AACTTATTTTAAATATGCCTAGTAACACATATGTAGTATAATTTCTAGAAA CAAACATCTAATAAGTATATAATCCTGTGAAAATATGAGGCTTGATAATA TTAGGTTGTCACGATGAAGCATGCTAGAAGCTGTAACAGAATACATAGAG AATAATGAGGAGTTTATGATGGAACCTTAAATATATAATGTTGCCAGCGA TTTTAGTTCAATATTTGTTACTGTTATCTATCTGCTGTATATGGAATTCTTT TAATTCAAACGCTGAAAAGAATCAGCATTTAGTCTTGCCAGGCACACCCA GTTGGTTGGTTTTTTTTGCTTTAAGTTGCATGATCTTTCTGCAGGAAAT CTGATGATGGATAGGGGGCAAATCTTTTTCCCCTTTCTGTTAATAGTCATC ACATTTCTATGCCAAACAGGAACAATCCATAACTTTAGTCTTAATGTACAC ATTGCATTTTGATAAAATTAATTTTGTTGTTTCCTTTGAGGTTGATCGTTGT GTTGTTGTTTTGCTGCACTTTTTACTTTTTTGCGTGTGGAGCTGTATTCCCG AGACCAACGAAGCGTTGGGATACTTCATTAAATGTAGCGACTGTCAACAG CGTGCAGGTTTTCTGTTTCTGTGTGGGGTCAACCGTACAATGGTGTGG GAGTGACGATGTGAATATTTAGAATGTACCATATTTTTTGTAAATTAT TTATGTTTTCTAAACAAATTTATCGTATAGGTTGATGAAACGTCATGTGT TTTGCCAAAGACTGTAAATATTTATTTATGTGTTCACATGGTCAAAATTTC ACCACTGAAACCCTGCACTTAGCTAGAACCTCATTTTTAAAGATTAACAAC AAAA **SEO ID NO: 94** 

### Adlican

>gi|18390318|ref|NM\_015419.1| Homo sapiens adlican (DKFZp564I1922), mRNA | qPCR assay on demand context match [694..718]

ATGCCCAAGCGCGCGCACTGGGGGGCCCTCTCCGTGGTGCTGATCC TGCTTTGGGGCCATCCGCGAGTGGCGCTGGCCTGCCCGCATCCTTGTGCCT GCTACGTCCCAGCGAGGTCCACTGCACGTTCCGATCCCTGGCTTCCGTGC CCGCTGGCATTGCTAGACACGTGGAAAGAATCAATTTGGGGTTTAATAGC ATACAGGCCCTGTCAGAAACCTCATTTGCAGGACTGACCAAGTTGGAGCT ACTTATGATTCACGGCAATGAGATCCCAAGCATCCCCGATGGAGCTTTAA GAGACCTCAGCTCTCTCAGGTTTTCAAGTTCAGCTACAACAAGCTGAGA GTGATCACAGGACAGACCCTCCAGGGTCTCTCTAACTTAATGAGGCTGCA CATTGACCACAACAAGATCGAGTTTATCCACCCTCAAGCTTTCAACGGCTT AACGTCTCTGAGGCTACTCCATTTGGAAGGAAATCTCCTCCACCAGCTGCA CCCCAGCACCTTCTCCACGTTCACATTTTTGGATTATTTCAGACTCTCCACC ATAAGGCACCTCTACTTAGCAGAGAACATGGTTAGAACTCTTCCTGCCAG CATGCTTCGGAACATGCCGCTTCTGGAGAATCTTTACTTGCAGGGAAATCC GTGGACCTGCGATTGTGAGATGAGATGGTTTTTGGAATGGGATGCAAAAT CCAGAGGAATTCTGAAGTGTAAAAAGGACAAAGCTTATGAAGGCGGTCA GTTGTGTGCAATGTGCTTCAGTCCAAAGAAGTTGTACAAACATGAGATAC ACAAGCTGAAGGACATGACTTGTCTGAAGCCTTCAATAGAGTCCCCTCTG

AGACAGAACAGGAGCAGGAGTATTGAGGAGGAGCAAGAACAGGAAGAG GATGGTGGCAGCCAGCTCATCCTGGAGAAATTCCAACTGCCCCAGTGGAG CATCTCTTTGAATATGACCGACGAGCACGGGAACATGGTGAACTTGGTCT GTGACATCAAGAAACCAATGGATGTGTACAAGATTCACTTGAACCAAACG GATCCTCCAGATATTGACATAAATGCAACAGTTGCCTTGGACTTTGAGTGT CCAATGACCCGAGAAAACTATGAAAAGCTATGGAAATTGATAGCATACTA CAGTGAAGTTCCCGTGAAGCTACACAGAGAGCTCATGCTCAGCAAAGACC CCAGAGTCAGCTACCAGTACAGGCAGGATGCTGATGAGGAAGCTCTTTAC TACACAGGTGTGAGAGCCCAGATTCTTGCAGAACCAGAATGGGTCATGCA GCCATCCATAGATATCCAGCTGAACCGACGTCAGAGTACGGCCAAGAAGG TGCTACTTCCTACTACACCCAGTATTCTCAAACAATATCCACCAAAGATA CAAGGCAGGCTCGGGCAGAAGCTGGGTAATGATTGAGCCTAGTGGAGCT GTGCAAAGAGATCAGACTGTCCTGGAAGGGGGTCCATGCCAGTTGAGCTG CAACGTGAAAGCTTCTGAGAGTCCATCTATCTTCTGGGTGCTTCCAGATGG CTCCATCCTGAAAGCGCCCATGGATGACCCAGACAGCAAGTTCTCCATTCT CAGCAGTGGCTGAGGATCAAGTCCATGGAGCCATCTGACTCAGGCT TGTACCAGTGCATTGCTCAAGTGAGGGATGAAATGGACCGCATGGTATAT AGGGTACTTGTGCAGTCTCCCTCCACTCAGCCAGCCGAGAAAGACACAGT GACAATTGGCAAGAACCCAGGGGAGTCGGTGACATTGCCTTGCAATGCTT TAGCAATACCCGAAGCCCACCTTAGCTGGATTCTTCCAAACAGAAGGATA ATTAATGATTTGGCTAACACATCACATGTATACATGTTGCCAAATGGAACT CTTTCCATCCCAAAGGTCCAAGTCAGTGATAGTGGTTACTACAGATGTGTG GCTGTCAACCAGCAAGGGGCAGACCATTTTACGGTGGGAATCACAGTGAC CAAGAAAGGGTCTGGCTTGCCATCCAAAAGAGGCAGACGCCCAGGTGCA AAGGCTCTTTCCAGAGTCAGAGAAGACATCGTGGAGGATGAAGGGGGCTC GGGCATGGGAGATGAAGAGACACTTCAAGGAGACTTCTGCATCCAAAG GACCAAGAGGTGTTCCTCAAAACAAAGGATGATGCCATCAATGGAGACA AGAAAGCCAAGAAAGGGAGAAGAAAGCTGAAACTCTGGAAGCATTCGGA AAAAGAACCAGAGACCAATGTTGCAGAAGGTCGCAGAGTGTTTGAATCTA GACGAAGGATAAACATGGCAAACAAACAGATTAATCCGGAGCGCTGGGC TGATATTTTAGCCAAAGTCCGTGGGAAAAATCTCCCTAAGGGCACAGAAG TACCCCGTTGATTAAAACCACAAGTCCTCCATCCTTGAGCCTAGAAGTCA CACCACCTTTTCCTGCTGTTTCTCCCCCCTCAGCATCTCCTGTGCAGACAGT AACCAGTGCTGAAGAATCCTCAGCAGATGTACCTCTACTTGGTGAAGAAG AGCACGTTTTGGGTACCATTTCCTCAGCCAGCATGGGGCTAGAACACAAC CACAATGGAGTTATTCTTGTTGAACCTGAAGTAACAAGCACACCTCTGGA GGAAGTTGTTGATGACCTTTCTGAGAAGACTGAGGAGATAACTTCCACTG AAGGAGACCTGAAGGGGACAGCAGCCCCTACACTTATATCTGAGCCTTAT GAACCATCTCCTACTCTGCACACATTAGACACAGTCTATGAAAAGCCCAC CCATGAAGAGACGCAACAGAGGGTTGGTCTGCAGCAGATGTTGGATCGT CACCAGAGCCCACATCCAGTGAGTATGAGCCTCCATTGGATGCTGTCTCCT TGGCTGAGTCTGAGCCCATGCAATACTTTGACCCAGATTTGGAGACTAAG TCACAACCAGATGAGGATAAGATGAAAGAAGACACCTTTGCACACCTTAC TCCAACCCCACCATCTGGGTTAATGACTCCAGTACATCACAGTTATTTGA GGATTCTACTATAGGGGAACCAGGTGTCCCAGGCCAATCACATCTACAAG GACTGACAGACAACATCCACCTTGTGAAAAGTAGTCTAAGCACTCAAGAC ACCTTACTGATTAAAAAGGGTATGAAAGAGATGTCTCAGACACTACAGGG AGGAAATATGCTAGAGGGAGACCCCACACACTCCAGAAGTTCTGAGAGTG AGGGCCAAGAGAGCAAATCCATCACTTTGCCTGACTCCACACTGGGTATA ATGAGCAGTATGTCTCCAGTTAAGAAGCCTGCGGAAACCACAGTTGGTAC

CCTCCTAGACAAGACACCACAACAGTAACAACAACACCAAGGCAAAAA GTTGCTCCGTCATCCACCATGAGCACTCACCCTTCTCGAAGGAGACCCAAC GGGAGAAGGAGATTACGCCCCAACAAATTCCGCCACCGGCACAAGCAAA **AAGCACCTGACATTAAGATTTCAAGTCAAGTGGAGAGTTCTCTGGTTCCTA** CAGCTTGGGTGGATAACACAGTTAATACCCCCAAACAGTTGGAAATGGAG AAGAATGCAGAACCCACATCCAAGGGAACACCACGGAGAAAACACGGGA AGAGGCCAAACAACATCGATATACCCCTTCTACAGTGAGCTCAAGAGCG TCCGGATCCAAGCCCAGCCCTTCTCCAGAAAATAAACATAGAAACATTGT TACTCCCAGTTCAGAAACTATACTTTTGCCTAGAACTGTTTCTCTGAAAAC TGAGGGCCCTTATGATTCCTTAGATTACATGACAACCACCAGAAAAATAT ATTCATCTTACCCTAAAGTCCAAGAGACACTTCCAGTCACATATAAACCCA CATCAGATGGAAAAGAAATTAAGGATGATGTTGCCACAAATGTTGACAAA CATAAAAGTGACATTTTAGTCACTGGTGAATCAATTACTAATGCCATACCA ACTTCTCGCTCCTTGGTCTCCACTATGGGAGAATTTAAGGAAGAATCCTCT CCTGTAGGCTTTCCAGGAACTCCAACCTGGAATCCCTCAAGGACGCCCA GCCTGGGAGGCTACAGACAGACATACCTGTTACCACTTCTGGGGAAAATC TTACAGACCCTCCCCTTCTTAAAGAGCTTGAGGATGTGGATTTCACTTCCG AGTTTTTGTCCTCTTTGACAGTCTCCACACCATTTCACCAGGAAGAAGCTG GTTCTTCCACAACTCTCTCAAGCATAAAAGTGGAGGTGGCTTCAAGTCAG GCAGAAACCACCACCTTGATCAAGATCATCTTGAAACCACTGTGGCTAT TCTCCTTCTGAAACTAGACCACAGAATCACACCCCTACTGCTGCCCGGAT GAAGGAGCCAGCATCCTCGTCCCCATCCACAATTCTCATGTCTTTGGGACA AACCACCACCACTAAGCCAGCACTTCCCAGTCCAAGAATATCTCAAGCAT CTAGAGATTCCAAGGAAAATGTTTTCTTGAATTATGTGGGGAATCCAGAA ACAGAAGCAACCCCAGTCAACAATGAAGGAACACAGCATATGTCAGGGC CAAATGAATTATCAACACCCTCTTCCGACCGGGATGCATTTAACTTGTCTA CAAAGCTGGAATTGGAAAAGCAAGTATTTGGTAGTAGGAGTCTACCACGT GGCCCAGATAGCCAACGCCAGGATGGAAGAGTTCATGCTTCTCATCAACT AACCAGAGTCCCTGCCAAACCCATCCTACCAACAGCAACAGTGAGGCTAC CTGAAATGTCCACACAAAGCGCTTCCAGATACTTTGTAACTTCCCAGTCAC CTCGTCACTGGACCAACAAACCGGAAATAACTACATATCCTTCTGGGGCT TTGCCAGAGAACAACAGTTTACAACTCCAAGATTATCAAGTACAACAAT TCCTCTCCCATTGCACATGTCCAAACCCAGCATTCCTAGTAAGTTTACTGA CCGAAGAACTGACCAATTCAATGGTTACTCCAAAGTGTTTGGAAATAACA ACATCCCTGAGGCAAGAAACCCAGTTGGAAAGCCTCCCAGTCCAAGAATT TTCCACAGTTGGGAGTCACCCGGAGACCCCAGATACCCACTTCTCCTGCCC CAGTAATGAGAGAGAAAAGTTATTCCAGGTTCCTACAACAGGATACAT TCCCATAGCACCTTCCATCTGGACTTTGGCCCTCCGGCACCTCCGTTGTTG CACACTCCGCAGACCACGGGATCACCCTCAACTAACTTACAGAATATCCC TATGGTCTCTCCACCCAGAGTTCTATCTCCTTTATAACATCTTCTGTCCAG TCCTCAGGAAGCTTCCACCAGAGCAGCTCAAAGTTCTTTGCAGGAGGACC TCCTGCATCCAAATTCTGGTCTCTTGGGGAAAAGCCCCAAATCCTCACCAA GTCCCACAGACTGTGTCCGTCACCGCTGAGACAGACACTGTGTTCCCCTG TGAGGCAACAGGAAAACCAAAGCCTTTCGTTACTTGGACAAAGGTTTCCA CAGGAGCTCTTATGACTCCGAATACCAGGATACAACGGTTTGAGGTTCTC AAGAACGGTACCTTAGTGATACGGAAGGTTCAAGTACAAGATCGAGGCCA GTATATGTGCACCGCCAGCAACCTGCACGGCCTGGACAGGATGGTGGTCT TGCTTTCGGTCACCGTGCAGCAACCTCAAATCCTAGCCTCCCACTACCAGG

ACGTCACTGTCTACCTGGGAGACACCATTGCAATGGAGTGTCTGGCCAAA GGGACCCCAGCCCCCAAATTTCCTGGATCTTCCCTGACAGGAGGGTGTG GCAAACTGTGTCCCCCGTGGAGAGCCGCATCACCCTGCACGAAAACCGGA CCCTTTCCATCAAGGAGGCGTCCTTCTCAGACAGAGGCGTCTATAAGTGC GTGGCCAGCAATGCAGCCGGGGGGGGACAGCCTGGCCATCCGCCTGCACGT GGCGGCACTGCCCCCGTTATCCACCAGGAGAAGCTGGAGAACATCTCGC TGCCCCGGGGCTCAGCATTCACATTCACTGCACTGCCAAGGCTGCGCCCC TGCCCAGCGTGCGCTGGGTGCTCGGGGACGGTACCCAGATCCGCCCCTCG CAGTTCCTCCACGGGAACTTGTTTTCTTTTCCCCAACGGGACGCTCTACATC CGCAACCTCGCGCCCAAGGACAGCGGGCGCTATGAGTGCGTGGCCGCCAA CCTGGTAGGCTCCGCGCGCAGGACGGTGCAGCTGAACGTGCAGCGTGCAG GTACGGAGGAACCCTCAAGCTGGACTGCAGCGCCTCGGGGGACCCCTGGC CGCGCATCCTCTGGAGGCTGCCGTCCAAGAGGATGATCGACGCGCTCTTC AGTTTTGATAGCAGAATCAAGGTGTTTGCCAATGGGACCCTGGTGGTGAA ATCAGTGACGGACAAAGATGCCGGAGATTACCTGTGCGTAGCTCGAAATA AGGTTGGTGATGACTACGTGGTGCTCAAAGTGGATGTGGTGATGAAACCG GCCAAGATTGAACACAAGGAGGAGAACGACCACAAAGTCTTCTACGGGG GTGACCTGAAAGTGGACTGTGTGGCCACCGGGCTTCCCAATCCCGAGATC TCCTGGAGCCTCCCAGACGGGAGTCTGGTGAACTCCTTCATGCAGTCGGA TGACAGCGGTGGACGCACCAAGCGCTATGTCGTCTTCAACAATGGGACAC TCTACTTTAACGAAGTGGGGATGAGGAGGAGGAGGACTACACCTGCTTT GCTGAAAATCAGGTCGGGAAGGACGAGATGAGAGTCAAGGTGG CCCTATGGAGACGTGGTCACTGTAGCCTGTGAGGCCAAAGGAGAACCCAT GCCCAAGGTGACTTGGTTGTCCCCAACCAACAAGGTGATCCCCACCTCCTC TGAGAAGTATCAGATATACCAAGATGGCACTCTCCTTATTCAGAAAGCCC AGCGTTCTGACAGCGCGAACTACACCTGCCTGGTCAGGAACAGCGCGGGA GAGGATAGGAAGACGTGTGGATTCACGTCAACGTCCAGCCACCCAAGAT CAACGGTAACCCCAACCCCATCACCACCGTGCGGAGATAGCAGCCGGGG GCAGTCGGAAACTGATTGACTGCAAAGCTGAAGGCATCCCCACCCCGAGG GTGTTATGGGCTTTTCCCGAGGGTGTGGTTCTGCCAGCTCCATACTATGGA AACCGGATCACTGTCCATGGCAACGGTTCCCTGGACATCAGGAGTTTGAG GAGGCGAGGTTGATCGTGCAGCTCACTGTCCTGGAGCCCATGGAGAAACC CATCTTCCACGACCCGATCAGCGAGAAGATCACGGCCATGGCGGCCACA CCATCAGCCTCAACTGCTCTGCCGCGGGGACCCCGACACCCAGCCTGGTG TGGGTCCTTCCCAATGGCACCGATCTGCAGAGTGGACAGCAGCTGCAGCG CTTCTACCACAAGGCTGACGGCATGCTACACATTAGCGGTCTCTCCTCGGT GGACGCTGGGCCTACCGCTGCGTGGCCCAATGCCGCTGGCCACACGG AGAGGCTGGTCTCCCTGAAGGTGGGACTGAAGCCAGAAGCAACAAGCA GTATCATAACCTGGTCAGCATCATCAATGGTGAGACCCTGAAGCTCCCCT GCACCCCCCGGGGCTGGGCAGGGACGTTTCTCCTGGACGCTCCCCAAT GGCATGCATCTGGAGGGCCCCCAAACCCTGGGACGCGTTTCTCTTCTGGA CAATGGCACCCTCACGGTTCGTGAGGCCTCGGTGTTTGACAGGGGTACCT ATGTATGCAGGATGGAGACGGAGTACGGCCCTTCGGTCACCAGCATCCCC GTGATTGTGATCGCCTATCCTCCCGGATCACCAGCGAGCCCACCCCGGTC ATCTACACCCGGCCCGGGAACACCGTGAAACTGAACTGCATGGCTATGGG GATTCCCAAAGCTGACATCACGTGGGAGTTACCGGATAAGTCGCATCTGA AGGCAGGGGTTCAGGCTCGTCTGTATGGAAACAGATTTCTTCACCCCCAG

GGATCACTGACCATCCAGCATGCCACACAGAGAGATGCCGGCTTCTACAA GTGCATGGCAAAAAACATTCTCGGCAGTGACTCCAAAACAACTTACATCC ACGTCTTCTGAAATGTGGATTCCAGAATGATTGCTTAGGAACTGACAACA AAGCGGGGTTTGTAAGGGAAGCCAGGTTGGGGAATAGGAGCTCTTAAATA ATGTGTCACAGTGCATGGTGGCCTCTGGTGGGTTTCAAGTTGAGGTTGATC TTGATCTACAATTGTTGGGAAAAGGAAGCAATGCAGACACGAGAAGGAG GGCTCAGCCTTGCTGAGACACTTTCTTTTGTGTTTACATCATGCCAGGGGC TTCATTCAGGGTGTCTGTGCTCTGACTGCAATTTTTCTTCTTTTGCAAATGC CACTCGACTGCCTTCATAAGCGTCCATAGGATATCTGAGGAACATTCATCA AAAATAAGCCATAGACATGAACACACCTCACTACCCCATTGAAGACGCA TCACCTAGTTAACCTGCTGCAGTTTTTACATGATAGACTTTGTTCCAGATT GACAAGTCATCTTTCAGTTATTTCCTCTGTCACTTCAAAACTCCAGCTTGC TCAAATCAGACGATGAGACTAGAAGGAGAAATACTTTCTGTCTTATTAAA ATTAATAAATTATTGGTCTTTACAAGACTTGGATACATTACAGCAGACATG GAAATATAATTTTAAAAAATTTCTCTCCAACCTCCTTCAAATTCAGTCACC ACTGTTATATTACCTTCTCCAGGAACCCTCCAGTGGGGAAGGCTGCGATAT TAGATTTCCTTGTATGCAAAGTTTTTGTTGAAAGCTGTGCTCAGAGGAGGT GAGAGGAGGAAGGAGAAAACTGCATCATAACTTTACAGAATTGAATC TAGAGTCTTCCCGAAAAGCCCAGAAACTTCTCTGCAGTATCTGGCTTGTC CATCTGGTCTAAGGTGGCTGCTTCTTCCCCAGCCATGAGTCAGTTTGTGCC CATGAATAATACACGACCTGTTATTTCCATGACTGCTTTACTGTATTTTTA AGGTCAATACTGTACATTTGATAATAAAATAATATTCTCCCAAAAAA AAA **SEO ID NO: 95** 

#### Cystatin SA

>gi|19882252|ref|NM\_001322.2| Homo sapiens cystatin SA (CST2), mRNA | qPCR forward\_primer match [302..320] | qPCR reverse\_primer match [393..370] | qPCR probe match [341..369]

**SEO ID NO: 96** 

## Cystatin SN

>gi|19882250|ref|NM\_001898.2| Homo sapiens cystatin SN (CST1), mRNA | qPCR forward\_primer match [358..376] | qPCR reverse\_primer match [449..426] | qPCR probe match [397..425]

GGGCTCCCTGCCTCGGGCTCTCACCCTCCTCCTCCTGCAGCTCCAGCT TTGTGCTCTGCCTCTGAGGAGACCATGGCCCAGTATCTGAGTACCCTGCTG CTCCTGCTGGCCACCCTAGCTGTGGCCCTGGCCTGGAGCCCCAAGGAGGA GGATAGGATAATCCCGGGTGGCATCTATAACGCAGACCTCAATGATGAGT GGGTACAGCGTGCCCTTCACTTCGCCATCAGCGAGTATAACAAGGCCACC AAAGATGACTACAGACGTCCGCTGCGGGTACTAAGAGCCAGGCAACA GACCGTTGGGGGGGTGAATTACTTCTTCGACGTAGAGGTGGGCCGCACCA TATGTACCAAGTCCCAGCCCAACTTGGACACCTGTGCCTTCCATGAACAGC CAGAACTGCAGAAGAACAGTTGTGCTCTTTCGAGATCTACGAAGTTCCC TGGGAGAACAGAAGGTCCCTGGTGAAATCCAGGTGTCAAGAATCCTAGGG **ATCTGTGCCAGGCCATTCGCACCAGCCACCACCCACTCCCACCCCTGTAG** TGCTCCCACCCTGGACTGGTGGCCCCCACCCTGCGGGAGGCCTCCCCATG TGCCTGCGCCAAGAGACAGACAGAGAGGCTGCAGGAGTCCTTTGTTGCT CAGCAGGGCGCTCTCCCTCCTTCCTTCTTGCTTCTAATAGCCCTGGT ACATGGTACACACCCCCCACCTCCTGCAATTAAACAGTAGCATCGCCTCC **SEO ID NO: 97** CTCTGAAAAAAAAAAAAAAAAAAAAA

### Lysyl Oxidase-Like Enzyme 2

>gi|4505010|ref|NM\_002318.1| Homo sapiens lysyl oxidase-like 2 (LOXL2), mRNA | qPCR forward\_primer match [2205..2223] | qPCR reverse\_primer match [2286..2269] | qPCR probe match [2261..2229]

ACTCCAGCGCGCGGCTACCTACGCTTGGTGCTTGCTTTCTCCAGCCA TCGGAGACCAGAGCCGCCCCTCTGCTCGAGAAAGGGGCTCAGCGGCGGC GGAAGCGGAGGGGACCACCGTGGAGAGCGCGGTCCCAGCCCGGCCACT GCGGATCCTGAAACCAAAAGCTCCTGCTGCTTCTGTACCCCGCCTGTCC CTCCCAGCTGCGCAGGGCCCCTTCGTGGGATCATCAGCCCGAAGACAGGG ATGGAGAGGCCTCTGTGCTCCCACCTCTGCAGCTGCCTGGCTATGCTGGCC CTCCTGTCCCCCTGAGCCTGGCACAGTATGACAGCTGGCCCCATTACCCC GAGTACTTCCAGCAACCGGCTCCTGAGTATCACCAGCCCCAGGCCCCCGC CAACGTGGCCAAGATTCAGCTGCGCCTGGCTGGCAGAAGAGGAAGCAC AGCGAGGCCGGGTGGAGGTGTACTATGATGGCCAGTGGGGCACCGTGTG CGATGACGACTTCTCCATCCACGCTGCCCACGTCGTCTGCCGGGAGCTGG GCTATGTGGAGGCCAAGTCCTGGACTGCCAGCTCCTCCTACGGCAAGGGA GAAGGCCCATCTGGTTAGACAATCTCCACTGTACTGGCAACGAGGCGAC CGGAGGATGTCGGTGTGCGCGACAAAAGGATTCCTGGGTTCAAA TTTGACAATTCGTTGATCAACCAGATAGAGAACCTGAATATCCAGGTGGA GGACATTCGGATTCGAGCCATCCTCTCAACCTACCGCAAGCGCACCCCAG TGATGGAGGGCTACGTGGAGGTGAAGGGGGCAAGACCTGGAAGCAGAT CTGTGACAAGCACTGGACGGCCAAGAATTCCCGCGTGGTCTGCGGCATGT TTGGCTTCCCTGGGGAGAGGACATACAATACCAAAGTGTACAAAATGTTT GCCTCACGGAGGAAGCAGCGCTACTGGCCATTCTCCATGGACTGCACCGG

CACAGAGGCCCACATCTCCAGCTGCAAGCTGGGCCCCCAGGTGTCACTGG TGTGTGCCTGGGCAGGTCTTCAGCCCTGACGGACCCTCGAGATTCCGGAA AGCATACAAGCCAGAGCAACCCCTGGTGCGACTGAGAGGCGGTGCCTACA TCGGGGAGGCCGCGTGGAGGTGCTCAAAAATGGAGAATGGGGGACCGT CTGCGACGACAGTGGGACCTGGTGTCGGCCAGTGTGGTCTGCAGAGAGC TGGGCTTTGGGAGTGCCAAAGAGGCAGTCACTGGCTCCCGACTGGGGCAA GGGATCGGACCCATCCACCTCAACGAGATCCAGTGCACAGGCAATGAGAA GTCCATTATAGACTGCAAGTTCAATGCCGAGTCTCAGGGCTGCAACCACG AGGAGGATGCTGTGTGAGATGCAACACCCCTGCCATGGGCTTGCAGAAG AAGCTGCGCCTGAACGGCCGCCGAATCCCTACGAGGGCCGAGTGGAGGT GCTGGTGGAGAAACGGGTCCCTTGTGTGGGGGATGGTGTGTGGCCAAA ACTGGGGCATCGTGGAGGCCATGGTGGTCTGCCGCCAGCTGGGCCTGGGA TTCGCCAGCAACGCCTTCCAGGAGACCTGGTATTGGCACGGAGATGTCAA CAGCAACAAGTGGTCATGAGTGGAGTGAAGTGCTCGGGAACGGAGCTG TCCCTGGCGCACTGCCCCCACGGGGAGGACGTGGCCTGCCCCAGGG CGGAGTGCAGTACGGGGCCGGAGTTGCCTGACAGAAACCGCCCCTGACC TGGTCCTCAATGCGGAGATGGTGCAGCAGCACCTACCTGGAGGACCGG CCCATGTTCATGCTGCAGTGTGCCATGGAGGAGAACTGCCTCTCGGCCTCA GCCGCGCAGACCCACCACGGGCTACCGCCGGCTCCTGCGCTTCTC CTCCCAGATCCACAACAATGGCCAGTCCGACTTCCGGCCCAAGAACGGCC GCCACGCGTGGATCTGGCACGACTGTCACAGGCACTACCACAGCATGGAG GTGTTCACCCACTATGACCTGCAACCTCAATGGCACCAAGGTGGCAGA GGGCCACAAGGCCAGCTTCTGCTTGGAGGACACAGAATGTGAAGGAGAC ATCCAGAAGAATTACGAGTGTGCCAACTTCGGCGATCAGGGCATCACCAT GGGCTGCTGGGACATGTACCGCCATGACATCGACTGCCAGTGGGTTGACA TCACTGACGTGCCCCTGGAGACTACCTGTTCCAGGTTGTTATTAACCCCA ACTTCGAGGTTGCAGAATCCGATTACTCCAACAACATCATGAAATGCAGG AGCCGCTATGACGGCCACCGCATCTGGATGTACAACTGCCACATAGGTGG TTCCTTCAGCGAAGAGACGGAAAAAAAGTTTGAGCACTTCAGCGGGCTCT TAAACAACCAGCTGTCCCCGCAGTAAAGAAGCCTGCGTGGTCAACTCCTG TCTTCAGGCCACACCACATCTTCCATGGGACTTCCCCCCAACAACTGAGTC TGAACGAATGCCACGTGCCCTCACCCAGCCCGGCCCCCACCCTGTCCAGA CCCCTACAGCTGTGTCTAAGCTCAGGAGGAAAGGGACCCTCCCATCATTC ATGGGGGGCTGCTACCTGACCCTTGGGGCCTGAGAAGGCCTTGGGGGGGT GGGGTTTGTCCACAGAGCTGCTGGAGCAGCACCAAGAGCCAGTCTTGACC GGGATGAGGCCCACAGACAGGTTGTCATCAGCTTGTCCCATTCAAGCCAC CGAGCTCACCACAGACACAGTGGAGCCGCGCTCTTCTCCAGTGACACGTG GACAAATGCGGGCTCATCAGCCCCCCAGAGAGGGTCAGGCCGAACCCCA TTTCTCCTCCTCTTAGGTCATTTTCAGCAAACTTGAATATCTAGACCTCTCT TCCAATGAAACCCTCCAGTCTATTATAGTCACATAGATAATGGTGCCACGT GTTTTCTGATTTGGTGAGCTCAGACTTGGTGCTTCCCTCTCCACAACCCCC ACCCCTTGTTTTCAAGATACTATTATTATATTTCACAGACTTTTGAAGCA CAAATTTATTGGCATTTAATATTGGACATCTGGGCCCTTGGAAGTACAAAT CTAAGGAAAAACCAACCCACTGTGTAAGTGACTCATCTTCCTGTTGTTCCA ATTCTGTGGGTTTTTGATTCAACGGTGCTATAACCAGGGTCCTGGGTGACA GGGCGCTCACTGAGCACCATGTGTCATCACAGACACTTACACATACTTGA AACTTGGAATAAAAGAAAGATTTATG SEQ ID NO: 98

### Thyroglobulin

>gi|33589851|ref[NM\_003235.3| Homo sapiens thyroglobulin (TG), mRNA | qPCR forward\_primer match [886..905] | qPCR reverse\_primer match [962..941] | qPCR probe match [915..939]

GCAGTGGTTTCTCCTCCTCCCAGGAAGGGCCAGGAAAATGGC CCTGGTCCTGGAGATCTTCACCCTGCTGGCCTCCATCTGCTGGGTGTCGGC CAATATCTTCGAGTACCAGGTTGATGCCCAGCCCCTTCGTCCCTGTGAGCT GCAGAGGGAAACGGCCTTTCTGAAGCAAGCAGACTACGTGCCCCAGTGTG CAGAGGATGGCAGCTTCCAGACTGTCCAGTGCCAGAACGACGGCCGCTCC TGCTGGTGTGGGTGCCAACGGCAGTGAAGTGCTGGGCAGCAGCAGCC AGGACGGCCTGTGGCTTGTCTGTCATTTTGTCAGCTACAGAAACAGCAGA GTCAGGATTCAGGGGACTACGCGCCTGTTCAGTGTGATGTGCAGCATGTC CAGTGCTGGTGTGGACGCAGAGGGGATGGAGGTGTATGGGACCCGCCA GCTGGGGAGGCCAAAGCGATGTCCAAGGAGCTGTGAAATAAGAAATCGT CGTCTTCTCCACGGGGTGGGAGATAAGTCACCACCCCAGTGTTCTGCGGA GGGAGAGTTTATGCCTGTCCAGTGCAAATTTGTCAACACCACAGACATGA TGATTTTTGATCTGGTCCACAGCTACAACAGGTTTCCAGATGCATTTGTGA CCTTCAGTTCCTTCCAGAGGAGGTTCCCTGAGGTATCTGGGTATTGCCACT GTGCTGACAGCCAAGGGCGGGAACTGGCTGAGACAGGTTTGGAGTTGTTA TTCACTGAAACCACCCTGTACCGGATACTGCAGAGACGGTTCCTCGCAGTT CAATCAGTCATCTCTGGCAGATTCCGATGCCCCACAAAATGTGAAGTGGA GCGGTTTACAGCAACCAGCTTTGGTCACCCCTATGTTCCAAGCTGCCGCCG AAATGGCGACTATCAGGCGGTGCAGTGCCAGACGGAAGGGCCCTGCTGGT GTGTGGACGCCCAGGGGAAGGAAATGCATGGAACCCGGCAGCAAGGGGA GCCGCCATCTTGTGCTGAAGGCCAATCTTGTGCCTCCGAAAGGCAGCAGG TGTTCTCTCCCAGAGAAAAGATGGGCCTCTCCAAGAGTAGCCAGATTT GCCACATCCTGCCCACCACGATCAAGGAGCTCTTTGTGGACTCTGGGCTT CTCCGCCCAATGGTGGAGGGACAGAGCCAACAGTTTTCTGTCTCAGAAAA TCTTGCCCTTCAGTTTACCACCAACCCAAAGAGACTCCAGCAAAACCTTTT TGGAGGGAAATTTTTGGTGAATGTTGGCCAGTTTAACTTGTCTGGAGCCCT TGGCACAAGAGGCACATTTAACTTCAGTCAATTTTTCCAGCAACTTGGTCT TGCAAGCTTCTTGAATGGAGGGAGACAAGAAGATTTGGCCAAGCCACTCT CTGTGGGATTAGATTCAAATTCTTCCACAGGAACCCCTGAAGCTGCTAAG AAGGATGGTACTATGAATAAGCCAACTGTGGGCAGCTTTGGCTTTGAAAT TAACCTACAAGAGAACCAAAATGCCCTCAAATTCCTTGCTTCTCTCTGGA GCTTCCAGAATTCCTTCTTCTTGCAACATGCTATCTCTGTGCCAGAAGA TGTGGCAAGAGATTTAGGTGATGTGATGGAAACGGTACTCGACTCCCAGA CCTGTGAGCAGACACCTGAAAGGCTATTTGTCCCATCATGCACGACAGAA GGAAGCTATGAGGATGTCCAATGCTTTTCCGGAGAGTGCTGGTGTGTGAA TTCCTGGGGCAAAGAGCTTCCAGGCTCAAGAGTCAGAGATGGACAGCCAA GGTGCCCACAGACTGTGAAAAGCAAAGGGCTCGCATGCAAAGCCTCATG GGCAGCCAGCCTGCTCCACCTTGTTTGTCCCTGCTTGTACTAGTGAG GGACATTTCCTGCCTGTCCAGTGCTTCAACTCAGAGTGCTACTGTTTGAT GCTGAGGGTCAGGCCATTCCTGGAACTCGAAGTGCAATAGGGAAGCCCAA

GAAATGCCCCACGCCTGTCAATTACAGTCTGAGCAAGCTTTCCTCAGGA CCTACATCCCACAGTGCAGCACCGATGGGCAGTGGAGACAAGTGCAATGC **AATGGGCCTCCTGAGCAGGTCTTCGAGTTGTACCAACGATGGGAGGCTCA** GAACAAGGGCCAGGATCTGACGCCTGCCAAGCTGCTAGTGAAGATCATGA GCTACAGAGAAGCAGCTTCCGGAAACTTCAGTCTCTTTATTCAAAGTCTGT ATGAGGCTGGCCAGCAAGATGTCTTCCCGGTGCTGTCACAATACCCTTCTC TGCAAGATGTCCCACTAGCAGCACTGGAAGGGAAACGGCCCCAGCCCAG GGAGAATATCCTCCTGGAGCCCTACCTCTTCTGGCAGATCTTAAATGGCCA ACTCAGCCAATACCCGGGGTCCTACTCAGACTTCAGCACTCCTTTGGCACA TTTTGATCTTCGGAACTGCTGGTGTGTGGATGAGGCTGGCCAAGAACTGG AAGGAATGCGGTCTGAGCCAAGCAAGCTCCCAACGTGTCCTGGCTCCTGT GAGGAAGCAAAGCTCCGTGTACTGCAGTTCATTAGGGAAACGGAAGAGA TTGTTTCAGCTTCCAACAGTTCTCGGTTCCCTCTGGGGGAGAGTTTCCTGG TGGCCAAGGGAATCCGGCTGAGGAATGAGGACCTCGGCCTTCCTCCGCTC TTCCCGCCCGGGAGGCTTTCGCGGAGTTTCTGCGTGGGAGTGATTACGCC ATTCGCCTGGCGCTCAGTCTACCTTAAGCTTCTATCAGAGACGCCGCTTT TCCCGGACGACTCGGCTGGAGCATCCGCCCTTCTGCGGTCGGGCCCCTAC ATGCCACAGTGTGATGCGTTTGGAAGTTGGGAGCCTGTGCAGTGCCACGC TGGGACTGGCACTGCTGGTGTGTAGATGAGAAAGGAGGGTTCATCCCTG GCTCACTGACTGCCGCTCTCTGCAGATTCCACAGTGCCCGACAACCTGCG AGAAATCTCGAACCAGTGGGCTGCTTTCCAGTTGGAAACAGGCTAGATCC AGGAGAATATGCCAGGCTGCAGGCATCGGGGGCTGGCACCTGGTGTGTGG ACCTGCATCAGGAGAAGAGTTGCGGCCTGGCTCGAGCAGCAGTGCCCAG TGCCCAAGCCTCTGCAATGTGCTCAAGAGTGGAGTCCTCTCTAGGAGAGT CAGCCAGGCTATGTCCCAGCCTGCAGGGCAGAGGATGGGGGCTTTTCCC CAGTGCAATGTGACCAGGCCCAGGGCAGCTGCTGGTGTCATGGACAGC GGAGAAGAGGTGCCTGGGACGCGTGACCGGGGGCCAGCCCGCCTGTG ACAATCCTGTGTGAGACAATCTCGGGCCCCACAGGCTCTGCCATGCAGCA GTGCCAATTGCTGTGCCGCCAAGGCTCCTGGAGCGTGTTTCCACCAGGGC CATTGATATGTAGCCTGGAGAGCGGACGCTGGGAGTCACAGCTGCCTCAG CCCGGGCCTGCCAACGCCCCAGCTGTGGCAGACCATCCAGACCCAAGG GCACTTTCAGCTCCAGCTCCCGCCGGGCAAGATGTGCAGTGCTGACTACG CGGGTTTGCTGCAGACTTTCCAGGTTTTCATATTGGATGAGCTGACAGCCC GCGGCTTCTGCCAGATCCAGGTGAAGACTTTTGGCACCCTGGTTTCCATTC CTGTCTGCAACACTCCTCTGTGCAGGTGGGTTGTCTGACCAGGGAGCGTT TAGGAGTGAATGTTACATGGAAATCACGGCTTGAGGACATCCCAGTGGCT TCTCTTCCTGACTTACATGACATTGAGAGAGCCTTGGTGGGCAAGGATCTC CTTGGGCGCTTCACAGATCTGATCCAGAGTGGCTCATTCCAGCTTCATCTG GACTCCAAGACGTTCCCAGCGGAAACCATCCGCTTCCTCCAAGGGGACCA CTTTGGCACCTCTCCTAGGACACGGTTTGGGTGCTCGGAAGGATTCTACCA AGTCTTGACAAGTGAGGCCAGTCAGGACGGACTGGGATGCGTTAAGTGCC ATGAAGGAAGCTATTCCCAAGATGAGGAATGCATTCCTTGTCCTGTTGGA TTCTACCAAGAACAGGCAGGGAGCTTGGCCTGTGTCCCATGTCCTGTGGG CAGAACGACCATTTCTGCCGGAGCTTTCAGCCAGACTCACTGTCACTGA CTGTCAGAGGAACGAAGCAGGCCTGCAATGTGACCAGAATGGCCAGTATC GAGCCAGCCAGAAGGACAGGGCCAGTGGGAAGGCCTTCTGTGTGGACGG CGAGGGCGGAGGCTGCCATGGTGGGAAACAGAGGCCCCTCTTGAGGAC

TCACAGTGTTTGATGATGCAGAAGTTTGAGAAGGTTCCAGAATCAAAGGT GATCTTCGACGCCAATGCTCCTGTGGCTGTCAGATCCAAAGTTCCTGATTC TGAGTTCCCCGTGATGCAGTGCTTGACAGATTGCACAGAGGACGAGGCCT GCAGCTTCTTCACCGTGTCCACGACGAGCCAGAGATTTCCTGTGATTTCT **ATGCTTGGACAAGTGACAATGTTGCCTGCATGACTTCTGACCAGAAACGA** GATGCACTGGGGAACTCAAAGGCCACCAGCTTTGGAAGTCTTCGCTGCCA GGTGAAAGTGAGGAGCCATGGTCAAGATTCTCCAGCTGTGTATTTGAAAA AGGGCCAAGGATCCACCACAACACTTCAGAAACGCTTTGAACCCACTGGT TTCCAAAACATGCTTTCTGGATTGTACAACCCCATTGTGTTCTCAGCCTCA GGAGCCAATCTAACCGATGCTCACCTCTTCTGTCTTCTTGCATGCGACCGT GATCTGTGTTGCGATGGCTTCGTCCTCACACAGGTTCAAGGAGGTGCCATC ATCTGTGGGTTGCTGAGCTCACCCAGTGTCCTGCTTTGTAATGTCAAAGAC TGGATGGATCCCTCTGAAGCCTGGGCTAATGCTACATGTCCTGGTGTGACA TATGACCAGGAGAGCCACCAGGTGATATTGCGTCTTGGAGACCAGGAGTT CATCAAGAGTCTGACACCCTTAGAAGGAACTCAAGACACCTTTACCAATT TTCAGCAGGTTTATCTCTGGAAAGATTCTGACATGGGGTCTCGGCCTGAGT CTATGGGATGTAGAAAAAACACAGTGCCAAGGCCAGCATCTCCAACAGA AGCAGGTTTGACAACAGAACTTTTCTCCCCTGTGGACCTCAACCAGGTCAT TGTCAATGGAAATCAATCACTATCCAGCCAGAAGCACTGGCTTTTCAAGC ACCTGTTTTCAGCCCAGCAGGCAAACCTATGGTGCCTTTCTCGTTGTGC AGGAGCACTCTTTCTGTCAGCTCGCAGAGATAACAGAGAGTGCATCCTTG TACTTCACCTGCACCCTCTACCCAGAGGCACAGGTGTGTGATGACATCATG GAGTCCAATACCCAGGGCTGCAGACTGATCCTGCCTCAGATGCCAAAGGC CCTGTTCCGGAAGAAGTTATACTGGAAGATAAAGTGAAGAACTTTTACA CTCGCCTGCCGTTCCAAAAACTGATGGGGATATCCATTAGAAATAAAGTG CCCATGTCTGAAAAATCTATTTCTAATGGGTTCTTTGAATGTGAACGACGG TGCGATGCGGACCCATGCTGCACTGGCTTTGGATTTCTAAATGTTTCCCAG TTAAAAGGAGAGAGGTGACATGTCTCACTCTGAACAGCTTGGGAATTCA GATGTGCAGTGAGGAGAATGGAGGAGCCTGGCGCATTTTGGACTGTGGCT CTCCTGACATTGAAGTCCACACCTATCCCTTCGGATGGTACCAGAAGCCCA TTGCTCAAAATAATGCTCCCAGTTTTTGCCCTTTGGTTGTTCTGCCTTCCCT CACAGAGAAAGTGTCTCTGGAATCGTGGCAGTCCCTGGCCCTCTCTCAGT GGTTGTTGATCCATCCATTAGGCACTTTGATGTTGCCCATGTCAGCACTGC TGCCACCAGCAATTTCTCTGCTGTCCGAGACCTCTGTTTGTCGGAATGTTC CCAACATGAGGCCTGTCTCATCACCACTCTGCAAACCCAACTCGGGGCTG TGAGATGTATGTTCTATGCTGATACTCAAAGCTGCACACATAGTCTGCAGG AAGCCAGGAATCTCTCTGCTCAGCTATGAGGCATCTGTACCTTCTGTGCCC ATTTCCACCATGCCGGCTGCTGGGCAGGTCCCAGGCCATCCAGGTGGG TACCTCATGGAAGCAAGTGGACCAGTTCCTTGGAGTTCCATATGCTGCCCC GCCCTGGCAGAGAGGCACTTCCAGGCACCAGAGCCCTTGAACTGGACAG GCTCCTGGGATGCCAGCAAGCCAAGGCCAGCTGCTGGCAGCCAGGCACC TTCATCCTCAGAATGTGGCCCCTAACGCGTCTGTGCTGGTGTTCTTCCAC AACACCATGGACAGGAGGAGAGTGAAGGATGGCCGGCTATCGACGGCT CCTTCTTGGCTGCTGTTGGCAACCTCATCGTGGTCACTGCCAGCTACCGAG TGGGTGTCTTCGGCTTCCTGAGTTCTGGATCCGGAGAGGTGAGTGGCAACT GGGGGCTGCTGGACCAGGTGGCGGCTCTGACCTGGGTGCAGACCCACATC CGAGGATTTGGCGGGGACCCTCGGCGCGTGTCCCTGGCAGCAGACCGTGG CGGGGCTGATGTGGCCAGCATCCACCTTCTCACGGCCAGGGCCACCAACT

CCCAACTTTCCGGAGAGCTGTGCTGATGGGAGGCTCCGCACTCTCCCCGG CCGCCGTCATCAGCCATGAGAGGGCTCAGCAGCAGCAATTGCTTTGGCA AAGGAGGTCAGTTGCCCCATGTCATCCAGCCAAGAAGTGGTGTCCTGCCT CCGCCAGAAGCCTGCCAATGTCCTCAATGATGCCCAGACCAAGCTCCTGG CCGTGAGTGGCCCTTTCCACTACTGGGGTCCTGTGATCGATGGCCACTTCC TCCGTGAGCCTCCAGCCAGAGCACTGAAGAGGTCTTTATGGGTAGAGGTC GATCTGCTCATTGGGAGTTCTCAGGACGACGGGCTCATCAACAGAGCAAA GGCTGTGAAGCAATTTGAGGAAAGTCGAGGCCGGACCAGTAGCAAAACA GCCTTTTACCAGGCACTGCAGAATTCTCTGGGTGGCGAGGACTCAGATGC CCGCGTCGAGGCTGCTACATGGTATTACTCTCTGGAGCACTCCACGGA TGACTATGCCTCCTTCTCCCGGGCTCTGGAGAATGCCACCCGGGACTACTT TATCATCTGCCCTATAATCGACATGGCCAGTGCCTGGGCAAAGAGGGCCC GAGGAAACGTCTTCATGTACCATGCTCCTGAAAACTACGGCCATGGCAGC CTGGAGCTGCTGGCGGATGTTCAGTTTGCCTTGGGGCTTCCCTTCTACCCA GCTACGAGGGCAGTTTTCTCTGGAGGAGAAGAGCCTGTCGCTGAAAAT CATGCAGTACTTTTCCCACTTCATCAGATCAGGAAATCCCAACTACCCTTA TGAGTTCTCACGGAAAGTACCCACATTTGCAACCCCCTGGCCTGACTTTGT ACCCGTGCTGGTGGAGAGAACTACAAGGAGTTCAGTGAGCTGCTCCCCA ATCGACAGGCCTGAAGAAGCCGACTGCTCCTTCTGGTCCAAGTACATC GAGTGAAGAGGAGTTGACGCTGGATCTGGGCTAAGAGAAGATCTC CTAAGCCTCCAGGAACCAGGCTCTAAGACCTACAGCAAGTGACCAGCCCT TGAGCTCCCCAAAAACCTCACCGAGGCTGCCCACTATGGTCATCTTTTTC TCTAAAATAGTTACTTACCTTCAATAAAGTATCTACATGCGGTG

**SEQ ID NO: 99** 

# **Transforming Growth Factor, Beta 1**

>gi|10863872|ref|NM\_000660.1| Homo sapiens transforming growth factor, beta 1 (Camurati-Engelmann disease) (TGFB1), mRNA | qPCR forward\_primer match [1651..1668] | qPCR reverse\_primer match [1539..1557] | qPCR probe match [1687..1713]

GCAGGGGGACGCCCGTCCGGGGCACCCCCCGGCTCTGAGCCGCCCG CGGGGCCGGCCCGGAGCGGAGGAGGAGTCGCCGAGGAGCAGC AGGAAAAAACTTTTGAGACTTTTCCGTTGCCGCTGGGAGCCGGAGGCGC GGGGACCTCTTGGCGCGACGCTGCCCCGCGAGGAGGCAGGACTTGGGGAC TACACGGCGTCCCTCAGGCGCCCCCATTCCGGACCAGCCCTCGGGAGTCG CCGACCGGCCTCCCGCAAAGACTTTTCCCCAGACCTCGGGCGCACCCCCT GCACGCCGCCTTCATCCCCGGCCTGTCTCCTGAGCCCCCGCGCATCCTAGA CCCTTTCTCCCAGGAGACGGATCTCTCTCCGACCTGCCACAGATCCCCT ATTCAAGACCACCACCTTCTGGTACCAGATCGCGCCCATCTAGGTTATTT CCGTGGGATACTGAGACACCCCGGTCCAAGCCTCCCCTCCACCACTGCG CCCTTCTCCCTGAGGAGCCTCAGCTTTCCCTCGAGGCCCTCCTACCTTTTGC CGGGAGACCCCCAGCCCTGCAGGGGGGGGCCTCCCCACCACCACCAGCC

CTGACGCCTGGCCGCCGCCGCGGGACTATCCACCTGCAAGACTATCGA CATGGAGCTGGTGAAGCGGAAGCGCATCGAGGCCATCCGCGGCCAGATCC TGTCCAAGCTGCGGCTCGCCAGCCCCGAGCCAGGGGGAGGTGCCGCCC GGCCGCTGCCGAGGCCGTGCTCGCCCTGTACAACAGCACCCGCGACCG GGTGGCCGGGGAGAGTGCAGAACCGGAGCCGAGCCTGAGGCCGACTAC TACGCCAAGGAGGTCACCCGCGTGCTAATGGTGGAAACCCACAACGAAAT CTATGACAAGTTCAAGCAGAGTACACACAGCATATATATGTTCTTCAACA CATCAGAGCTCCGAGAAGCGGTACCTGAACCCGTGTTGCTCTCCCGGGCA GAGCTGCGTCTGAGGAGGCTCAAGTTAAAAGTGGAGCACGTGGA GCTGTACCAGAAATACAGCAACAATTCCTGGCGATACCTCAGCAACCGGC TGCTGGCACCCAGCGACTCGCCAGAGTGGTTATCTTTTGATGTCACCGGAG TTGTGCGGCAGTGGTTGAGCCGTGGAGGGGAAATTGAGGGCTTTCGCCTT AGCGCCACTGCTCTGTGACAGCAGGGATAACACACTGCAAGTGGACAT CAACGGGTTCACTACCGGCCGCCGAGGTGACCTGGCCACCATTCATGGCA TGAACCGGCCTTTCCTGCTTCTCATGGCCACCCCGCTGGAGAGGGCCCAGC ATCTGCAAAGCTCCCGGCACCGCGAGCCCTGGACACCAACTATTGCTTC AGCTCCACGGAGAAGAACTGCTGCGTGCGGCAGCTGTACATTGACTTCCG CAAGGACCTCGGCTGGAAGTGGATCCACGAGCCCAAGGGCTACCATGCCA ACTTCTGCCTCGGGCCCTGCCCCTACATTTGGAGCCTGGACACGCAGTACA GCAAGGTCCTGGCCCTGTACAACCAGCATAACCCGGGCGCCTCGGCGGCG CCGTGCTGCCGCAGGCGCTGGAGCCGCTGCCCATCGTGTACTACGT GGGCCGCAAGCCCAAGGTGGAGCAGCTGTCCAACATGATCGTGCGCTCCT GCAAGTGCAGCTGAGGTCCCGCCCCGCCCCGCCCCGGCAGGCCCG GCCCCACCCGCCCCGCCCCGCTGCCTTGCCCATGGGGGCTGTATTTAAG GACACCGTGCCCCAAGCCCACCTGGGGCCCCATTAAAGATGGAGAGAGG ACTGCGGATCTCTGTGTCATTGGGCGCCTGCCTGGGGTCTCCATCCCTGAC TGCACTATTCCTTTGCCCGGCATCAAGGCACAGGGGACCAGTGGGGAACA CTACTGTAGTTAGATCTATTTATTGAGCACCTTGGGCACTGTTGAAGTGCC TTACATTAATGAACTCATTCAGTCACCATAGCAACACTCTGAGATGGCAG GGACTCTGATAACACCCATTTTAAAGGTTGAGGAAACAAGCCCAGAGAGG TTAAGGGAGGAGTTCCTGCCCACCAGGAACCTGCTTTAGTGGGGGATAGT GAAGAAGACAATAAAAGATAGTAGTTCAGGCCAGGCGGGGTGCTCACGC CTGTAATCCTAGCACTTTTGGGAGGCAGAGATGGGAGGATACTTGAATCC AGGCATTTGAGACCAGCCTGGGTAACATAGTGAGACCCTATCTCTACAAA ACACTTTTAAAAAATGTACACCTGTGGTCCCAGCTACTCTGGAGGCTAAG GTGGGAGGATCACTTGATCCTGGGAGGTCAAGGCTGCAG

**SEQ ID NO: 100** 

#### Serine Proteinase Inhibitor, Clade H, Member 1

>gi|32454740|ref|NM\_001235.2| Homo sapiens serine (or cysteine) proteinase inhibitor, clade H (heat shock protein 47), member 1, (collagen binding protein 1) (SERPINH1), mRNA | qPCR assay on demand context match [184..208]

TCTTTGGCTTTTTTTGGCGGAGCTGGGGCCCCTCCGGAAGCGTTTCCAACTTTCCAGAAGTTTCTCGGGACGGGCAGGAGGGGGTGGGGACTGCCA

TATATAGATCCCGGGAGCAGGGGAGCGGGCTAAGAGTAGAATCGTGTCGC GGCTCGAGAGCGAGAGTCACGTCCCGGCGCTAGCCCAGCCCGACCCAGGC TCAGCGCCTTCTGCCTCCTGGAGGCGCCCTGGCCGCCGAGGTGAAGAAA CCTGCAGCCGCAGCAGCTCCTGGCACTGCGGAGAAGTTGAGCCCCAAGGC GGCCACGCTTGCCGAGCGCAGCGCCGGCCTGGCCTTCAGCTTGTACCAGG CCATGCCAAGGACCAGGCAGTGGAGAACATCCTGGTGTCACCCGTGGTG GTGGCCTCGTCGCTAGGGCTCGTGTCGCTGGCCGCCAAGGCGACCACGGC GTCGCAGGCCAAGGCAGTGCTGAGCGCCGAGCAGCTGCGCGACGAGGAG GTGCACGCCGGCCTGGGCGAGCTGCTGCGCTCACTCAGCAACTCCACGGC GCGCAACGTGACCTGGAAGCTGGGCAGCCGACTGTACGGACCCAGCTCAG TGAGCTTCGCTGATGACTTCGTGCGCAGCAGCAGCAGCACTACAACTGC GAGCACTCCAAGATCAACTTCCGCGACAAGCGCAGCGCGCTGCAGTCCAT CAACGAGTGGCCGCGCAGACCACCGACGCAAGCTGCCCGAGGTCACC AAGGACGTGGAGCGCACGGACGCCCTGCTAGTCAACGCCATGTTCTT CAAGCCACACTGGGATGAGAAATTCCACCACAAGATGGTGGACAACCGTG GCTTCATGGTGACTCGGTCCTATACCGTGGGTGTCATGATGATGCACCGGA CAGGCCTCTACAACTACTACGACGACGAGAAGGAAAAGCTGCAAATCGTG GAGATGCCCCTGGCCCACAAGCTCTCCAGCCTCATCATCCTCATGCCCCAT CACGTGGAGCCTCTCGAGCGCCTTGAAAAGCTGCTAACCAAAGAGCAGCT GAAGATCTGGATGGGGAAGATGCAGAAGAAGGCTGTTGCCATCTCCTTGC CTGGGCCTGACTGAGGCCATTGACAAGAACAAGGCCGACTTGTCACGCAT GTCAGGCAAGAAGGACCTGTACCTGGCCAGCGTGTTCCACGCCACCGCCT TTGAGTTGGACACAGATGGCAACCCCTTTGACCAGGACATCTACGGGCGC GAGGAGCTGCGCAGCCCCAAGCTGTTCTACGCCGACCACCCCTTCATCTTC CTAGTGCGGGACACCCAAAGCGGCTCCCTGCTATTCATTGGGCGCCTGGT CCGGCCTAAGGGTGACAAGATGCGAGACGAGTTATAGGGCCTCAGGGTGC ACACAGGATGCAGGAGCATCCAAAGGCTCCTGAGACACATGGGTGCT ATTGGGGTTGGGGGGGGGGTGAGGTACCAGCCTTGGATACTCCATGGGGT GGGGGTGGAAAAACAGACCGGGGTTCCCGTGTGCCTGAGCGGACCTTCCC AGCTAGAATTCACTCCACTTGGACATGGGCCCCAGATACCATGATGCTGA CCTGAAAGTCCCAGATCAAGCCTGCCTCAATCAGTATTCATATTTATAGCC CCTCTTCTGACACTAAAACACCTCAGCTGCCTCCCCAGCTCTATCCCAACC TCTCCCAACTATAAAACTAGGTGCTGCAGCCCCTGGGACCAGGCACCCCC AGAATGACCTGGCCGCAGTGAGGCGGATTGAGAAGGAGCTCCCAGGAGG GGCTTCTGGGCAGACTCTGGTCAAGAAGCATCGTGTCTGGCGTTGTGGGG ATGAACTTTTTGTTTCTTCCTTTTTTAGTTCTTCAAAGATAGGGAGG GAAGGGGAACATGAGCCTTTGTTGCTATCAATCCAAGAACTTATTTGTA CATTTTTTTTCAATAAACTTTTCCAATGACATTTTGTTGGAGCGTGGAA AAAA **SEQ ID NO: 101** 

#### Serine Proteinase Inhibitor, Clade B, Member 5

>gi|4505788|ref|NM\_002639.1| Homo sapiens serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 5 (SERPINB5), mRNA | qPCR

forward\_primer match [36..56] | qPCR reverse\_primer match [106..86] | qPCR probe match [60..80]

GGCACGAGTTGTCCTCCTCGCTTGCCTGTTCCTTTTCCACGCATTTT CCAGGATAACTGTGACTCCAGGCCCGCAATGGATGCCCTGCAACTAGCAA ATTCGGCTTTTGCCGTTGATCTGTTCAAACAACTATGTGAAAAGGAGCCAC TGGGCAATGTCCTCTCTCCAATCTGTCTCCACCTCTGTCACTTGC TCAAGTGGGTGCTAAAGGTGACACTGCAAATGAAATTGGACAGGTTCTTC ATTTTGAAAATGTCAAAGATATACCCTTTGGATTTCAAACAGTAACATCGG ATGTAAACAAACTTAGTTCCTTTTACTCACTGAAACTAATCAAGCGGCTCT ACGTAGACAAATCTCTGAATCTTTCTACAGAGTTCATCAGCTCTACGAAGA GACCCTATGCAAAGGAATTGGAAACTGTTGACTTCAAAGATAAATTGGAA GAAACGAAAGGTCAGATCAACAACTCAATTAAGGATCTCACAGATGGCCA CTTTGAGAACATTTTAGCTGACAACAGTGTGAACGACCAGACCAAAATCC TTGTGGTTAATGCTGCCTACTTTGTTGGCAAGTGGATGAAGAAATTTCCTG ACCAGTGCAGATGAACATGGAGGCCACGTTCTGTATGGGAAACATTG ACAGTATCAATTGTAAGATCATAGAGCTTCCTTTTCAAAATAAGCATCTCA GCATGTTCATCCTACTACCCAAGGATGTGGAGGATGAGTCCACAGGCTTG GAGAAGATTGAAAAACAACTCAACTCAGAGTCACTGTCACAGTGGACTAA TCCCAGCACCATGGCCAATGCCAAGGTCAAACTCTCCATTCCAAAATTTA AGGTGGAAAAGATGATTGATCCCAAGGCTTGTCTGGAAAATCTAGGGCTG AAACATATCTTCAGTGAAGACACATCTGATTTCTCTGGAATGTCAGAGAC CAAGGGAGTGGCCCTATCAAATGTTATCCACAAAGTGTGCTTAGAAATAA CTGAAGATGGTGGGGATTCCATAGAGGTGCCAGGAGCACGGATCCTGCAG CACAAGGATGAATTGAATGCTGACCATCCCTTTATTTACATCATCAGGCAC AACAAAACTCGAAACATCATTTCTTTGGCAAATTCTGTTCTCCTTAAGTG GCATAGCCCATGTTAAGTCCTCCCTGACTTTTCTGTGGATGCCGATTTCTG TAAACTCTGCATCCAGAGATTCATTTTCTAGATACAATAAATTGCTAATGT TGCTGGATCAGGAAGCCGCCAGTACTTGTCATATGTAGCCTTCACACAGA TAGACCTTTTTTTTCCAATTCTATCTTTTGTTTCCTTTTTTCCCATAAGA CAATGACATACGCTTTTAATGAAAAGGAATCACGTTAGAGGAAAAATATT TATTCATTATTTGTCAAATTGTCCGGGGTAGTTGGCAGAAATACAGTCTTC CACAAAGAAATTCCTATAAGGAAGATTTGGAAGCTCTTCTTCCCAGCAC TATGCTTTCCTTTTGGGATAGAGAATGTTCCAGACATTCTCGCTTCCCTG AAAGACTGAAGAAAGTGTAGTGCATGGGACCCACGAAACTGCCCTGGCTC CAGTGAAACTTGGGCACATGCTCAGGCTACTATAGGTCCAGAAGTCCTTA TGTTAAGCCCTGGCAGGCAGGTGTTTATTAAAATTCTGAATTTTGGGGATT TTCAAAAGATAATTTTACATACACTGTATGTTATAGAACTTCATGGATC AGATCTGGGGCAGCAACCTATAAATCAACACCTTAATATGCTGCAACAA ATGTAGAATATTCAGACAAAATGGATACATAAAGACTAAGTAGCCCATAA GGGGTCAAAATTTGCTGCCAAATGCGTATGCCACCAACTTACAAAAACAC TTCGTTCGCAGAGCTTTTCAGATTGTGGAATGTTGGATAAGGAATTATAGA CCTCTAGTAGCTGAAATGCAAGACCCCAAGAGGAAGTTCAGATCTTAATA TAAATTCACTTTCATTTTTGATAGCTGTCCCATCTGGTCATGTGGTTGGCAC TAGACTGGTGGCAGGGGCTTCTAGCTGACTCGCACAGGGATTCTCACAAT AGCCGATATCAGAATTTGTGTTGAAGGAACTTGTCTCTTCATCTAATATGA TAGCGGGAAAAGGAGAGAAACTACTGCCTTTAGAAAATATAAGTAAAG TGATTAAAGTGCTCACGTTACCTTGACACATAGTTTTTCAGTCTATGGGTT TAGTTACTTTAGATGCAAGCATGTAACTTATATTAATAGTAATTTGTAAA

GTTGGGTGGATAAGCTATCCCTGTTGCCGGTTCATGGATTACTTCTCTATA AAAAATATATTTACCAAAAAATTTTGTGACATTCCTTCTCCCATCTCTT CCTTGACATGCATTGTAAATAGGTTCTTCTTGTTCTGAGATTCAATATTGA ATTTCTCCTATGCTATTGACAATAAAATATTATTGAACTACC

**SEQ ID NO: 102** 

#### Carcinoembryonic Antigen-Related Cell Adhesion Molecule 5

>gi|11386170|ref|NM\_004363.1| Homo sapiens carcinoembryonic antigenrelated cell adhesion molecule 5 (CEACAM5), mRNA | qPCR assay on demand context match [2128..2152]

CTCAGGGCAGAGGAGGAAGGACAGACCAGACAGTCACAGC AGCCTTGACAAAACGTTCCTGGAACTCAAGCTCTTCTCCACAGAGGAGGA CAGAGCAGACAGAGACCATGGAGTCTCCCTCGGCCCCTCCCCACAGA TGGTGCATCCCTGGCAGAGGCTCCTGCTCACAGCCTCACTTCTAACCTTC TGGAACCCGCCCACCGCCAAGCTCACTATTGAATCCACGCCGTTCAAT GTCGCAGAGGGGAAGGAGGTGCTTCTACTTGTCCACAATCTGCCCCAGCA TCTTTTTGGCTACAGCTGGTACAAAGGTGAAAGAGTGGATGGCAACCGTC AAATTATAGGATATGTAATAGGAACTCAACAAGCTACCCCAGGGCCCGCA TACAGTGGTCGAGAGATAATATACCCCAATGCATCCCTGCTGATCCAGAA CATCATCCAGAATGACACAGGATTCTACACCCTACACGTCATAAAGTCAG ATCTTGTGAATGAAGAAGCAACTGGCCAGTTCCGGGTATACCCGGAGCTG CCCAAGCCTCCATCTCCAGCAACACTCCAAACCCGTGGAGGACAAGGA GTGGGTAAACAATCAGAGCCTCCCGGTCAGTCCCAGGCTGCAGCTGTCCA ATGGCAACAGGACCCTCACTCTATTCAATGTCACAAGAAATGACACAGCA AGCTACAAATGTGAAACCCAGAACCCAGTGAGTGCCAGGCGCAGTGATTC AGTCATCCTGAATGTCCTCTATGGCCCGGATGCCCCCACCATTTCCCCTCT AAACACATCTTACAGATCAGGGGAAAATCTGAACCTCTCCTGCCACGCAG CCTCTAACCCACCTGCACAGTACTCTTGGTTTGTCAATGGGACTTTCCAGC AATCCACCAAGAGCTCTTTATCCCCAACATCACTGTGAATAATAGTGGAT CCTATACGTGCCAAGCCCATAACTCAGACACTGGCCTCAATAGGACCACA GTCACGACGATCACAGTCTATGCAGAGCCACCCAAACCCTTCATCACCAG CAACACTCCAACCCGTGGAGGATGAGGATGCTGTAGCCTTAACCTGTG AACCTGAGATTCAGAACACAACCTACCTGTGGTGGGTAAATAATCAGAGC CTCCCGGTCAGTCCCAGGCTGCAGCTGTCCAATGACAACAGGACCCTCAC TCTACTCAGTGTCACAAGGAATGATGTAGGACCCTATGAGTGTGGAATCC AGAACGAATTAAGTGTTGACCACAGCGACCCAGTCATCCTGAATGTCCTC TATGGCCCAGACGACCCCACCATTTCCCCCTCATACACCTATTACCGTCCA GGGGTGAACCTCAGCCTCTCCTGCCATGCAGCCTCTAACCCACCTGCACA TTATCTCCAACATCACTGAGAAGAACAGCGGACTCTATACCTGCCAGGCC AATAACTCAGCCAGTGGCCACAGCAGGACTACAGTCAAGACAATCACAGT CTCTGCGGAGCTGCCCAAGCCCTCCATCTCCAGCAACACTCCAAACCCG TGGAGGACAAGGATGCTGTGGCCTTCACCTGTGAACCTGAGGCTCAGAAC GCTGCAGCTGTCCAATGGCAACAGGACCCTCACTCTATTCAATGTCACAA 

AACCGCAGTGACCCAGTCACCCTGGATGTCCTCTATGGGCCGGACACCCC CATCATTTCCCCCCAGACTCGTCTTACCTTTCGGGAGCGAACCTCAACCT CTCCTGCCACTCGGCCTCTAACCCATCCCGCAGTATTCTTGGCGTATCAA TGGGATACCGCAGCACACACACAGTTCTCTTTATCGCCAAAATCACGC CAAATAATAACGGGACCTATGCCTGTTTTGTCTCTAACTTGGCTACTGGCC GCAATAATTCCATAGTCAAGAGCATCACAGTCTCTGCATCTGGAACTTCTC CTGGTCTCTCAGCTGGGGCCACTGTCGGCATCATGATTGGAGTGCTGGTTG GGGTTGCTCTGATATAGCAGCCCTGGTGTAGTTTCTTCATTTCAGGAAGAC TGACAGTTGTTTTGCTTCTTCCTTAAAGCATTTGCAACAGCTACAGTCTAA AATTGCTTCTTTACCAAGGATATTTACAGAAAAGACTCTGACCAGAGATC GAGACCATCCTAGCCAACATCGTGAAACCCCATCTCTACTAAAAATACAA AAATGAGCTGGGCTTGGTGGCGCGCACCTGTAGTCCCAGTTACTCGGGAG GCTGAGGCAGGAGAATCGCTTGAACCCGGGAGGTGGAGATTGCAGTGAG CCCAGATCGCACCACTGCACTCCAGTCTGGCAACAGAGCAAGACTCCATC TCAAAAAGAAAAGAAAGAAGACTCTGACCTGTACTCTTGAATACAAGTT TTCATGGGACTAAATGAACTAATGAGGATTGCTGATTCTTTAAATGTCTTG TTTCCCAGATTTCAGGAAACTTTTTTTTTTTTAAGCTATCCACTCTTACAGC AATTTGATAAAATATACTTTTGTGAACAAAAATTGAGACATTTACATTTTC TCCTATGTGGTCGCTCCAGACTTGGGAAACTATTCATGAATATTTATATT GTATGGTAATATAGTTATTGCACAAGTTCAATAAAAATCTGCTCTTTGTAT AACAGAAAAA **SEQ ID NO: 103** 

### **Matrix Metalloproteinase 2**

>gi|11342665|ref|NM\_004530.1| Homo sapiens matrix metalloproteinase 2 (gelatinase A, 72kDa gelatinase, 72kDa type IV collagenase) (MMP2), mRNA | qPCR forward\_primer match [1713..1732] | qPCR reverse\_primer match [1793..1775] | qPCR probe match [1751..1773]

TGTTTCCGCTGCATCCAGACTTCCTCAGGCGGTGGCTGGAGGCTGC GCATCTGGGGCTTTAAACATACAAAGGGATTGCCAGGACCTGCGGCGGCG GCGGCGGCGGGGGCTGGGGCGCGGGGCCGGACCATGAGCCGCTGA GCCGGCAAACCCCAGGCCACCGAGCCAGCGACCCTCGGAGCGCAGCC CTGCGCCGCGACCAGGCTCCAACCAGGCGGCGAGGCGGCCACACGCAC CGAGCCAGCGACCCCGGGCGACGCGGGGCCAGGGAGCGCTACGATG GAGGCGCTAATGGCCCGGGGCGCGCTCACGGGTCCCCTGAGGGCGCTCTG TCTCCTGGGCTGCCTGAGCCACGCCGCCGCGCGCGCGTCGCCATCAT CAAGTTCCCCGGCGATGTCGCCCCCAAAACGGACAAAGAGTTGGCAGTGC AATACCTGAACACCTTCTATGGCTGCCCCAAGGAGAGCTGCAACCTGTTT GTGCTGAAGGACACACTAAAGAAGATGCAGAAGTTCTTTGGACTGCCCCA GACAGGTGATCTTGACCAGAATACCATCGAGACCATGCGGAAGCCACGCT GCGGCAACCCAGATGTGGCCAACTACAACTTCTTCCCTCGCAAGCCCAAG TGGGACAAGAACCAGATCACATACAGGATCATTGGCTACACACCTGATCT GGACCCAGAGACAGTGGATGATGCCTTTGCTCGTGCCTTCCAAGTCTGGA GCGATGTGACCCCACTGCGGTTTTCTCGAATCCATGATGGAGAGGCAGAC ATCATGATCAACTTTGGCCGCTGGGAGCATGGCGATGGATACCCCTTTGA CGGTAAGGACGGACTCCTGGCTCATGCCTTCGCCCCAGGCACTGGTGTTG

GGGGAGACTCCCATTTTGATGACGATGAGCTATGGACCTTGGGAGAAGGC CAAGTGGTCCGTGTGAAGTATGGCAACGCCGATGGGGAGTACTGCAAGTT CCCCTTCTTGTTCAATGGCAAGGAGTACAACAGCTGCACTGATACTGGCC GCAGCGATGGCTTCCTCTGGTGCTCCACCACCTACAACTTTGAGAAGGAT GGCAAGTACGGCTTCTGTCCCCATGAAGCCCTGTTCACCATGGGCGGCAA CGCTGAAGGACAGCCTGCAAGTTTCCATTCCGCTTCCAGGGCACATCCTA TGACAGCTGCACCACTGAGGGCCGCACGGATGGCTACCGCTGGTGCGGCA CCACTGAGGACTACGACCGCGACAAGAAGTATGGCTTCTGCCCTGAGACC GCCATGTCCACTGTTGGTGGGAACTCAGAAGGTGCCCCCTGTGTCTTCCCC TGACGGAAAGATGTGGTGTGCGACCACAGCCAACTACGATGACGACCGCA AGTGGGGCTTCTGCCCTGACCAAGGGTACAGCCTGTTCCTCGTGGCAGCC CACGAGTTTGGCCACGCCATGGGGCTGGAGCACTCCCAAGACCCTGGGGC CCTGATGGCACCCATTTACACCTACACCAAGAACTTCCGTCTGTCCCAGGA TGACATCAAGGGCATTCAGGAGCTCTATGGGGCCTCTCCTGACATTGACCT TGGCACCGGCCCCACCCCCACACTGGGCCCTGTCACTCCTGAGATCTGCA AACAGGACATTGTATTTGATGGCATCGCTCAGATCCGTGGTGAGATCTTCT TCTTCAAGGACCGGTTCATTTGGCGGACTGTGACGCCACGTGACAAGCCC ATGGGGCCCCTGCTGGTGGCCACATTCTGGCCTGAGCTCCCGGAAAAGAT TGATGCGGTATACGAGGCCCCACAGGAGGAGAAGGCTGTGTTCTTTGCAG GGAATGAATACTGGATCTACTCAGCCAGCACCCTGGAGCGAGGGTACCCC AAGCCACTGACCAGCCTGGGACTGCCCCCTGATGTCCAGCGAGTGGATGC CGCCTTTAACTGGAGCAAAAACAAGAAGACATACATCTTTGCTGGAGACA AATTCTGGAGATACAATGAGGTGAAGAAGAAAATGGATCCTGGCTTTCCC AAGCTCATCGCAGATGCCTGGAATGCCATCCCCGATAACCTGGATGCCGT CGTGGACCTGCAGGGCGGCGGTCACAGCTACTTCTTCAAGGGTGCCTATT ACCTGAAGCTGGAGAACCAAAGTCTGAAGAGCGTGAAGTTTGGAAGCATC AAATCCGACTGGCTAGGCTGCTGAGCTGGCCCTGGCTCCCACAGGCCCTT CCTCTCCACTGCCTTCGATACACCGGGCCTGGAGAACTAGAGAAGGACCC GGAGGGCCTGCCGTGCCTTCAGCTCTACAGCTAATCAGCATTCTC ACTCCTACCTGGTAATTTAAGATTCCAGAGAGTGGCTCCTCCCGGTGCCCA AGAATAGATGCTGACTGTACTCCTCCCAGGCGCCCCTTCCCCCTCCAATCC CACCAACCTCAGAGCCACCCTAAAGAGATCCTTTGATATTTTCAACGCA GCCTGCTTTGGGCTGCCTGGTGCCACACTTCAGGCTCTTCTCCTTTC ACAACCTTCTGTGGCTCACAGAACCCTTGGAGCCAATGGAGACTGTCTCA AGAGGGCACTGGTGGCCCGACAGCCTGGCACAGGGCAGTGGGACAGGGC ATGCCAGGTGGCCACTCCAGACCCCTGGCTTTTCACTGCTGCCTTA GGGTCTTGTTTTTTTTCCACTTAGAAATTGCATTTCCTGACAGAAGGACT CAGGTTGTCTGAAGTCACTGCACAGTGCATCTCAGCCCACATAGTGATGG TTCCCCTGTTCACTCACTTAGCATGTCCCTACCGAGTCTCTTCTCCACTGG TCAACCATTCCCCATGGGAAATGTCAACAAGTATGAATAAAGACACCTAC **TGAGTGGC SEQ ID NO: 104** 

# Proprotein Convertase Subtilisin/Kezin Type 5

>gi|20336245|ref]NM\_006200.2| Homo sapiens proprotein convertase subtilisin/kexin type 5 (PCSK5), mRNA | qPCR forward\_primer match [2677..2697] | qPCR reverse\_primer match [2821..2801] | qPCR probe match [2737..2765]

CGGAGGGAGCGTGGGAGCGAGCAAGCGAGCGTTTGGAGCCCGGG CCAGCAGAGGGGCGCCGGTCGCTGCCTGTACCGCTCCCGCTGGTCATC TCCGCCGCGCTCGGGGGCCCCGGGAGGGCGAGACCGAGTCGGAGAGTC CAGCCTCCTCCTGCGTCCGAGCCGGGGAGCATCGCCGAGCGCCCCACGGG CCGGAGAGCTGGGAGCACAGGTCCCGGCAGCCCCAGGGATGGTCTAGGA GCCGGCGTAAGGCTCGCTGCTCCTGCCGGGGCTAGCCGCCTCCTG CCGATCGCCCGGGGCTGCGAGCTGCGGGGCCCGGGGCTGCTCGCCGGGC GGCGCAGGCCGGAGAAGTTAGTTGTGCGCGCCCTTAGTGCGCGGAACCAG CCAGCGAGCGAGCAGCGAGCGCGGGACCATGGGCTGGGGGAGC CGCTGCTGCCCGGGACGTTTGGACCTGCTGTGCGTGCTGCTGCTC GGCAGTCAAAATCGCCGGGGGCTTCCCGGAGGCCAACCGTATCGCCAGCA AGTACGGATTCATCAACATAGGACAGATAGGGGCCCTGAAGGACTACTAC CACTTCTACCATAGCAGGACGATTAAAAGGTCAGTTATCTCGAGCAGAGG GACCCACAGTTTCATTTCAATGGAACCAAAGGTGGAATGGATCCAACAGC AAGTGGTAAAAAAGCGGACAAAGAGGGATTATGACTTCAGTCGTGCCCA GTCTACCTATTTCAATGATCCCAAGTGGCCCAGCATGTGGTATATGCACTG CAGTGACAATACACATCCCTGCCAGTCTGACATGAATATCGAAGGAGCCT GGAAGAGGCTACACGGGAAAGAACATTGTGGTCACTATCCTGGATGAC GGAATTGAGAGAACCCATCCAGATCTGATGCAAAACTACGATGCTCTGGC AAGTTGCGACGTGAATGGGAATGACTTGGACCCAATGCCTCGTTATGATG CAAGCAACGAGAACAAGCATGGGACTCGCTGTGCTGGAGAAGTGGCAGC CGCTGCAAACAATTCGCACTGCACAGTCGGAATTGCTTTCAACGCCAAGA TCGGAGGAGTGCGAATGCTGGACGGAGATGTCACGGACATGGTTGAAGC AAAATCAGTTAGCTTCAACCCCCAGCACGTGCACATTTACAGCGCCAGCT GGGGCCCGGATGATGGCAAGACTGTGGACGGACCAGCCCCCTCACC CGGCAAGCCTTTGAAAACGGCGTTAGAATGGGGCGGAGAGGCCTCGGCTC TGTGTTTGTTTGGGCATCTGGAAATGGTGGAAGGAGCAAAGACCACTGCT CCTGTGATGGCTACACCAACAGCATCTACACCATCTCCATCAGCAGCACT GCAGAAAGCGGAAAGAAACCTTGGTACCTGGAAGAGTGTTCATCCACGCT GGCCACAACCTACAGCAGCGGGGGGTCCTACGATAAGAAAATCATCACTA CAGATCTGAGGCAGCGTTGCACGGACAACCACACTGGGACGTCAGCCTCA GCCCCATGGCTGCAGGCATCATTGCGCTGGCCCTGGAAGCCAATCCGTTT CTGACCTGGAGAGACGTACAGCATGTTATTGTCAGGACTTCCCGTGCGGG ACATTTGAACGCTAATGACTGGAAAACCAATGCTGCTGGTTTTAAGGTGA GCCATCTTATGGATTTGGACTGATGGACGCAGAAGCCATGGTGATGGAG AGACCGACAAATCAAGACAATCCGCCCTAACAGTGCAGTGCGCTCCATCT ACAAAGCTTCAGGCTGCTCGGATAACCCCAACCGCCATGTCAACTACCTG GGCCATCTACCTGACCTCGCCCTCTGGAACTAGGTCTCAGCTTTTGGCCAA CAGGCTATTTGATCACTCCATGGAAGGATTCAAAAACTGGGAGTTCATGA CCATTCATTGCTGGGGAGAAAGAGCTGCTGGTGACTGGGTCCTTGAAGTT

TATGATACTCCCTCTCAGCTAAGGAACTTTAAGACTCCAGGTAAATTGAA AGAATGGTCTTTGGTCCTCTACGGCACCTCCGTGCAGCCATATTCACCAAC CAATGAATTTCCGAAAGTGGAACGGTTCCGCTATAGCCGAGTTGAAGACC CCACAGACGACTATGGCACAGAGGATTATGCAGGTCCCTGCGACCCTGAG TGCAGTGAGGTTGGCTGTGACGGGCCAGGACCAGACCACTGCAATGACTG TTTGCACTACTACAAGCTGAAAAACAATACCAGGATCTGTGTCTCCA GCTGCCCCCTGGCCACTACCACGCCGACAAGAAGCGCTGCAGGAAGTGT GCCCCAACTGTGAGTCCTGCTTTGGGAGCCATGGTGACCAATGCATGTCC TGCAAATATGGATACTTTCTGAATGAAGAAACCAACAGCTGTGTTACTCA CTGCCCTGATGGGTCATATCAGGATACCAAGAAAAATCTTTGCCGGAAAT GCAGTGAAAACTGCAAGACATGTACTGAATTCCATAACTGTACAGAATGT AGGGATGGGTTAAGCCTGCAGGGATCCCGGTGCTCTGTCTCCTGTGAAGA TGGACGGTATTTCAACGGCCAGGACTGCCAGCCCTGCCACCGCTTCTGCG CCACTTGTGCTGGGGCAGGAGCTGATGGGTGCATTAACTGCACAGAGGGC TACTTCATGGAGGATGGGAGATGCGTGCAGAGCTGTAGTATCAGCTATTA CTTTGACCACTCTTCAGAGAATGGATACAAATCCTGCAAAAAATGTGATA TCAGTTGTTTGACGTGCAATGGCCCAGGATTCAAGAACTGTACAAGCTGC CCTAGTGGGTATCTCTTAGACTTAGGAATGTGTCAAATGGGAGCCATTTGC AAGGATGCAACGGAAGAGTCCTGGGCGGAAGGAGGCTTCTGTATGCTTGT GAAAAGAACAATCTGTGCCAACGGAAGGTTCTTCAACAACTTTGCTGCA AAACATGTACATTCAAGGCTGAGCAGCCATCTTAGATTTCTTTGTTCCTG AG **SEQ ID NO: 105** 

# Carboxypeptidase N, polypeptide 2, 83kD

>gi|18554966|ref|XM\_087358.1| Homo sapiens carboxypeptidase N, polypeptide 2, 83kD (CPN2), mRNA

ATGGGTTGTGACTGCTTCGTCCAGGAGGTGTTCTGCTCAGATGAGG AGCTTGCCACCGTCCCGCTGGACATCCCGCCATATACGAAAAACATCATC TTTGTGGAGACCTCGTTCACCACATTGGAAACCAGAGCTTTTGGCAGTAAC CCCAACTTGACCAAGGTGGTCTTCCTCAACACTCAGCTCTGCCAGTTTAGG CCGGATGCCTTTGGGGGGCTGCCCAGGCTGGAGGACCTGGAGGTCACAGG CAGTAGCTTCTTGAACCTCAGCACCAACATCTTCTCCAACCTGACCTCGCT GGGCAAGCTCACCTCAACTTCAACATGCTGGAGGCTCTGCCCGAGGGTC TTTTCCAGCACCTGGCTGCCCTGGAGTCCCTCCACCTGCAGGGGAACCAGC TCCAGGCCCTGCCCAGGAGGCTCTTCCAGCCTCTGACCCATCTGAAGACA CTCAACCTGGCCCAGAACCTCCTGGCCCAGCTCCCGGAGGAGCTGTTCCA CCCACTCACCAGCCTGCAGACCCTGAAGCTGAGCAACAACGCGCTCTCTG GTCTCCCCAGGGTGTGTTTGGCAAACTGGGCAGCCTGCAGGAGCTCTTCC TGGACAGCAACACATCTCGGAGCTGCCCCCTCAGGTGTTCTCCCAGCTCT TCTGCCTAGAGAGGCTGTGGCTGCAACGCAACGCCATCACGCACCTGCCG CTCTCCATCTTTGCCTCCTGGGTAATCTGACCTTTCTGAGCTTGCAGTGG AACATGCTTCGGGTCCTGCCGGCCTCTTTGCCCACACCCCATGCCTG GTTGGCCTGTCTCTGACCCATAACCAGCTGGAGACTGTCGCTGAGGGCAC CTTTGCCCACCTGTCCAACCTGCGTTCCCTCATGCTCTCATACAATGCCATT ACCCACCTCCAGCTGGCATCTTCAGAGACCTGGAGGAGTTGGTCAAACT CTACCTGGGCAGCACACCTTACGGCGCTGCACCCAGCCCTCTTCCAGA ACCTGTCCAAGCTGGAGCTGCTCAGCCTCCCAAGAACCAGCTGACCACA CTTCCGGAGGCATCTTCGACACCAACTACAACCTGTTCAACCTGGCCCTG

CACGGTAACCCCTGGCAGTGCGACTGCCACCTGGCCTACCTCTTCAACTGG
CTGCAGCAGTACACCGATCGGCTCCTGAACATCCAGACCTACTGCGCTGG
CCCTGCCTACCTCAAAGGCCAGGTGGTGCCCGCCTTGAATGAGAAGCAGC
TGGTGTGTCCCGTCACCCGGGACCACTTGGGCTTCCAGGTCACGTGGCCG
GACGAAAGCAAGGCAGGGGCAGCTGGGATCTGGCTGTGCAGGAAAGGG
CAGCCCGGAGCCAGTGCACCTACAGCAACCCCGAGGGCACCGTGGTGCTC
GCCTGTGACCAGGCCCAGTGTCGCTGGCTGAACGTCCAGCTCTCTCCTTGG
CAGGGCTCCCTGGGACTGCAGTACAATGCTAGTCAGGAGTGGGACCTGAG
GTCGAGCTGCGGTTCTCTGCGGCTCACCGTGTCTATCGAGGCTCGGGCAGC
AGGGCCCTAGTAGCAGCGCATACAGGAGCTGGGGAAGGGGGCTTTGGGG
CCTGCCCACGCGACAGGTAGGGGCGGAGGGGAGCTGAGTCTCCGAAGCTT
GGCTTT
SEQ ID NO: 106

## . Hyaluronan and proteoglycan link protein 4

>gi|30794471|ref|NM\_023002.1| Homo sapiens hyaluronan and proteoglycan link protein 4 (HAPLN4), mRNA

CGGGGCCGCGCGCAAGATGGTGTGCGCTCGGGCGCCCTCGG TCCCGCCGCTCTGGGCCGCGCCTGGGGCGTCCTGCTCACAGCCC CTGCGGGGGCGCAGCGTGCCGGAAGAAGGTCGTGCACGTGCTGGAGGG TGAGTCGGCTCGGTAGTGGTACAGACAGCGCCTGGGCAGGTGGTAAGCC ACCGTGGTGGCACCATCGTCTTGCCCTGCCGCTACCACTATGAGGCAGCC GCCCACGGTCACGACGCGTCCGGCTCAAGTGGACAAAGGTGGTGGACCC GCTGGCCTTCACCGACGTCTTCGTGGCACTAGGCCCCCAGCACCGGGCATT CGGCAGCTACCGTGGGCGGCTGAGCTGCAGGGCGACGGGCCTGGGGAT GCCTCCCTGGTCCTCCGCAACGTCACGCTGCAAGACTACGGGCGCTATGA GTGCGAAGTCACCAATGAGCTGGAAGATGACGCTGGCATGGTCAAGCTGG ACCTGGAAGGCGTGGTCTTTCCCTACCACCCCCGTGGAGGCCGATACAAG CTGACCTTCGCGGAGCGCAGCGCGCGCGCGCGCGAGCAGGACGGCATCCT GGCATCTGCAGAACAGCTGCACGCGGCCTGGCCGCGACGGCCTGGACTGGT GCAACGCGGCTGGTTGCGCGACGGCTCAGTGCAATACCCCGTGAACCGG CCCCGGGAGCCCTGCGGCGCCTGGGGGGGACCGGGAGTGCAGGGGGCG GCGGTGATGCCAACGGGGCCTGCGCAACTACGGGTATCGCCATAACGCC GAGGAACGCTACGACGCCTTCTGCTTCACGTCCAACCTGCCGGGGCGCGT GTGTGCTGCGCGTGGCGGCCGTGGCCAAGGTGGGGCAGCTGTTCGCCG GGCAGTGCGCGCTACCCCATCGTGAACCCGCGAGCGCGCTGCGGAGGCCG CAGGCCTGGTGTGCGCAGCCTCGGCTTCCCGGACGCCACCCGACGGCTCT TCGGCGTCTACTGCTACCGCGCTCCAGGAGCACCGGACCCGGCACCTGGC GGCTGGGGCTGGGCGGCGGCGGCGGCTGGGCAGGGGGCGCGC GCGATCCTGCTGCCTGGACCCCTCTGCACGTCTAGGCTGGAGTAGGCGG ACAGCCAGGCGCTTGACCACTGGTCTAGAGCCCTGTGGTCCCCTGGAGC CTGGCCACGCCTTGAAGCCCTGGACACTGGCCACATTCCCTGTGGTCCCT TACAAACTAACTGTGCCCCTGGGGTCCCTGAAGACTGGCTAGTCCTGGCA GAACAGTACTTTGGAGTTCCCTGGAGCCTGGCCAGCCCTCACCTCTTCTGG ATAGAGGATTCCCCAACTCCCCAACTTTCTCCATGAGGGTCACGCCCCCT GAGGACCTCAGGAGGCCAGCAGAACCCGCAGGCTCCTGAAGACTGGCCA CGCCTCCTGAGACCACTTGGAAACAGACCAACTGCCCCGTGGTCGCCTG GTGGCTGGACCCCGGGATTGACTAGAGACCGGCCGTACACCTTCTGCAT

CTCACTGGAGACTGAACACTAGTCCCTTGCGGTCACGTGGGACACTGGGC GCCTCCTCCCCCCCCCCCCCCGGGGGGGGGGGGACTTCAGGG TCACTCCCGTGGTCACATGGAGGTTGTGGGCCGAGGCGCTTATTTCCCT TATGGTGACCTGAGTCCTGGAGACTCCCATTCTCCCCCCTCTCCCTGAGAGT CCCCTGCAGTTTCTGGGTAACAGGGCACACCCCTCTAGTTTCATGGGCGAG CACCCCATCTGCCACCTCAGACTGACACACAGCCAGCTGGCTCACTTACT GGGGCCACGTCCACCCCTCAGATATTTCTTTGAAGGGAGAGCAAACCC ACCCTGTCCTCTGACGTCCCTTTCCCAACTGTCACCAAACAGACCATCTTC CCAGGCCTGGGGACCGGTAAGATCCATGTCACTAGTTATGCAGAGCAGTT GCCTTGGGTCCCACTGTCACCAAGGCAACCAGTCCTGCTGCTACCTGTCAC CTAGAGTCACACCCCTTCCCTCATCAGGCACACCCATGAAGACAGTGC CTCCCTCCAGCTGTAACCATGGATACCACACATTTCTCATCTCATTGG CCCCACCCAGAGACCTCCACCTCAACTTCTGGCTGTCCCTACCCTGACT CACCGCCATGGAGATCACCCTCCCGAAGCTGTCGCCAGGGTGACCCAAC ATCCAGTTCTCCGGCTCTCACCATGGAAACAAACTGTCCCTGTCCCCAGGC CCACTCCAGTTCCAGACCACCCTCCATGCTCCACCCCCAGGCGGTTTGGAC CCCACCACTGTTGCCATGGTGACCAAACTCTGGAGTCCGAGGTAACAGAA CACCTGTCCCCTAGGCTTTTCCTTGTGGACAACGGGGCCCTGTTCACCAA GCTGTTGCCATAGAGACTGTCAACGTTGTCCTCATGACAACCAGACTTCCA GTTCTCAGGAACTTCTCATTGTGGGCCAGAAGTCCTGGGTGCCTCCTACTA GGGCTACCCTACTGCACCCCATCAGGGGCCTGATGGCTGCCCCTTCCCCAG ACAGGCTGGACTTCTGGAGCTGCTAAGCCACCCTCCGTTTGCACGTTAAC TCTATGCCGGATAGCAGCTGTGCACGAGACAATCTTGCAACACCCGGGCA TGTTTGTCGTCCTACAAATGAGGAAACCGAGCCTATGGCGTGCCCTG GTCTGTTGAGATATGCAAGCACTGAGCTCCTCTTTTGTCCTCTGAGACCCC CCCTCCTTAGAGATCCAGGAGGGATGGAATGTTCTTTAAAATTCAACACC CACCAGGCTCTAAGCGGCGATCTGTGCTAAGAGGTCAGGACCCAGCCGAA GTCCTCGGCGTTGACAGGCAGCTGGGGGGACATGATCCATGGACAAGGCC ATCCCGGCCGTGGGAGACCCCAGTCCCGAAGTCTTGCCTGCAGGAGTACT GGGGTCCCCTGGGGCCCTCTTTACTGTCACGTCATCTCTAGGAAACCTAT CTCTGAGTTTTGGGACCAGGTCGGTTTGGGTTTGAATTCTGCCTCTTCTTGC TCACTGTGTGACCAAGTGACAAACTCCTTCTGAACCTGTGTTCTCCCACTG TACCAGGGCTGTTCTGTGGTCCCCGTGAGTGCCAAGCATACAGTAGGGGC **TCAATAAATCCTTGT SEQ ID NO: 107** 

#### **Immunohistochemistry**

8uM frozen sections were cut from tissue blocks and mounted onto APES slides. The tissue was then fixed in acetone for 10 minutes before being air-dried. The slides were then soaked in 0.3% hydrogen peroxide in methanol for 10 minutes and washed in phosphate-buffered saline (PBS). Non-specific binding sites were blocked by incubating the slides in 20% serum from the appropriate animal and washing again in PBS. Primary antibody diluted in PBS containing 1% serum was then added to the slides. After incubation for 1 hour, the slides were again washed in PBS before incubating with the secondary antibody for a further 1 hour. After final washing in

PBS, the secondary antibody was detected with diaminobenzidine tetrahydrochloride dissolved in Tris buffered saline (TBS), before being washed in TBS and water. The slides were then counter stained in haemotoxylin and viewed under a light microscope.

In certain embodiments, gastric tumors can be localized in situ using stains based on cancer markers of this invention. At least one marker may be forming amyloid structures that can be visualized using Congo red or equivalent, non-specific amyloid stains.

# Tests for Gastric Cancer Markers in Body Fluids

In several embodiments, assays for GTM can be desirably carried out on samples obtained from blood, plasma, serum, peritoneal fluid obtained for example using peritoneal washes, or other body fluids, such as urine, lymph, cerebrospinal fluid, gastric fluid or stool samples.

In general, methods for assaying for oligonucleotides, proteins and peptides in these fluids are known in the art. Detection of oligonucleotides can be carried out using hybridization methods such as Northern blots, Southern blots or microarray methods, or qPCR. Methods for detecting proteins include such as enzyme linked immunosorbent assays (ELISA), protein chips having antibodies, suspension beads radioimmunoassay (RIA), Western blotting and lectin binding. However, for purposes of illustration, fluid levels of a GTM can be quantified using a sandwich-type enzyme-linked immunosorbent assay (ELISA). For plasma assays, a 5 uL aliquot of a properly diluted sample or serially diluted standard GTM and 75 uL of peroxidase-conjugated anti-human GTM antibody are added to wells of a microtiter plate. After a 30 minute incubation period at 30°C, the wells are washed with 0.05% Tween 20 in phosphate-buffered saline (PBS) to remove unbound antibody. Bound complexes of GTM and anti-GTM antibody are then incubated with o-phenylendiamine containing H<sub>2</sub>O<sub>2</sub> for 15 minutes at 30°C. The reaction is stopped by adding 1 M H<sub>2</sub>SO<sub>4</sub>, and the absorbance at 492 nm is measured with a microtiter plate reader.

It can be appreciated that anti-GTM antibodies can be monoclonal antibodies or polyclonal antisera. It can also be appreciated that any other body fluid can be suitably studied.

Certain markers are known to be present in plasma or serum. These include osteopontin (Hotte et al., Cancer 95(3): 507-510 (2002)), prostate-specific antigen

(Martin et al., Prostate Cancer Prostatic Dis. (March 9, 2004) (Pub Med No: PMID: 15007379), thyroglobulin (Hall et al., Laryngoscope 113(1):77-81 (2003); Mazzaferri et al., J. Clin. Endocrinol. Metab. 88(4):1433-14421 (2003), matrix metalloproteinase-2 and -9 (Kuo et al., Clin. Chem. Acta. 294(1-2):157-168 (2000), CEA and TIMP1 (Pellegrini et al., Cancer Immunol. Immunother. 49(7):388-394 (2000). Thus, because some of the above markers are also useful markers for GTM, plasma, serum or other fluid assays are already available for their detection and quantification. Because many proteins are either (1) secreted by cells, (2) sloughed from cell membranes, or (3) are lost from cells upon cell death, other GTM are also present in body fluids, such as plasma, serum and the like. Therefore, in embodiments of this invention, detection of GTM in conveniently obtained samples will be useful and desirable and can be a basis for diagnosis of gastric cancer.

## Western Analysis

Proteins were extracted from gastric tissue using a TriReagent and guanidine HCl extraction method. The non-aqueous phase from the TriReagent extraction of RNA was mixed with 1.5vols of ethanol and centrifuged to remove DNA and OCT medium. 0.5mls of supernatant was mixed with 0.75ml isopropanol, incubated at room temperature for 10 minutes, and then centrifuged. The pellet was washed three times in 1ml 0.3M guanidine HCl in 95% ethanol and once in ethanol alone, then resuspended in 50ul 1% SDS.

Proteins were quantified and electrophoresed on SDS polyacrylamide gels using standard methods. Briefly, the separated proteins were transferred to PVDF membrane using the BioRad trans-blot electrophoretic transfer cell using standard methodology. The membranes were then blocked with a solution containing non-fat milk powder for 30 minutes before being incubated with primary antibody for 2 hours at room temperature. After washing, the membrane was incubated with secondary antibody for 1 hour at room temperature. After final washes, bound antibody was visualized using the ECL detection system (Amersham Biosciences).

Detection of markers in the serum can be accomplished by providing a sample of serum using known methods and then subjecting the serum sample to analysis, either using oligonucleotide probes or antibodies directed against the protein of interest. Immunoblotting, including Western blotting analysis can be especially useful to determine whether alternatively expressed proteins are present in the serum.

Additionally, other body fluids may contain markers, and include peritoneal fluid, cerebrospinal fluid and the like. It is not necessary for a marker to be secreted, in a physiological sense, to be useful. Rather, any mechanism by which a marker protein or gene enters the serum can be effective in producing a detectable, quantifiable level of the marker. Thus, normal secretion of soluble proteins from cells, sloughing of membrane proteins from plasma membranes, secretion of alternatively spliced forms of mRNA or proteins expressed therefrom, cell death (either apoptotic) can produce sufficient levels of the marker to be useful. There is increasing support for the use of serum markers as tools to diagnose and/or evaluate efficacy of therapy for a variety of cancer types.

Yoshikawa et al., (Cancer Letters, 151: 81-86 (2000) describes tissue inhibitor of matrix metalloproteinase-1 in plasma of patients with gastric cancer.

Rudland et al., (Cancer Research 62: 3417-3427 (2002) describes osteopontin as a metastasis associated protein in human breast cancer.

Buckhaults et al., (Cancer Research 61:6996-7001 (2002) describes certain secreted and cell surface genes expressed in colorectal tumors.

Kim et al., (JAMA 287(13):1671-1679 (2002) describes osteopontin as a potential diagnostic biomarker for ovarian cancer.

Hotte et al., (AJ. American Cancer Society 95(3):507-512 (2002) describes plasma osteopontin as a protein detectable in human body fluids and is associated with certain malignancies.

Martin et al., (Prostate Cancer Prostatic Dis. March 9, 2004 (PMID: 15007379) (Abstract) described use of human kallikrein 2, prostate-specific antigen (PSA) and free PSA as markers for detection of prostate cancer.

Hall et al (Laryngoscope 113(1):77-81 (2003) (PMID: 12679418) (Abstract) described predictive value of serum thyroglobulin in thyroid cancer.

Mazzaferri et al., (J. Clin. Endocrinol. Metab. 88(4):1433-1441 (2003) (Abstract) describes thyroglobulin as a potential monitoring method for patients with thyroid carcinoma.

Whitley et al, (Clin. Lab. Med. 24(1):29-47 (2004) (Abstract) describes thyroglobulin as a serum marker for thyroid carcinoma.

Kuo et al (Clin. Chim. Acta. 294(1-2):157-168 (2000) (Abstract) describes serum matrix metalloproteinase-2 and -9 in HCF- and HBV-infected patients.

Koopman et al., (Cancer Epidemiol. Biomarkers Prev 13(3):487-491 (2004) (Abstract) describes osteopontin as a biomarker for pancreatic adenocarcinoma.

Pellegrini et al., (Cancer Immunol. Immunother. 49(7):388-394 (2000) (Abstract) describes measurement of soluble carcinoembryonic antigen and TIMP1 as markers for pre-invasive colorectal cancer.

Thus, we have identified numerous genes and/or proteins that are useful for developing reagents, devices and kits for detecting and evaluating gastric cancer. One or more markers of gastric can be used, either singly or in combination to provide a reliable molecular test for gastric cancer.

#### **EXAMPLES**

The examples described herein are for purposes of illustrating embodiments of the invention. Other embodiments, methods and types of analyses are within the scope of persons of ordinary skill in the molecular diagnostic arts and need not be described in detail hereon. Other embodiments within the scope of the art are considered to be part of this invention.

#### Example 1: Identification of Markers for Gastric Malignancy

Figure 2 depicts a table that shows results of studies using 38 markers for gastric malignancy selected using the above criteria. The Figure 2 includes the symbol for the gene ("symbol"), the MWG oligo number, the NCBI mRNA reference sequence number, the protein reference sequence number, the fold change between tumor and non-tumor gene expression, the fold change rank relative to other genes in the microarray analysis, the results of an original, unadjusted Student's t-test, the results of the Bonferroni-adjusted p value and the results of the 2-sample Wilcoxon test.

The median fold change (tumor: non malignant tissue) for these 34 genes ranged from 1.6 to 7 and the median change in fold change rank ranged from -16,995 to -25,783. The maximum possible change in fold change rank was -29,718. For each of the markers shown, the statistical significance of their specificity as cancer markers was found to be extremely high. The Bonferroni-adjusted p values were, in general, all below 10<sup>-6</sup> or less, indicating that diagnosis using these markers is very highly associated with gastric cancer.

The three cystatins (CST1, CST2, and CST4) are highly homologous and represented by the same oligonucleotide on the microarray and unless otherwise stated, are referred to collectively as "CST1,2,4."

All proteins depicted in Figure 2 were predicted to have signal peptides using the SMART package (European Molecular Biology Laboratory). The signal peptides are known to target synthesized proteins to the extracellular compartment and can therefore be secreted into the interstitial fluid, from which they can have access to the blood. In fact, some proteins of this invention have been detected in serum.

Each of the genes depicted in Figure 2 exhibited a change in intensity rank greater than the two oligonucleotides on the array corresponding to CEA, the marker most frequently used in clinical practice to monitor gastric cancer progression.

### Example 2: qPCR Analysis

More sensitive and accurate quantitation of gene expression was obtained for a subset of the genes shown in Figure 3 using qPCR. RNA from 46 tumor and 49 non-malignant samples was analyzed for 23 genes identified by the microarray analysis (Figure 2) and results are shown in Figure 3. Figure 3 includes the gene symbol, median fold change between cancer and normal tissue, and the % of tumor samples with expression levels greater than the 95<sup>th</sup> percentile of expression levels in non-malignant samples. 12 tumor samples and 9 normal samples were excluded from the analysis because of high (>75%) normal cell contamination, a high degree of necrosis (>40%), or poor hybridization signal on the microarrays. The median fold change (tumor tissues compared to the median non-malignant tissue expression) for these 23 genes ranged from 3 to 525 fold (Figure 3).

The level of expression of genes ASPN, CST1,2,4, LOXL2, TIMP1, SPP1, SFRP4, INHBA, THBS2 and SPARC was greater in tumors than the 95<sup>th</sup> percentile of the non-malignant range for ≥0% of cases (Figure 3). For the remainder of genes, the expression in tumors was greater than the 95<sup>th</sup> percentile in >50% of samples. Each tumor over-expressed at least seven genes greater than the 95<sup>th</sup> percentile indicating that combinations of markers will lead to comprehensive coverage of all gastric tumors.

## Example 3: Validation of Array Data Using qPCR

Array data was validated using quantitative, real-time PCR (qPCR) on the tumor and non-malignant samples with probes for 24 genes. Of all 24 genes studied, 20 showed a strong correlation between the two techniques. Four of these analyses are show in Figures 4a – 4d, which depict graphs of the relative expression for the 4 selected cancer markers detected using array and qPCR methods. For each graph in Figure 4, the horizontal axis represents the array log2 fold change in gene expression, and the vertical axis represents the qPCR log2 fold change in gene expression. We found that there was a strong correlation between the two methods, as indicated by the co-variant relationship between the methods. The strong correlation indicates that both microarray fold change analysis and qPCR are suitable methods for detecting changes in the expression of gastric cancer marker genes and therefore can be used as an accurate, sensitive screening method. It can also be appreciated from Figures 4a – 4d that qPCR can be more sensitive at detecting changes in expression than are array methods. Thus, in situations in which early detection is especially desirable, qPCR may be especially useful.

Figures 5a – 5w depict histograms comparing frequency of observation of expression of each of a series of 23 genes (vertical axis) and the log2 fold change in expression for that gene (horizontal axis), for both normal tissue (open bars) and tumor tissues (black bars). We found surprisingly that for each of these 23 genes, there was substantial separation in the frequency distributions between normal and tumor tissue, as reflected by the low degree of overlap between the frequency distribution curves. For example, Figure 5b depicts the results for CST 1, 2, 4, for which there was only one normal sample observed to have an expression level in the tumor range. In other cases (e.g., Figure 5n; for PRS11) each frequency distribution curve was relatively narrow and there was a degree of overlap. However, even for this marker, the median log2 fold change showed a substantial separation of the amount of gene expression. In other cases, (e.g., Figure 5a; ASPN), although there was some overlap, there was a clear separation of the median log2 fold expression between normal and tumor samples.

Figure 6 depicts a histogram of the number of genes exhibiting a significantly increased expression ("over-expression") in tumor samples compared to normal samples (vertical axis) and the individual samples tested. In each case, the tumor sample exhibited multiple genes with elevated expression levels. The lowest number

of genes having increased expression was 7, found in sample E123. This finding indicates that, in situations in which multiple genes are over-expressed relative to normal tissue, the reliability of cancer detection can be very high, making diagnosis of cancer more certain. However, in some cases, elevation of expression of a single marker gene is sufficient to lead to the diagnosis of cancer.

Our previous comparison with the serum marker most frequently used currently for detection of gastric cancer, CEA, was based on difference in intensity rank of array data between tumors and normal samples. This comparison was verified using qPCR data for the markers and CEA.

Figures 7a-7c depict graphs of the relative log2 expression (compared to a reference RNA preparation) of markers in individual tumor samples and non-malignant samples compared to the expression of the gene for the tumor marker, CEA. CEA is the serum marker currently most used to monitor progression of gastric cancer. The zero point is defined to be the median normal expression for each marker. It can be seen that there is extensive overlap between the expression of the CEA gene (CEACAM5) in tumor samples and normal samples. This overlap is markedly less in the gastric cancer markers ASPN, CSPG2, CST1,2,4, IGFBP7, INHBA, LOXL2, LUM, SFRP4, SPARC, SPP1, THBS2, TIMP1, adlican, LEPRE1, and EFEMP2. For the other markers in Figures 7b-7c, ASAH1, SFRP2, GGH, MMP12, KLK10, TG, PRSS11 and TGFBI, the overlap between the tumor expression range and the non-malignant tissue expression range is greater than the overlap for the above markers, but still less than that of CEA, indicating that all of the herein described new markers are quantitatively better than CEA, and therefore can provide more reliable diagnosis.

To minimize effects of variable tissue handling, tumor:normal (non-malignant) fold changes were calculated using qPCR data from tumor and non-malignant tissue samples derived from the same patient. Such paired analysis corrects for differences in background levels of gene expression in different individuals and minimizes the effects of tissue handling on RNA quality. For example, if the resected stomach was at room temperature for an hour, the transcripts from the normal and tumor samples will be degraded to the same extent.

Figure 8 summarizes the T:N expression levels determined by qPCR for the markers, but used paired data (i.e., tumor and non-malignant samples) from the same individual. Figure 8 also includes expression data for six genes that were not included in Figure 3. The additionally studied genes are MMP2, CGR11, TGFB1, PCSK5,

SERPINB5, and SERPINH1. Identifying information and probes are shown in Figures 1 and 2. Figure 8 shows the median T:N fold change and the maximum T:N fold change for 29 gastric cancer markers in these 40 patients with "paired" samples. 27 of the 29 markers have a median T:N difference greater than or equal to the prior art marker, CEA. 29/29 of the markers have a higher percentage of paired samples in which the expression in the tumor sample exceeds the expression in the normal sample.

Figures 9a – 9d depict scatter dot plots of data from tumor and normal tissue from the same individuals. Each point represents the fold-change, within patient, in expression of the markers in tumor tissue relative to the expression in non-malignant tissue. All of the markers studied have better discrimination of tumor from non-tumor tissue than CEA. Three markers, CST1,2,4, ASPN and SFRP4 showed 100% discrimination between the paired tumor and normal samples. That is, for those markers, every tumor tissue had greater expression than did the corresponding non-tumor tissue from the same individual. In many other markers, for example, Adlican, CSPG2, EFEMP2, IGFBP7, INHBA, LOXL2, LUM, SERPINH1, SPARC, SPP1, TGFbI, THBS2 and TIMP1, each had only 2 or 3 individual points for which tumor tissue expression was less than that of the non-tumor tissue. Thus, for those markers, the likelihood that any one pair of tumor and non-tumor tissue would produce a false negative is relatively low (e.g., 3 of 40 or 7.5%; 2 of 40 or 5%, 1 of 40 or 2.5%). Thus, even if the other markers listed immediately above were used, use of multiple samples from an individual patient would produce reliable diagnostic information.

The gene sequences of these markers, and the location of the primers and probes used to detect them, are shown herein above.

To determine if over-expression of the marker genes is independent of the stage of the gastric tumors, the paired T:N log2 fold changes were plotted against the tumor stage (Figures 10a – 10ad). No stage dependency of expression on tumor stage was observed for 26 of the markers listed in Figure 8. These markers were similarly over-expressed in early stage as well as late stage tumors. However, KLK10 showed more consistent over-expression in stage 1 and stage 2 tumors, and PCSK5 and SERPINB5 showed more consistent over-expression in stage 4 tumors. KLK10, PCSK5 and SERPINB5 therefore can be used in determining the stage of gastric tumors.

In a similar analysis, paired T:N log2 fold changes were plotted against the Lauren classification of the tumor (either diffuse type or intestinal type). Figures 11a – 11ad show that each of the 29 GTMs discriminated between tumor and non-tumor tissue, regardless of whether the type of tumor was intestinal (I) or diffuse (D).

## Example 4: Use of Multiple Markers

As described above, certain markers exhibit an ability to discriminate tumor from non-tumor tissue in 100% of the samples. Other markers, also described above, can be used in combination to achieve very high degrees of discrimination of tumor tissue from non-tumor tissue. Figure 12 depicts a 3-dimensional plot of the expression of 3 markers, SERPINH1, CST1,2,4 and INHBA, expressed as log2 T:N fold changes for a series of gastric tumor samples and non-malignant gastric samples. There is complete separation between the two groups of samples.

The reliability of successful discrimination of tumor and non-tumor samples using marker combinations is further illustrated by a statistical analysis summarized in Figure 13. This analysis compared the normal distributions of data generated using the qPCR gene expression from paired tumor and non-malignant samples, shows the effect of increasing the numbers of markers used to discriminate between tumor and non-malignant samples on test sensitivity (with a fixed specificity of 95%). Although few of the 29 markers (as shown in Figure 8) have a sensitivity of greater than 90, 95, or 99% when used alone in this analysis, the combination of two or three markers enabled high sensitivity to be reached with large numbers of combinations. For example, 50 combinations of three markers would discriminate between tumor and non-malignant samples with a sensitivity of ≥9% and specificity of ≥5%.

# Example 5: Detection of Gastric Tumor Marker Proteins

In yet further embodiments, GTM proteins can be detected as a basis for diagnosis. In certain situations, the concentration of mRNA in a particular sample, such as a sample containing no cells, it may be difficult to use either microarray or qPCR methods to detect elevations in gene expression. Thus, in certain embodiments, detection of GTM proteins can be accomplished using antibodies directed against either the entire protein, a fragment of the protein (peptide) or the protein core. Methods for detecting and quantifying expression of proteins and peptides are known in the art and can include methods relying on specific antibodies raised against the

protein or peptide. Monoclonal antibodies and polyclonal antisera can be made using methods that are well known in the art and need not be described herein further.

To demonstrate that GTM proteins can be used to discriminate tumor from non-tumor tissue, commercial antibodies were obtained against SPARC (R&D Systems; cat # AF941), THBS2 (Santa Cruz Biotechnology Inc; cat # sc-7655), CSPG2 (Calbiochem; cat # 428060) and IGFBP7 (R&D Systems; cat # AF1334). An additional polyclonal antibody was raised in rabbits (Alpha Diagnostic International Inc; San Antonio) against the cystatin SN peptide sequence 50-66 (C) FAISEYNKATKDDYYRR.

SEQ ID NO: 108.

These antibodies were used in either immunohistochemistry or Western analysis of tumor and non-malignant gastric tissue. Each of these markers showed strong tumor:normal differences at the protein level. This confirmed that the over-expression observed at the RNA level for these genes also occurred at the protein level.

Figure 14 shows a Western blot analyses of total protein extracted from two pairs of tumor and non-malignant tissues using antibodies against the proteins encoded by SPARC, CST1 (cystatin SN), IGFBP7 and THBS2. For each marker, the signal is significantly higher in the tumor samples than the non-malignant samples.

The antibody raised against cystatin SN detected three major bands, corresponding to molecular weights of approximately 34, 45 and 65kDa respectively. The lowest molecular weight band is shown in Figure 14. The protein species were larger than the control cystatin SN protein, suggesting that the protein produced by tumors has undergone post-translational modifications or multimerization. Regardless of the mechanism responsible for the differences in molecular weights of CST proteins, Figure 14 demonstrated that CST expression was low in the non-tumor tissue, but was easily observed in blots of tumor-derived proteins.

Figure 14 also showed that SPARC protein is expressed substantially to a greater degree in tumor tissue than in non-tumor tissue. The SPARC protein had gel mobility slower than the form of this protein that was detected in serum (Figure 15), also indicating the occurrence of different post-translational modifications in proteins produced by malignant gastric cells. Regardless of the mechanism(s) responsible for any such modification, the finding that SPARC is over-expressed in tumor tissue relative to non-malignant tissue indicates that SPARC is a useful protein marker.

Similarly, IGFBP7 and THBS2 show over-expression in tumor tissue relative to non-malignant tissue.

Immunohistochemical analysis of tumor and non-malignant tissue was carried out using antibodies against the proteins encoded by CSPG2 (versican) and CST1 (cystatin SN). Immunohistochemical analysis of tissue with antibodies against versican identified strong staining in the extracellular matrix of tumor tissue, but not non-malignant tissue. With the anti-cystatin SN antibodies, strong staining was observed in the area around the outside of the tumor cells. In non-malignant cells, the staining with this antibody was weaker, and observed only on the mucosal surface of the tissue and the lining of the gastric pits. This demonstrated that in non-malignant cells, cystatin SN protein is directed out of the cell onto the mucosal surface and not into the extracellular spaces. Therefore, not only is the cystatin SN protein being produced in higher amounts in tumor tissue than non-malignant tissue, but, unlike the protein produced by the non-malignant tissue, the tumor cystatin SN is in direct contact with the tissue vasculature. To extend these observations, cystatin SN was immunoprecipitated from the supernatant of the gastric cancer cell line, AGS with a monoclonal antibody (R&D Systems; cat # MAB1285) (Figure 16). Large amounts of cystatin SN were detected in the supernatant, confirming that this protein is produced by, and secreted from, gastric epithelial cells.

#### Example 6: Analysis of Tumor Markers in Serum

For a marker to be useful for rapid screening, it is desirable for the marker to be present in the serum in sufficient levels for detection. Certain proteins described in Figure 8 can be secreted into the blood at detectable levels from gastric cancers. One marker known to be secreted from gastric tumors into blood in detectable levels is TIMP1. However, if a protein is secreted or shed from any surface of a cell other than a mucosal surface, it will have contact with the interstitial fluid. From there, it can pass either directly into the blood supply through a capillary or via the lymph system. Thus, any shed GTM will be present in blood. Osteopontin, thyroglobulin, and members of the MMP and kallikrein families have previously been described to be elevated in the serum of patients with a range of epithelial cancers, but not gastric cancer. TIMP1 has, however, previously been observed to be elevated in the serum of gastric cancer patients. These findings suggest that the selection criteria for markers in this study, namely over-expression of secreted proteins in tumor tissue but not non-

malignant tissue, can be effectively used to detect markers in the serum, and thus can be of substantial use clinically, without the need for tissue or organ biopsies.

From Figure 15, it is apparent that the serum SPARC has a different molecular weight (depicted here in the Western blot) with the tumor SPARC having a lower molecular weight than the SPARC produced by blood cells. Thus, even though SPARC is produced by tumor and non-tumor blood cells, the presence of tumor SPARC can be determined using molecular size, such as determined using Western analysis, or with an antibody specific for the glycosylated protein produced by the tumor cells.

In another study, we detected cystatin SN in the supernatant of a gastric cancer cell line, AGS. Figure 16 depicts a Western analysis of media alone or a supernatant from AGS cells in culture. The right hand lane of Figure 16 shows a dense band corresponding to cystatin SN protein.

Thus, we conclude from Figure 10 that GTM of this invention are suitable for diagnosing gastric cancers at early, middle or late stages of progression of the disease.

Although certain marker proteins can be glycosylated, variations in the pattern of glycosylation can, in certain circumstances, lead to mis-detection of forms of GTMs that lack usual glycosylation patterns. Thus, in certain embodiments of this invention, GTM immunogens can include deglycosylated GTM or deglycosylated GTM fragments. Deglycosylation can be accomplished using one or more glycosidases known in the art. Alternatively, GTM cDNA can be expressed in glycosylation-deficient cell lines, such as prokaryotic cell lines, including E. coli, thereby producing non-glycosylated proteins or peptides. It can also be appreciated that the level and quality of glycosylation can be sensitive to the presence of essential precursors for sugar side-chains. Thus, in the absence of an essential sugar, "normal" glycosylation may not occur, but rather, shorter or missing side chain sugars may be found. Such "glycosylation variants" can be used as immunogens to produce antibodies specific for different types of marker genes.

Additionally, certain GTMs may form homo-or heterodimers or other types of multimeric forms. For example, inhibin beta A is a 47 kDa protein that can form homodimers of 97 kDa molecular weight (activin A) and 92 kDa heterodimers with the 45 kDa protein inhibin beta B (the heterodimers are known as activin AB). Thus, it can be appreciated that Western analysis or other type of assay that provides molecular weight need not be limited to only detection of a monomeric form of a

GTM. Rather, one can readily appreciate that any form of a GTM can be detected, regardless of the molecular weight. Thus, detection of a multimeric form of a GTM can be readily used to diagnose the presence of gastric cancer. Further, for those GTM that are selective for stage (1-4) or type of gastric tumor (diffuse or intestinal), detection of a multimeric form can provide suitable target for evaluating stage or type of gastric cancer.

Once an antibody or antiserum against a GTM is produced, such antibody preparations can be used for in a variety of ways. First, enzyme-linked immunosorbent assay (ELISA) or radioimmunoassay (RIA) methods can be used to quantify GTM proteins or peptides. Immunodetection can be accomplished in tissue samples using immunohistochemistry. These methods are all known in the art and need not be described further herein.

#### **Example 7: Vectors Containing GTM Oligonucleotides**

Other embodiments of this invention include vectors useful for in vitro expression of marker genes or portions thereof ("marker peptides") or fragments of marker gene products. For example, vectors can be made having oligonucleotides for encoding GTMs therein. Many such vectors can be based on standard vectors known in the art. This invention also includes vectors that can be used to transfect a variety of cell lines to prepare GTM-producing cell lines, which can be used to produce desired quantities of GTMs for development of specific antibodies or other reagents for detection of GTMs or for standardizing developed assays for GTMs.

It is to be understood that to manufacture such vectors, an oligonucleotide containing the entire open reading frame or a portion of such an open reading frame encoding a portion of the protein to be expressed can be inserted into a vector containing a promoter region, one or more enhancer regions operably linked to the oligonucleotide sequence, with an initiation codon, an open reading frame, and a stop codon. Methods for producing expression vectors are known in the art and need not be repeated herein.

It can also be appreciated that one or more selectable markers can be inserted into an expression vector to permit the expansion of cell lines selected to contain the expression vector of interest. Moreover, one can also insert leader sequences known in the art, in frame, to direct secretion, internal storage or membrane insertion of the protein or protein fragment in the expressing cell.

## Example 8: Cells Transfected with GTM-Containing Vectors

In still further embodiments, cells are provided that can express GTMs, GTM fragments or peptide markers. Both prokaryotic and eukaryotic cells can be so used. For example, E. coli (a prokaryotic cell) can be use to produce large quantities of GTMs lacking in mature glycosylation (if the particular GTM normally is glycosylated). COS cells, 293 cells and a variety of other eukaryotic cells can be used to produce GTMs that are glycosylated, or have proper folding and therefore, three-dimensional structure of the native form of the GTM protein. Methods for transfecting such cells are known in the art and need not be described further herein.

### Example 9: Kits

Based on the discoveries of this invention, several types of test kits can be produced. First, kits can be made that have a detection device pre-loaded with a detection molecule (or "capture reagent"). In embodiments for detection of GTM mRNA, such devices can comprise a substrate (e.g., glass, silicon, quartz, metal, etc) on which oligonucleotides as capture reagents that hybridize with the mRNA to be detected. In some embodiments, direct detection of mRNA can be accomplished by hybridizing mRNA (labeled with cy3, cy5, radiolabel or other label) to the oligonucleotides on the substrate. In other embodiments, detection of mRNA can be accomplished by first making complementary DNA (cDNA) to the desired mRNA. Then, labeled cDNA can be hybridized to the oligonucleotides on the substrate and detected.

Regardless of the detection method employed, comparison of test GTM expression with a standard measure of expression is desirable. For example, RNA expression can be standardized to total cellular DNA, to expression of constitutively expressed RNAs (for example, ribosomal RNA) or to other relatively constant markers.

Antibodies can also be used in kits as capture reagents. In some embodiments, a substrate (e.g., a multiwell plate) can have a specific GTM capture reagent attached thereto. In some embodiments, a kit can have a blocking reagent included. Blocking reagents can be used to reduce non-specific binding. For example, non-specific oligonucleotide binding can be reduced using excess DNA from any convenient source that does not contain GTM oligonucleotides, such as salmon sperm DNA.

Non-specific antibody binding can be reduced using an excess of a blocking protein such as serum albumin. It can be appreciated that numerous methods for detecting oligonucleotides and proteins are known in the art, and any strategy that can specifically detect GTM associated molecules can be used and be considered within the scope of this invention.

In embodiments relying upon antibody detection, GTM proteins or peptides can be expressed on a per cell basis, or on the basis of total cellular, tissue, or fluid protein, fluid volume, tissue mass (weight). Additionally, GTM in serum can be expressed on the basis of a relatively high-abundance serum protein such as albumin.

In addition to a substrate, a test kit can comprise capture reagents (such as probes), washing solutions (e.g., SSC, other salts, buffers, detergents and the like), as well as detection moieties (e.g., cy3, cy5, radiolabels, and the like). Kits can also include instructions for use and a package.

Although this invention is described with reference to specific embodiments thereof, it can be appreciated that other embodiments involving the use of the disclosed markers can be used without departing from the scope of this invention.

#### INDUSTRIAL APPLICABILITY

Methods for detecting GTM family members include detection of nucleic acids using microarray and/or real time PCR methods and detection of proteins and peptides. The compositions and methods of this invention are useful in the manufacture of diagnostic devices and kits, diagnosis of disease, evaluating efficacy of therapy, and for producing reagents suitable for measuring expression of GTM family members in biological samples.